

## UTILIZATION OF POST HARVEST TECHNOLOGIES AMONG YAM FARMERS IN SELECTED LOCAL GOVERNMENT AREAS OF NIGER STATE, NIGERIA

Hassan<sup>1\*</sup>, S., Salihu<sup>2</sup>, I. T., Abdullahi<sup>2</sup>, A., and Aliyu<sup>1</sup>, A.

<sup>1</sup>Department of Agricultural Economics and Extension Technology,  
Federal University of Technology, Minna, P.M.B. 65, Niger State, Nigeria

<sup>2</sup>Department of Agricultural Extension and Rural Development  
Federal University of Technology, Minna, P.M.B. 65, Niger State, Nigeria

\*Corresponding Author's E-mail: [organotim31@gmail.com](mailto:organotim31@gmail.com), GSM: [07061683781](tel:07061683781)

### ABSTRACT

*Yam is an important tuber crop in Nigeria. However, its availability is affected by rate of deterioration arising from poor postharvest handling. Hence, farmers use various techniques to minimize the loss. Therefore, this study examined the utilization of post harvest technologies among yam farmers in selected local Government areas of Niger state, Nigeria. Data were collected from 195 respondents and analyzed with descriptive statistics. The result revealed that, the mean age of the respondent was 35.0 years and highly experienced in food crops production with mean of 18.0 years experience. Majority of the respondents (79%) had formal education. However data analysis reveals that yam barn storage technique ranked first in terms of types and level of utilization by the farmers with the highest weighted mean (WM=3.87) while, open sided shelve was the least (WM=1.43). It is therefore recommended that extension agents should actively disseminate information on improved storage techniques to yam farmers in the study area as well as the need to use such techniques.*

**KEYWORDS:** Technology, Storage technique, Postharvest losses, Yam barn, Utilization.

### INTRODUCTION

Agriculture is the most important economic sector of any Nation saddled with the responsibility of meeting the food requirements of the rather fast growing global population (Andersen, 2010). The situation is direr with the projected world population expected to reach 10.5 billion by the year 2050 as stated by the United Nation (UN, 2013) thereby, adding to the global food security concerns. This projection, translates into 33% more human mouths to feed, with the greatest demand growth in the poor communities of the world. Food supplies would therefore need to increase by 60% as estimated at 2005 food production levels in order to meet the food demand in 2050 (Alexandratos and Bruinsma, 2012). To this end, food availability and accessibility can be increased by increasing production, improving distribution and reducing the losses by Food Agricultural Organization (FAO, 2010). Thus, reduction of post-harvest food losses is a critical component of ensuring future global food security.

Post harvest handling and storage of yam is an important aspect of economic development in Nigeria. Accurate financial worth of yam industry in Nigeria is hard to come by, but it was estimated at two hundred billion naira (200b1) per annum with estimated losses from storage over half the expected revenue (FAO, 2010). These losses were mostly attributed to rot caused by bacteria, fungi and nematode. The bulk of

these pathogens causing yam tuber rots are soil borne but manifestation of the tuber disease are observable mostly during storage (Okigbo, 2004).

Similarly, weight loss during storage in traditional or clamp storage can reach 10-12% in the first three months and 30-60% after 6 months of storage and in West Africa alone; this amounted to an annual loss of one million tons of tuber (FAO, 2003). It is important to also point out that maximization of profit can only be achieved when farmers are well equipped with required technological knowledge and skills of yam storage. This act is necessary because one of the major constraints upon establishing effective storage and processing management approaches for smallholder farmers is the lack of adequate information sources to increase farmers' knowledge level in the practices of yam tuber processing and storage. In fact, the need to understand farmers' knowledge systems has been recognized as a basis for development of proper storage and processing technologies that are sustainable and adaptable to local farmer's environmental condition.

However, despite the necessity of producing more food to meet the ever-increasing global population, most of agricultural produce are lost during postharvest handling and that up to 50-70% losses are estimated between production area and consumption point because of lack of storage facilities, limited access to processing technologies, unstable market prices, poor

market opportunities and non utilization of postharvest technologies (Olayemiet *al.*, 2012) and (Owolade, 2011). Post-harvest losses will lead to reduction in farmer's income, food insecurity, poor nutritional value and lack of input for the next production (Olayemiet *al.*, 2012).

In recent times research on post harvest preservation has been given little support by both the government and the private sectors. To minimize post-harvest losses, improved methods of storage have therefore been developed. However, the poor economic status of most Nigerian farmers has inhibited the adoption and usage of most agricultural technologies (Alimi and Zango, 2016). It was hoped that farmers' usage of these technologies would lead to reduction in food losses, improved income and enhanced food security (Okiedo and Onemolease, 2009). Nevertheless, the widespread and continued use of traditional storage practices by small-scale farmers despite considerable losses usually associated with these methods need investigation.

Over the past decades, significant focus by researchers (such as: Gbemisola, 2009; shehu, 2010) and resources have been allocated to increasing food production. For instance, 95% of the research investments during the past 30 years were reported to have focused on increasing productivity, and only about 5% was directed towards reducing losses (Kader, 2005; Kader and Roller, 2004). Although, increasing agricultural productivity is critical for ensuring global food security but this may not be sufficient. The global food supply is currently being challenged by limited post-harvest preservation techniques as a result of a wide gap that exist between actual achievement and achievable potential in the yam farming industry. This study seeks to fill the identified research gap. In view of the above, this study is initiated to assess the utilization of post harvest technologies among yam farmers in selected areas of Niger state, Nigeria. The specific objectives are to:

- i. describe the socio economic characteristics of yam farmers
- ii. identify the types of post harvest technologies used by the yam farmers
- iii. examine the level of utilization of post harvest technologies by yam farmers
- iv. identify the constraints limiting the utilization of post harvest technologies by yam farmers in the study area.

## METHODOLOGY

The study was conducted in Niger State of Nigeria. Niger State is located between latitudes 8°11'2 N and

11° 20'2 N and longitude 4° 30'2 E and 7° 20'2 E (Ojoet *al.*, 2013). The State is located in the North central zone along the Middle Belt region of Nigeria. The state has a population of about 3,954,772 individuals as stated by the National Population Commission (NPC, 2006). Niger state is classified as one of the largest states in the country spanning over 86,000km<sup>2</sup> in land area with 80% of the land mass conducive for agriculture (Tologbonse, 2008). With 9.30% of the total land area of the country, Niger state is not only divided into three agricultural zones under climatic features containing nearly all classes of soils of the savannah regions of West Africa (Tologbonse, 2008). In order to achieve the study objectives, multiple sampling techniques were employed. This involves purposive selection of zone B of Niger state because of the preponderance of yam production in this zone. The second stage involved the purposive selection of three Local Government Areas known for higher yam production in the zone namely; Paikoro, Shiroro, and Bosso Local Government Areas. The third stage involved the random selection of two districts from the selected Local Government Areas. The fourth stage involved the random selection of two villages from each of the selected Local Government Areas. The fifth stage involved a proportionate random selection of 5% from the registered yam farmers in each of the selected villages. Therefore, a total of 195 yam farmers were considered as the sample size for this study. Interview schedule was used to elicit data from the respondents. The responses were analyzed using frequency counts, percentages and mean score to describe the socio economic characteristics and the types of post harvest technologies used by yam farmers. However a 4-point Likert type of scale was employed for objective (iii) and scored as follows: Always Used (AU) = 4, Frequently Used (FU) = 3, Sometimes Used (SU) = 2, Never Used (NU) = 1. A mean score of 2.5 was obtained and the decision rule is that any mean ( ) scores  $\geq 2.5$  indicate used, while scores  $< 2.5$  not used. Similarly a 3-point Likert type of scale was employed for objective (iv) and scored as follows: Very Serious (VS) = 3, Serious (S) = 2, and Not Serious (NS) = 1. The mean score was obtained and the decision rule is that any mean ( ) scores  $\geq 2.0$ , indicate serious constraint while scores  $< 2.0$  indicate Not serious constraint.

## RESULTS AND DISCUSSION

### Socio economic characteristics of the respondents

The results in Table 4.1 showed that majority (92.4%) of the respondents were between the age of 21 and 50 years, while only 3.1% were less than 20 years and 4.6% above 50 years. The mean age of the respondents was 35 years. The implication of the mean age on utilization of post-harvest technologies is that the young farmers can take risk by utilizing new technologies than the older farmers. This supports the

findings of Jabil and Abdu (2012) who stated that young farmers are willing to use new technologies than the older ones and they are not averse to risk. Similarly, (69.2%) of the farmers had been into yam production for over 20 years and all the respondents had an average farming experience of 18 years. This implies that the farmers are quite experienced in yam cultivation. This compares favorably with the findings of Falola *et al.*, (2017); Oluwatosin (2011) that indicated that the yam farmers had a significant level of expertise in yam production. Similarly, more than half (59.0%) of the respondents had a household size of about 3-6 members, while the mean of the household size was found to be five (5), implying a large household size. This is in agreement with Girohet *et al.* (2012) who reported that farmers with large household size tend to channel more of their income to food consumption expenditure rather than to save and invest in improved storage technique. The result further revealed that majority (79%) of the respondents had formal education. This implies that respondents are educated enough to know and understand the complexities involved in improved technology to adopt it. This assertion is in contrast to Tor *et al.* (2017); Onemolease (2005), who reported that a low educational background not exceeding primary education may impede acceptance of improved post harvest technologies, since education facilitates farmers' utilization of innovations.

#### **Types of post harvest technologies used by yam farmers**

Post harvest technologies are referred to as technologies commonly used for the purpose of storing and processing by the yam farmers in the study area. The comprehensive list of technologies was developed and farmers were asked to indicate the technologies they use. Tables 2 showed that majority of the respondents use the following post harvest technologies: Storage in barn (98.5%), Processing into pounded yam (97.4%), Processing into fried yam (96.9%) and Storage in mud hut (88.7%). While open sided shelf stores (32.3%) and ventilated pit (28.7%) are the least storage methods used. This result agrees with that of Akangbe *et al.* (2012) who stated that storage of yam tubers in barns was the major storage method utilized while ventilated pit and open sided shelf stores were the least methods used by the respondents in the study area.

#### **Level of utilization of post harvest technologies by yam farmers**

Table 3 indicates that storage of yam tubers in barns was the major storage method used by the respondents in the study area (mean = 3.87). However, in terms of frequency of utilization, (91.3%) of the respondents always use yam barns, followed by processing into

pounded yam (90.2%) with the mean of 3.85, then processing into fried yam (77.9) with (mean = 3.68). Storage in open sided shelf stores were the least used storage techniques with (mean = 1.43). This implies that majority of the respondents depend on manual or traditional technology to execute some vital processes. This result agrees with that of Falola *et al.* (2017); Okoedo and Onemolease (2009) who stated that storage of yam tubers in barns was the major storage method utilized by the respondents in the study area. It also tallies with the findings of Akangbe *et al.*, (2012) who reported open sided store was the least adopted method by farmers in the storage of yam in Asa Local Government Area of Kwara State.

Summarily, level of utilization of post harvest technologies in the study area as shown in table 4 revealed that utilization level is low (51.3%) . This result may be attributed to lack of access and effective information on these improved technologies. Thus, respondents rely on traditional methods for yam storage and processing.

#### **Constraints limiting the utilization of post harvest technologies by yam farmers**

Analysis on table 5 revealed some of the constraints limiting the utilization of post harvest technologies. The result showed that the incidence of pest and diseases and lack of improved technology ranked first and third. This may be because of the quality deterioration of yam as a result of poor storage strategy till the time of sales. Most farmers in the study area use traditional method of storing yam. This result collaborated with the findings of Abubakar and Nasiru (2017) who stated that the yam barn are locally made or constructed which give room to micro-organisms and rodent to destroy yam tubers stored. High cost of labour has been identified to limit effective use of the local storage methods especially the storage barn. This is corroborated by Nwaigwe *et al.*, (2015) who reported that the construction of barn requires a lot of work and effort and is more expensive than other local storage methods. This finding agrees with that of Olayemiet *et al.*, (2011) who stated that bruising and spoilage, high cost of transportation, inadequate storage facilities and menace of theft are responsible for losses incurred by farmers and marketers. However the result seems to disagree with the research of Okoedo and onemolease (2009) that did not recognize theft of tubers as a significant cause of postharvest losses in

yam production.

### **CONCLUSION AND RECOMMENDATIONS**

The study examined the utilization of post harvest technologies among yam farmers in selected local government areas of Niger state, Nigeria. Based on the findings of this study, it was concluded that respondents use local post harvest technologies and utilization of modern postharvest technologies is low despite the losses due to the use of traditional techniques. It is therefore recommended that extension agents should actively disseminate information on improved post harvest to yam farmers in the study area as well as the need to use such technologies. New yam postharvest technologies should be made available to the farmers in the study area at subsidized rates so as to encourage them to use such technologies.



**Table 1: Socio-economic characteristic of respondents**

<b>Variable</b>	<b>Frequency</b>	<b>Per</b>
<b>Age</b>		
below 21 years	6	3.1
21-30 years	68	34.9
31-40 years	53	27.2
41-50 years	59	30.3
51-60 years	9	4.6
<b>Farming experience</b>		
below 11 years	46	23.6
11-20 years	89	45.6
21-30 years	42	21.5
31-40 years	15	7.7
41-50 years	3	1.5
<b>Household size</b>		
below 3 members	41	21.0
3-6 members	115	59.0

above 6  
members

20.0

**Level of  
education**



non formal 41  
Primary 43  
Secondary 58  
Tertiary 51  
Masters 2

21.0  
22.3  
29.7  
26.2  
1.0

Source: ICAAT, 2018

**Table 2 Types of post harvest technologies used by yam farmers**  
**Technologies Frequency**

Storage in barn 192 98.5

Storage in pit 115 59.0

Curing method 82 42.1

Storage in mud hut 173 88.7

Storage in open sided shelf stores 63 32.3

Storage in elevated store shed 70 35.9

---

Storage in ventilated pit	56	28.7
Storage in thatched roof pit	74	37.9
Processing into flour	181	92.8
Processing into chips	123	63.1

Processing into fried yam	189	96.9
Processing into pounded yam	190	97.4

Source: ICAAT, 2018

**Table 3 level of utilization of post harvest technologies by yam farmers**

Technologies	AU	FU	SU	NU	WM	Rank
Storage in barn	178(91.3)	12(6.2)	2(1.0)	3(1.5)	3.87	1 <sup>st</sup>
Processing into pounded yam	176(90.2)	13(6.7)	1(0.5)	5(2.6)	3.85	2 <sup>nd</sup>
Processing into fried yam	152(77.9)	30(15.4)	7(3.6)	6(3.1)	3.68	3 <sup>rd</sup>
Processing into flour	130(66.7)	44(22.6)	7(3.6)	14(7.2)	3.49	4 <sup>th</sup>
Storage in mud hut	110(56.4)	49(25.1)	14(7.2)	22(11.3)	3.27	5 <sup>th</sup>
Processing into chips	19(9.7)	30(15.4)	74(37.9)	72(36.9)	1.96	6 <sup>th</sup>
Storage in pit	13(6.7)	36(18.5)	66(33.8)	80(41.0)	1.91	7 <sup>th</sup>
Storage in thatched roof pit	17(8.7)	19(9.7)	38(19.5)	121(62.1)	1.66	8 <sup>th</sup>
Curing method	6(3.1)	15(7.7)	61(31.3)	113(57.9)	1.56	9 <sup>th</sup>
Storage in elevated store shed	5(2.6)	10(5.1)	55(28.2)	125(64.1)	1.46	10 <sup>th</sup>
Storage in ventilated pit	4(2.1)	23(11.8)	29(14.9)	139(71.3)	1.45	11 <sup>th</sup>
Storage in open sided shelf stores	5(2.6)	10(5.1)	48(24.6)	132(67.7)	1.43	12 <sup>th</sup>

Source: ICAAT, 2018

**Note:** Always Used (AU); Frequently Used (FU); Sometimes Used (SU); Never Used (NU) WM=Weighted Mean

**Table 4 categorization of respondents' level of utilization of post harvest technologies**

Utilization level	Frequency	Percentage
Low utilization (0 - 0.33)	100	51.3
Moderate utilization (0.34 - 0.66)	63	32.3
High utilization (0.67 - 1.00)	32	16.4
<b>Total</b>	<b>195</b>	<b>100.0</b>

Source: ICAAT, 2018

**Table 5 constraint faced by yam farmers**

Constraints	VS	S	NS	WM	Rank
Pests and diseases	161(82.6)	30(15.4)	4(2.1)	2.81	1 <sup>st</sup>
Poor transport network	146(74.9)	49(25.1)	0(0)	2.75	2 <sup>nd</sup>
Lack of improved technology	150(76.9)	33(16.9)	12(6.2)	2.71	3 <sup>rd</sup>
Long distance to market	140(71.8)	49(25.1)	6(3.1)	2.69	4 <sup>th</sup>
Insufficient working capital	141(72.3)	47(24.1)	7(3.6)	2.69	4 <sup>th</sup>
High cost of transportation	141(72.3)	41(21.0)	13(6.7)	2.66	6 <sup>th</sup>
Lack of credit	123(63.1)	53(27.2)	19(9.7)	2.58	7 <sup>th</sup>
High cost of labour	121(62.1)	66(33.8)	8(4.1)	2.58	7 <sup>th</sup>
Theft of yam	121(62.1)	65(33.3)	9(4.6)	2.57	9 <sup>th</sup>
Low government support	116(59.5)	58(29.7)	21(10.8)	2.49	10 <sup>th</sup>
Limited land	94(48.2)	67(34.4)	34(17.4)	2.31	11 <sup>th</sup>
Labor unavailability	83(42.6)	85(43.6)	27(13.8)	2.29	12 <sup>th</sup>
Injury on yam	64(32.8)	104(53.3)	27(13.8)	2.19	13 <sup>th</sup>
Poor storage facilities	60(30.8)	109(55.9)	26(13.3)	2.17	14 <sup>th</sup>
Illiteracy	65(33.3)	93(47.7)	37(19.0)	2.14	15 <sup>th</sup>
Lack of storage/ Processing facilities	65(33.3)	88(45.1)	42(21.5)	2.12	16 <sup>th</sup>
Poor management skill	68(34.9)	77(39.5)	50(25.6)	2.09	17 <sup>th</sup>
Poor buyers	50(25.6)	83(42.6)	62(31.8)	1.94	18 <sup>th</sup>
Over storage	40(20.5)	98(50.3)	57(29.2)	1.91	19 <sup>th</sup>
Lack of extension contact	59(30.3)	59(30.3)	77(39.5)	1.91	19 <sup>th</sup>

Source: ICAAT, 2018

**Note:** VS-Very Serious; S-Serious; NS-Not Serious; WM=Weighted Mean**REFERENCES**

Abubakar, M. and Nasiru, M. I. (2017). Assessment of post-harvest losses of yam in selected district of Karu local government area, Nasarawa state. *FUW Trends in Science & Technology Journal*, retrieved from [www.ftstjournal.com](http://www.ftstjournal.com).

Akangbe, J.A., Oloruntoba, O. O., Ayanda, I.F. and Komolafe, S. E. (2012). An Analysis of Yam Storage Strategy to Promote Food Security. *Ethiopian Journal of Environmental Studies and Management* 5 (4): 550-558

- Akinbile, L. A. (2007). Standardization of socioeconomic status (SES) scale for farm family in south-west Nigeria. *Journal of social sciences*, 14(3): 221-227.
- Alexandratos, N. and J. Bruinsma. (2012). World agriculture towards 2030/2050: the saving water. From Field to Fork-Curbing Losses and Wastage in the Food Chain 2012 revision. Working paper: FAO: ESA No. 12-03, p.4.
- Alimi, H. M. and Zango, M. I. (2016) The Influence of Socio-Economic Variables On Adoption Behaviour Towards Tadco Improved Rice Parboiling Technique Among Rice Parboilers In Kura Processing Areas Of Kano State, Nigeria. *Agrosearch*. 16 (2): 51-60
- Andersen, P., (2010). The State of Food Security in Pakistan: Future Challenges and Coping Strategies, 903–923.
- Falola, A., Salami, M. F., Bello, A. A. and Olaoye T. A. (2017). Effect of yam storage techniques usage on farm income in kwara state, Nigeria. *Agrosearch (2017)* 17(1): 54– 65.
- FAO. (2003), Food and Agricultural Organization 2003. Data bases. F.A.O. report (1998).
- Food and Agricultural Organization. (2010). Nigeria Food Security report 2010. Food Security Statistics, Nigeria (FAOSTAT) [www.fao.org](http://www.fao.org).
- Gbemisola, O. and Paul, W. (2009). Rural nonfarm activities and agricultural crop production in Nigeria. *Wiley Blackwell journal*, 40(2): 189-201
- Giroh, D. Y., Gal, T. N. and Minampah, C. J. (2012). Analysis of the Determinants of Savings among Gum Arabic Collectors in Selected Local Government Areas of Adamawa State, Nigeria. *New York Science Journal*, 5(11): 1-6
- Olayemi, F. F., Adegbola, J. A., Bamishaiye, E. I. of horticulture crops in Nigeria. *Journal and Awagu, E. F.* (2011) Assessment of *Horticulture*. 1 (9): 14-16. postharvest losses of some selected crops in eight local government areas of Rivers Shehu, J. F., Iyortyer, I. T., Mshelia, S. I. and Jongur, state Nigeria, *Asian Journal of Rural A. A. U.* (2010). Determinants of yam
- Jabil, I. Y and Abdu, U. D. (2012). Socio-economic characteristics of farmers and the factors that hampers their Adoption of Agricultural Technologies in Northern Central Zones of Plateau State, Nigeria. *International journal of agriculture economics and extension*, 4 (1): 5.
- Kader, A. A. (2005). “Increasing food availability by reducing postharvest losses of fresh produce.” *Acta Horticulture* 68(2): 21692176.
- Kader, A. A. and R. S. Rolle. 2004. The Role of Postharvest Management in Assuring the Quality and Safety Horticultural Crops. Food and Agriculture Organization. *Agricultural Services Bulletin* 15(2): 52 .
- National Population Commission, NPC (2006). Nigeria Population Census Report, NPC Abuja.
- Nwaigwe, K. N., Okafor, V. C., Asonye, G. U. and Nwokocho, J. C. (2015). *Analysis of Tuber Storage Techniques in Africa: A Review*. ASABE Annual International Meeting, July 26-29, New Orleans, Louisiana.
- Ojo, M. A., Nmadu, J. N., Tanko, L. and Olaleye, R. S. (2013). Multinomial Logit Analysis of Factors Affecting the Choice of Enterprise among Small-holder Yam and Cassava Farmers in Niger State, Nigeria. *journal of agricultural science*, 4(1): 7-12
- Okigbo, R. N. (2004). A review of biological control methods for postharvest yams *Dioscoreaspp.* In storage in South Eastern Nigeria. *King Mongkut's institute of technology Ladkrabangscience journal*. 4(1): 207 - 215.
- Okoedo-Okojie, D. U. and Onemolease, E. A. (2009). Factors Affecting the Adoption of Yam Storage Technologies in the Northern Ecological Zone of Edo State, Nigeria. *Journal of Human Ecology*. 27(2): 155160.

- Development*, 20(11): 1-11. production and technical efficiency among yam farmers in Benue State
- Oluwatosin F. M. (2011): Measuring Technical Efficiency of Yam Farmers in Nigeria: A Stochastic Parametric Approach. *Agricultural Journal* 6 (2): 40-46.
- Onemolease, E. A (2005). Imp act of the Agricultural Development Programme (ADP) Activities in Arable Crop Production on Rural Poverty Alleviation in Edo State, Nigeria. Ph.D Thesis (Unpublished). University of Benin, Benin City, Edo State, Nigeria pp123
- Owolade, S. O. (2011). Promotion of food Security through reduction in post-harvest losses Nigeria. *Journal of Social Science*, 24(2): 143-148.
- Tolgonse, D. (2008). Policy Issues In Meeting Rice Farmers Agricultural Information Needs in Niger State. *Journal of Agricultural Extension*. 12(2): 84-94.
- Tor, I. E., Iheanacho, A. C. and Okeke, A. M. (2017). Analysis of Socio – economic Challenges of Using Local Storage Systems for Root and Tuber Crops in Benue State, Nigeria. *International Journal of Innovative Agriculture & Biology Research* 5(4): 30-37.

