



Journal of

PHARMACY & BIORESOURCES

VOL. 2 NO 1

MARCH 2005

- 
- Medicinal plant research
 - Herbal medicines and cosmetics;
 - Development of drugs & Pharmaceuticals
 - Quality assurance of drugs
 - Safety and efficacy of drugs
 - Pharmacy practice
 - Veterinary pharmacy
 - Bioresources development & Conservation



EDITORIAL NOTICE

The Journal of Pharmacy and Bioresources (JPB) is published by the Faculty of Pharmaceutical Sciences, University of Jos, Jos. It covers scientific research work in all areas of Pharmaceutical and life sciences, including (but not restricted to): medicinal plant research; herbal medicines and cosmetics; development of drugs and pharmaceuticals; quality assurance of drugs; safety and efficacy of drugs; pharmacy practice; veterinary pharmacy; bioresources development and conservation. JPB is published in bi-annual editions (March and September). The Journal is available on the internet through the African Journal Online (AJOL) website. Please visit the journal at <http://www.ajol.info/journals/jpb> Abstracts of published articles are also available at <http://www.groups.yahoo.com/group/jpharmbiores> where you can join our yahoo group.

To this end, the publishers of JPB will be glad to receive articles for publication in the journal. Such articles must report original research work, which has not been published elsewhere and is not concurrently submitted elsewhere for publication. All manuscripts will be subjected to peer review process and those found to be of high quality would be published. Copyright of articles published shall reside exclusively with the Faculty of Pharmaceutical Sciences, University of Jos, Nigeria.

EDITORIAL BOARD

| | |
|-----------------------------|---|
| <i>Editor-in-Chief:</i> | Prof. E. N. Sokomba |
| <i>Managing Editor:</i> | Dr. N. A. Ochekepe |
| <i>Press Editor:</i> | Dr. T. O. E. Alemika |
| <i>Editorial Secretary:</i> | Mrs. J. D. Audu-Peter |
| <i>Members:</i> | Prof. T. A. Iranloye Prof. F. K. Okwuasaba |
| | Prof. K. I. Ekpenyong Prof. C. O. Akueshi |
| | Dr. M. A. Ibrahim Dr. J. A. Kolawole |
| | Dr. F. C. Ohiri Mr. P. O. Olorunfemi. |

PROMOTING ACCESS TO AFRICAN RESEARCH

Custom Search

AFRICAN JOURNALS ONLINE (AJOL)

JOURNALS

ADVANCED SEARCH

USING AJOL

RESOURCES

Journal of Pharmacy & Bioresources - Vol 2, No 2 (2005)

Journal Home > Archives > Vol 2, No 2 (2005)

Log in or Register to get access to full text downloads.

Username

Password

 Remember me

Login

or Register

JOURNAL HOME

ABOUT THIS JOURNAL

ADVANCED SEARCH

CURRENT ISSUE

ARCHIVES

 Open Access Subscription or Fee Access

Table of Contents

Articles

Comparative analgesic effects of paracetamol with paracetamol-caffeine formulation

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Abdulkadir U Zezi, Ibrahim Abdu-Aguye, Hamza A Yaro, Bello B Maiha, Nuhu M Danjuma

65-69

<http://dx.doi.org/10.4314/ajob.v2i2.32054>Effects of hexane extract of *Ricinus communis* L. (var. *minor*) on blood parameters in rats

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Rosaleen McNeil, Ekwere Ekwere, Sylvanus Dafur, Francis Okwuasaba

70-74

<http://dx.doi.org/10.4314/ajob.v2i2.32055>Preliminary studies of some pharmacological properties of *Croton zambesicus*

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Jude Okokon, Nnoye Nwajuaku, Aniekan Udokpoh

75-79

<http://dx.doi.org/10.4314/ajob.v2i2.32056>Evaluation of anti-snake venom activity of the aqueous root extract of *Securidaca longipedunculata* in rats

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Noel N Wannang, Athassan M Wudil, Maxwell LP Dapar, Lawal A Bichi

80-83

<http://dx.doi.org/10.4314/ajob.v2i2.32057>Hypotensive effect of *Baobab axillans* (Apocynaceae) leaves in normotensive rabbits

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Buniyamin A Ayinde, Fabian C Amaechina

84-88

<http://dx.doi.org/10.4314/ajob.v2i2.32058>Phytotherapeutic potentials and biochemical study of *Nyssa fructicans* (Wumb)

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Chukwuma SI Odoemena, Benjamin AJ Ekpo

89-92

<http://dx.doi.org/10.4314/ajob.v2i2.32059>Some *in vivo* and *in vitro* studies of the aqueous leaf extract of *Phyllanthus muellerianus* (Euphorbiaceae) in laboratory animals

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Joseph A Anuka, Ahmed H Yaro, Noel N Wannang, Ephraim B Ezenwanne, Ibrahim A Yakasai

93-99

<http://dx.doi.org/10.4314/ajob.v2i2.32060>Phytochemical analysis and inhibitory effect of *Pyrenacantha staudtii* leaf extract on isolated rat uterus

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Abiodun Falodun, Cyril O Usifoh, Zuleikha AM Nworgu

100-103

<http://dx.doi.org/10.4314/ajob.v2i2.32061>Preliminary evaluation for anti-trypanosomal activity of aqueous stem bark extract of *Crossopteryx febrifuga* in *Trypanosoma congolense* - infected rats

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Aminu B Yusuf, Binta Ilyasu, Abdulkadir Abubakar, Nnenna A Onyekwele, Augustine C Igweh, Florence N Ojiegbu, David Y Bot

111-115

<http://dx.doi.org/10.4314/ajob.v2i2.32062>

Phytochemical and antimicrobial screening of three Nigerian medicinal plants used to treat infectious diseases traditionally

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Philip A Idowu, Jones O Moody, Herbert A Odelola

116-119

<http://dx.doi.org/10.4314/ajob.v2i2.32063>

Antimicrobial activities of some volatile oils against some pathogenic organisms

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Ijeoma Ahonkhai, Buniyamin A Ayinde

120-123

<http://dx.doi.org/10.4314/ajob.v2i2.32064>Studies on the antimicrobial properties of formulated creams and ointments containing *Baphia nitida* heartwood extract

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Adeniji K Olowosulu, Yakubu KE Ibrahim, Partap G Bhatia

124-130

<http://dx.doi.org/10.4314/ajob.v2i2.32065>Antibacterial activity of the stem bark of *Boswellia dalzielii*

EMAIL FULL TEXT

DOWNLOAD FULL TEXT

Moses A Olukemi, Yvonne T Kandakar-Olukemi, John D Mawak

131-136



HOW TO USE AJOL...

- for Researchers
- for Librarians
- for Authors

OTHER RESOURCES...

- for Researchers
- for Journals
- for Authors
- for Policy Makers
- about Open Access
- FAQ's
- News
- AJOL jobs
- More about AJOL
- AJOL's Partners
- Contact AJOL
- Terms and Conditions of Use

CURRENT ISSUE



521 African Journals

- Browse By Category
- Browse Alphabetically
- Browse By Country
- List All Titles
- Open Access Titles

Featuring journals from 32 Countries:

- Algeria (5)
- Benin (2)
- Botswana (3)
- Burkina Faso (3)
- Cameroon (8)
- Congo, Republic (1)
- Côte d'Ivoire (4)
- Egypt, Arab Rep. (14)
- Eritrea (1)
- Ethiopia (30)
- Ghana (27)
- Kenya (29)
- Lesotho (1)
- Libya (2)
- Madagascar (1)
- Malawi (4)
- Mauritius (3)
- Mozambique (1)
- Nigeria (221)
- Rwanda (3)
- Senegal (6)
- Sierra Leone (1)
- South Africa (96)
- South Sudan (1)
- Sudan (3)
- Swaziland (3)
- Tanzania (19)
- Togo (1)
- Tunisia (2)
- Uganda (12)
- Zambia (2)
- Zimbabwe (12)

Like & GK

Share

<http://dx.doi.org/10.4314/jpb.v2i2.32076>

Brine shrimp toxicity of acidic fractions of *Boswellia daizileli* gum resin

[EMAIL FULL TEXT](#)

[DOWNLOAD FULL TEXT](#)

Taiwo E Alemika, Stephen O Ojerinde

137-140

<http://dx.doi.org/10.4314/jpb.v2i2.32077>

Effect of pH and ionic strength on the bioadhesive properties of *Prosopis africana* gum

[EMAIL FULL TEXT](#)

[DOWNLOAD FULL TEXT](#)

Anthony A Attama, Petra O Nnamani, Ogbonna Okorie

141-145

<http://dx.doi.org/10.4314/jpb.v2i2.32078>

4-Carboxyl-2, 6-dinitrobenzene diazonium ion (CDNBD), a new diazonium for the detection of phenol ether homologues

[EMAIL FULL TEXT](#)

[DOWNLOAD FULL TEXT](#)

Olajire A Adegoke, Olakunle S Idowu, Monsurat O Lawal, Ajibola A Olaniyi

146-161

<http://dx.doi.org/10.4314/jpb.v2i2.32079>

Synthesis and characterization of some novel azetidinone derivatives as anti-bacterial and anti-convulsant agents

[EMAIL FULL TEXT](#)

[DOWNLOAD FULL TEXT](#)

Aiyalu Rajasekaran, Sankaranarayanan Murugesan

162-168

<http://dx.doi.org/10.4314/jpb.v2i2.32080>

Prevalence and prognosis of cerebrovascular accident in Zaria, Kaduna State, Nigeria

[EMAIL FULL TEXT](#)

[DOWNLOAD FULL TEXT](#)

Nuhu M Danjuma, Abdulkadir U Zezi, Ibrahim Abdu-Aguye

169-173

<http://dx.doi.org/10.4314/jpb.v2i2.32081>

ISSN: 0189-8442

Abdul - Abur NTR



Vol. 2 no. 2, pp. 47-51 (September 2005)

<http://www.ajol.info/journals/jpb>

Journal of
**PHARMACY AND
BIORESOURCES**

Preliminary evaluation for anti-trypanosomal activity of aqueous stem bark extract of *Crossopteryx febrifuga* in *Trypanosoma congolense* - infected rats

Aminu B. Yusuf^{1*}, Binta Iliyasu, Abdulkadir Abubakar, Nnenna A. Onyekwelu, Augustine C. Igweh, Florence N. Ojiegbu and David Y. Bot

Biochemistry and Chemotherapy Division, Nigerian Institute for Trypanosomiasis Research (NITR), P.M B. 03, Vom, Plateau State, Nigeria.

Received 14th April 2004; Revised 30th March 2005; Accepted 15th July 2005

Abstract

Aqueous stem bark extract of *Crossopteryx febrifuga* was evaluated for anti-trypanosomal activity in rats. Twenty four rats divided into four groups (A, B, C, D) of six animals per group were used. Infection with *T. congolense* was established after a pre-latent periods of 10 and 11 days for curative (Group B) and prophylaxis (Group A) studies respectively. Both Group A and Group B were treated orally with 1500mg/kg of the crude extract for five consecutive days. Parasitemia, Packed Cell Volume (PCV), Red Blood Cells (RBC) counts and Differential leucocytes (WBC) counts were monitored throughout the studies. Phytochemical analysis of the crude aqueous extract was also carried out. Anaemia developed with establishment of infection in all the groups. Analysis of PCV, RBC and Differential WBC counts showed no significant difference ($P>0.05$) between treated and untreated animals. The crude extract contains saponins, tannins, reducing sugars, volatile oils and alkaloids as phytochemicals. It showed no significant prophylaxis or curative against *T. congolense* infection in rats as traditionally claimed. Although the plant may be used locally in combination with other plants or ingredients or even other orthodox drugs, further work is needed to fully assess the claim of the use of *Crossopteryx febrifuga* stem bark in the management or treatment of animal trypanosomiasis.

Keywords: *Crossopteryx febrifuga*; Trypanosomiasis; *Trypanosoma congolense*.

Introduction

Trypanosomiasis is a disease caused by parasitic protozoans called trypanosomes that cause sleeping sickness in humans and related diseases in domestic animals. Some 35 million people and 25 million cattle in Africa are at risk of infection with trypanosomes (Turner, 1982). Although, chemotherapy and chemo prophylaxis have continued to form

the most important and major aspect of the control and eradication of trypanosomiasis (Igweh and Onabanjo, 1989), the discovery and development of new drugs has been slow and far from satisfactory (Fairlamb, 1989). Resistance against existing trypanocides and toxicity are common occurrences (Onyekwelu, 1999). Thus, the need for new curative agents can not be over emphasized.

* Corresponding author.

Perhaps local medicinal plants could provide such agents with potential remedy against both human and animal trypanosomiasis. One of such plants is *Crossopteryx febrifuga*, claimed to be used by Fulani herdsmen in Northern Nigeria, for the treatment of Animal trypanosomiasis (Okpara and Ali, 2002).

The present studies were therefore carried out to evaluate the potential of *Crossopteryx febrifuga* in the treatment or management of *T. congolense* infected rats.

Experimental

Chemicals reagents. Anhydrous sodium dihydrogen phosphate (NaH_2PO_4) and anhydrous disodium hydrogen phosphate (Na_2HPO_4), sodium chloride, all purchased from British Drug House (BDH). All reagents for phytochemical analysis were of Analar grade.

Collection of plant. The stem bark of *C. febrifuga* (Hausa – Kashin Awaki; Fulani - Rumajegohi) was collected from Kaltungo LGA, Gombe State with the help of traditional medical practitioners. The plant was identified by Mr. Abdulkarim of the College of Forestry, Jos, Plateau State, Nigeria.

Preparation of aqueous extract. The extract was prepared according to the information obtained from the Fulani Herdsmen. The stem bark of *C. febrifuga* was air dried under ambient temperature and reduced to coarse powder using a pestle and mortar. Two hundred grammes (200g) of the powder was boiled in one liter of distilled water for 3 hrs. The aqueous extract obtained was decanted and filtered using muslin cloth and then Whatman's filter paper. The filtrate was carefully and slowly evaporated to dryness using hot water bath at 45°C . The dried extract was kept in a sealed container and refrigerated at 4°C until required.

Phytochemistry. The extract of *C. febrifuga* was tested for the presence of phytochemicals according to the method of Sofowora (1979). The pH was also determined using pH meter (Water proof pH scan 3, Eutech instruments)

Acute toxicity studies. The median lethal dose (LD_{50}) of the extract was determined by the method of Lorke (1983). This method involves two stages; stage one: 9 animals and dosages of 10mg/kg, 100mg/kg, 1000mg/kg are used. In stage two, 3 animals and dosages of 1600mg/kg, 2900mg/kg and 5000mg/kg are used. Thereafter, LD_{50} was calculated.

Experimental animals. Twenty four mature male albino rats were obtained from Parasitology Section, NITR, Vom. The animals were divided into four groups (A, B, C, D) of six animals each and allowed to acclimatize for two weeks. They were fed on Growers' mash (Vital Feeds, Nig. Ltd.) and water *ad libitum*.

Strain of parasite. *Trypanosoma congolense* (Karu strain) was obtained from Veterinary and Livestock Studies Division, NITR, Vom. The parasites were maintained in the laboratory by serial blood passage in donor rats.

Experimental design. Blood was collected from the tail of a heavily infected donor rat and immediately diluted with sterile phosphate saline buffer (pH 7.4). Parasites in the diluted blood which served as the inoculum were counted using a microscope and estimated by the method of Herbert and Lumsden (1976). Group A (prophylaxis) was treated orally with the extract at 1500mg/kg body weight for five days consecutively and then inoculated intraperitoneally with 0.1ml of the inoculum containing 1×10^5 parasites. Group C (prophylaxis control) was given normal saline for five consecutive days. Group B (curative) was inoculated intraperitoneally with 0.1ml of the inoculum containing 0.1ml of the parasite. When the parasites begin to appear in the peripheral

blood which indicated the establishment of infection, treatment commenced with the extract at 1500mg/kg body weight for five days consecutively. Group D (curative control) was given normal saline for five days simultaneously with group B treatment.

Determination of hematological parameters. Packed Cell Volume (PCV), Red Blood cells (RBC) counts and Differential leucocytes (WBC) counts were determined weekly. PCV was by microhaematocrit method, RBC counts using haemocytometer and Differential WBC counts as described by Schalm *et al.* (1975).

Results

Phytochemical analysis. The phytochemical analysis of the crude aqueous extract showed that it contains saponins (saponin glycosides), reducing sugars, tannins, volatile oils and alkaloids. It was also acidic with pH of 5.30.

Acute toxicity studies. The media lethal dose (LD₅₀) for the oral administration of the extract was found to be above 5000mg/kg body weight. No immediate or late sign of toxicity was observed, thus the extract was considered safe (Lorke, 1983).

Prophylaxis studies. After treatment for 5 days and then inoculation with the parasites,

animals were examined daily for parasitemia. Infection developed after a pre-latent period of 11 days in both treated group and untreated control. The animals died from 9-13 and 4-11 days post infection (Pi) for treated and untreated groups respectively. No significant difference ($P > 0.05$) was observed in the RBC count, PCV and Differential WBC counts between treated and untreated groups (Table 1, 3 and 5). However, a slight but insignificant ($P > 0.05$) increase in monocytes was observed in the treated group when compared to the control (Table 5).

Curative studies. Infection developed after a pre-latent period of 10 days in both treated and untreated groups. Treatment for 5 days commenced 10 days post inoculation and did not change the pattern of parasitemia as the parasites continued to multiply until all the animals died. The animals died between 9-11 and 8-13 days post-infection for treated and untreated groups respectively. There was no significant difference ($P > 0.05$) observed in the RBC counts, PCV and Differential WBC counts between the treated animals and untreated control (Table 2, 4 and 6). However, a slight but insignificant increase in lymphocytes was observed after the start of treatment.

Table 1: Prophylactic effect of *C. febrifuga* on RBC of *T. congolense* infected rats.

| Post inoculation days | RBC Count X 10 ¹² /L (mean ± S.D) | | P |
|-----------------------|--|----------|-------|
| | Group A | Group C | |
| 0 | 5.20±0.7 | 5.90±0.2 | >0.05 |
| 5 | 5.79±0.5 | 6.15±0.3 | >0.05 |
| 12 | 4.92±0.4 | 4.63±0.3 | >0.05 |
| 19 | 3.06±0.1 | 2.74±0.1 | >0.05 |

Key: Group A = Treated infected, Group C = Untreated infected control.

Table 2: Curative effect of *C. febrifuga* on RBC of *T. congolense* infected rats.

| Post inoculation days | RBC Count X 10 ¹² /L (mean ± S.D) | | P |
|-----------------------|--|-----------|-------|
| | Group A | Group C | |
| 0 | 5.05±0.20 | 4.98±0.30 | >0.05 |
| 6 | 5.41±0.50 | 5.40±0.30 | >0.05 |
| 13 | 4.39±0.40 | 3.37±0.70 | >0.05 |
| 20 | 5.01±0.00 | | >0.05 |

Key: Group B = Infected treated, Group D = Infected untreated control!

Table 3: Prophylactic effect of *C. febrifuga* on PCV of *T. congolense* infected rats.

| Post inoculation days | RBC Count $\times 10^{12}/l$ (mean \pm S.D) | | p |
|-----------------------|---|----------------|-------|
| | Group A | Group C | |
| 0 | 46.3 \pm 2.5 | 38.0 \pm 4.9 | >0.05 |
| 6 | 37.0 \pm 1.4 | 45.7 \pm 2.1 | >0.05 |
| 13 | 40.3 \pm 3.3 | 45.7 \pm 2.1 | >0.05 |
| 20 | 29.0 \pm 1.0 | 29.0 \pm 1.0 | >0.05 |

Key: Group A = Treated infected, Group C = Untreated infected control

Discussion

Treatment with *C. febrifuga* extract showed no significant effect on the number of parasites as the parasitemia continued to progress after infection was established. The parasitemia was milder in treated animals compared to untreated controls (Table 7). In both prophylaxis and curative studies, the pre-incident periods of treated groups and untreated controls were the same. This shows the absence of prophylaxis against *T. congolense* infection.

The PCV and RBC counts of the treated groups continued to drop till all the animals died as a result of anaemia. Expectedly, the drops in PCV and RBC counts correspond with the periods of progressive increase in parasitemia. The development of anaemia towards the terminal stage of the disease is thought to be as a result of the numerous increases of circulating trypanosomes causing lysis of the erythrocytes by the lashing action of the parasites flagellae (Esiebo *et al.*, 1982). The breakdown of plasma protein, and the activities of the enzymes neuraminidase and phospholipases produced by trypanosomes may have also been involved in this process, since they are reported to be released towards the late stage of infection (Esiebo *et al.*, 1982). The slight monocytosis in prophylaxis group was accompanied by mild neutrophilia, lymphopaenia and eosinopaenia. Monocytosis, lymphopaenia, and eosinophania are all consistent findings in trypanosomiasis (Emembe and Anosa, 1991).

The observed monocytosis is associated with the increased demands to remove particulate matter such as dead trypanosomes and lysed red blood cells from circulation (Nwosu and Ikeme, 1992).

Conclusion

Within the frame of this work, the aqueous stem bark extract of *C. febrifuga* has not shown significant anti-trypanosomal activity in rats infected with *T. congolense*. This seems to be contrary to the claims of the Fulani herdsmen. However, it is likely that the herdsmen use it in combination with other plant ingredients or even orthodox drugs, whereby the action of one plant or ingredient may potentiate that of the other synergistically to produce a desired effect. Thus, caution and care should be employed while seeking information on medicinal plants from local herdsmen or traditional herbalists, because such information obtained may be misguided or incomplete.

Nevertheless, further work will be needed to fully assess the claim of the use of *C. febrifuga* stem bark extract in the management or treatment of animal trypanosomiasis as this is a preliminary investigation.

Acknowledgement

Authors wish to acknowledge the technical assistance of Mr. Larfa Domtur, Miss Elizabeth Dang and Mr. Francis Doro.

References

- Emeribe, A.O. and Anosa, V.O.(1991). Haematology of experimental *T. b. gambiense* infection. II Erythrocyte and leucocyte changes. *Rev. Med. Vet Pays. Trop* 44 (1): 53.
- Esiebo, K.A., Saror, O.I., Hemode, A.A., and Hallaway, M.H. (1982). Variation in erythrocyte surface and free serum sialic acid concentrations during experimental *T. vivax* infection in cattle. *Res. Vet. Sci.* 32: 1-5.
- Fairlamb, A.H. (1989). Future prospects for the chemotherapy of Human trypanosomiasis: I. Novel approaches to the chemotherapy of trypanosomiasis. *Trans. Royal Soc. Trop. Med. Hygiene.* (1990) 84, 613-617.
- Herbert, W.J. and Lumsden, W.H. (1976). *Trypanosoma brucei* : A rapid matching method for estimating the host's parasitemia. *Experimental Parasitology* 40, 427 - 431.
- Igweh, A.C. and Onabanjo A.O. (1989). Chemocurative effects of *Annona senegalensis* in *T. b. brucei*. *Annals of Tropical Medicine and Parasitology.* 83(5), 527 - 534 (1989).
- Lorke, D.(1983). A new approach to acute toxicity testing. *Archives of Toxicology* (1983) 54: 275 - 287.
- Nwosu, C.O. and Ikeme, M.M (1992). Parasitemia and clinical manifestations in *T. brucei* infected dogs. *Rev. Med. Vet Pays. Trop.* 45(3-5): 273 - 277.
- Okpara, J.O. and Ali, G.A. (2002). Studies on ethnoveterinary preparations, treatments, and control of cattle diseases in Kanam LGA, Plateau state (unpublished).
- Onyekwelu, N.A. (1999). A Review. Toxicity of existing trypanocides. *West Afr. J. Pharmacol. Drug, Res.* Vol. 15 (1, 2).
- Schalm O.W., Jain N.C. and Carroll E.J. (1975). Materials and methods for the study of blood in veterinary hematology, Lea Febiger, Philadelphia. 47-50.
- Sofowora, A. (1979). African medicinal Plants. Univ. of Ife Press, Ile-Ife, Nig 400 pp.
- Turner, M.J. (1982). Biochemistry of variant surface glycoproteins of salivarian trypanosomes. *Acta Parasitol.*, 21, 61 - 153.