



# SOIL SCIENCE

SOCIETY OF NIGERIA (SSSN)



## PROCEEDINGS

OF THE

# 43<sup>RD</sup> ANNUAL CONFERENCE

SOIL SCIENCE SOCIETY OF NIGERIA (SSSN)

Date: 15th - 19th July, 2019

FOOD BASKET 2019



Theme:

# UNDERSTANDING NIGERIAN SOILS

FOR SUSTAINABLE FOOD AND NUTRITION  
SECURITY AND HEALTHY ENVIRONMENT

**Host:**

Department of Soil Science  
University of Agriculture, Makurdi

EDITED BY:

Jayeoba, J.O., Idoga, S., Olatunji, O., Jimin, A.A., Adaikwu, A. O.,  
Ibrahim, F. and Anikwe, M.A.N.



**PROCEEDINGS OF THE 43<sup>RD</sup> ANNUAL CONFERENCE  
OF THE SOIL SCIENCE SOCIETY OF NIGERIA**

***THEME:***

**UNDERSTANDING NIGERIAN SOILS FOR SUSTAINABLE  
FOOD AND NUTRITION SECURITY AND HEALTHY  
ENVIRONMENT**

**DEPARTMENT OF SOIL SCIENCE, COLLEGE OF AGRONOMY,  
FEDERAL UNIVERSITY OF AGRICULTURE, MAKURDI,  
BENUE STATE – NIGERIA**

**FOOD BASKET 2019**

**DATE: 15<sup>TH</sup> – 19<sup>TH</sup> JULY, 2019**

***EDITED BY:* Jayeoba, J.O., Idoga, S., Olatunji, O., Jimin, A.A., Adaikwu, A.O.,  
Ibrahim, F. and Anikwe, M.A.N.**

**© 2019 SOIL SCIENCE SOCIETY OF NIGERIA (SSSN)**

**ISBN: 978-978958-981-5**



©2019 NSSS

Published in 2019

SOIL SCIENCE SOCIETY OF NIGERIA (SSSN)  
Publishers of the Nigerian Journal of Soil Science

Jayeoba et al (editors). 2018. Understanding Nigerian Soils for Sustainable Food and Nutrition Security and Healthy Environment. Proceedings of the 43<sup>rd</sup> Annual Conference of the Soil Science Society of Nigeria held at the Department of Soil Science, College of Agronomy, Federal University of Agriculture, Makurdi, Benue State - Nigeria. 15<sup>th</sup> - 19<sup>th</sup> July, 2019.

### CITATION



**NATIONAL EXECUTIVE COMMITTEE OF SSSN (2019)**

Prof. B.A. Raji	President
Prof. D. O. Asawalam	Vice President
Prof. Peter I. Ogban	National Secretary
Prof. M. A. Hassan	Asst. National Secretary
Dr. (Mrs) T.O. Ande	Treasurer
Dr. E. E. Oku	Financial Secretary
Prof. M.A.N Anikwe	Editor-in-Chief
Prof. T. A. Okusami	Chairman, Awards Committee
Prof. V.O. Chude	Ex-Officio
Prof. O.O. Agbede	Ex-Officio
Prof. G. A. Oluwatosin	Ex-Officio

**ADMINISTRATIVE STAFF**

Miss Esther Ogbole	-	Administrative Secretary, SSSN
Miss Acquine Onwukwe	-	Administrative Secretary, NISS

**MEMBERS OF COUNCIL NIGERIA INSTITUTE OF SOIL SCIENCE**

Prof. A. O. Ogunkule	-	President
Prof. G. Nwaka	-	Vice President
Prof. Ibrahim Saminu	-	North East
Prof. S. Noma	-	North West
Prof. A. O. Osunde	-	North Central
Prof. C.L.A. Asadu	-	South East
Prof. T. O. Ibia	-	South South
Prof. V. O. Chude	-	Ex-Officio



## **FELLOWS OF THE SOIL SCIENCE SOCIETY OF NIGERIA**

1. Prof. Akinola A. Agboola
2. Prof. Walter O. Enwezor
3. Prof. Abayomi G. Ojanuga
4. Prof. G. Lombin
5. Prof. Rauf. A. Sobulo (L.)
6. Prof. David O. Ataga
7. Chief A. O. Nnodi (L.)
8. Prof. Eno Jumbo Udo (L.)
9. Otunba G. Daniel
10. Dr. Sam Ominyi Egwu
11. Mallam Adamu Bello
12. Prof. V. O. Chude
13. Prof. Ivara E. Esu (OFR)
14. Prof. A. O. Ogunkunle
15. Prof. A. O. Babalola
16. Prof. T. A. Okusami
17. Prof. S. O. Ojeniyi
18. Prof. Olusola O. Agbede
19. Prof. Patrick O. Aina
20. Prof. A. Olayinka
21. Prof. J. A. I. Omueti
22. Prof. Olu Obi
23. Prof. Emeritus G. O. Obigbesan
24. Chief C. O. Ezendu
25. Prof. B. A. Raji
26. Dr. Jimmy I. Ibanga (L.)
27. Prof. Marian G. Solomon
28. Dr. Umoru Omoti
29. Prof. F. O. R. Akamigbo
30. Prof. Adebayo Adewale
31. Prof. Emmanuel A. Aduayi
32. Prof. Caroline C. Mba
33. Prof. Akim Osa Osunde
34. Dr. (Mrs.) Foluke I. Oluwatoyinbo
35. Prof. M. A. N. Anikwe
36. Prof. Charles L. A. Asadu



37. Prof. Uche C. Amalu
38. Prof. J. A. Adediran
39. Prof. Abayomi S. Fasina
40. Prof. Bashiru H. Usman
41. Dr. Akinwumi Adesina
42. Prof. Abubakar M. Kundiri
43. Dr. Napoleon O. Aisueni
44. Prof. Nnaemeka Isirimah (L.)
45. Prof. G. Lekwa
46. Prof. T. O. Ibia
47. Dr. Bukar Tijani
48. Chief Godswill Akpabio
49. Dr. David Sasseville
50. Prof. J. D. Kwari
51. Prof. Geoffrey Ijoma C. Nwaka
52. Prof. Gabriel Akinboye Oluwatosin
53. Prof. Nnenna Nnannaya Oti
54. Dr. Anthony O. Ano
55. Dr. Emmanuel U. O. Okoye
56. Mr. Ojuola Olabode
57. Prof. Abdullahi Bala
58. Prof. Felix N. Ikpe
59. Prof. Lateef B. Taiwo
60. Prof. A. A. Amusan
61. Chief Audu Ogbeh
62. Dr. Heineken Lokpobiri (Ph. D Leeds)
63. Prof. Jibrin N. Jibrin
64. Prof. Saminu Ibrahim
65. Prof. D. O. Asawalam
66. Prof. Felix Salako
67. Prof. P. I. Ogban



### FOOD BASKET 2019 CENTRAL LOCAL ORGANIZING COMMITTEE

1.	Dr. Bemgba Anjembe	-	Chairman
2.	Dr. O. Olatunji	-	Member/Secretary
3.	Professor Shaibu Idoga	-	Member
4.	Professor M.I. Ogbaji	-	Member
5.	Professor James O. Jayeoba	-	Member
6.	Dr. Matthew Adamu	-	Member
7.	Dr. N.B. Ibrahim	-	Member
8.	Dr. S.O.I. Abagyeh	-	Member
9.	Dr. Alfred Jimin	-	Member
10.	Rev. Dr. S.O. Odoemena	-	Member
11.	Dr. P.I. Agber	-	Member
12.	Dr. Ali Aminu	-	Member
13.	Dr. A.I. Sengev	-	Member
14.	Dr. A.O. Adaikwu	-	Member
15.	Dr. S.A. Ejembi	-	Member
16.	Mr. Friday Ibrahim	-	Member
17.	Mr. M.M. Ibrahim	-	Member
18.	Mr. Aliyu Zubairu	-	Member
19.	Mr. E.T. Ali	-	Member
20.	Mrs B.I Agada	-	Member
21.	Mr. V.M. Samson	-	Member
22.	Mr. T. Iorlamen	-	Member
23.	Mrs. Maureen Kajo FMA&RD, Makurdi	-	Member
24.	Programme Manager BNARDA	-	Member
25.	Representative, Ministry of Agriculture, Makurdi	-	Member
26.	Representative LBRBDA	-	Member
27.	Dr. Abraham Ajon AOCA, Yandev	-	Member

### LOCAL ORGANIZING SUB-COMMITTEE FOOD BASKET 2019

#### FUND RAISING/FINANCE

1.	Dr. Bemgba Anjembe	-	Chairman
2.	Professor Shaibu Idoga	-	Member
3.	Professor M.I. Ogbaji	-	Member
4.	Dr. Ali Aminu	-	Member/Secretary
5.	PM BNARDA	-	Member



**ABSTRACTS-FULL PAPER COLLECTION/ EDITORIAL**

- |    |                            |                  |
|----|----------------------------|------------------|
| 1. | Professor James O. Jayeoba | Chairman         |
| 2. | Professor Shaibu Idoga     | Member           |
| 3. | Dr. O. Olatunji            | Member           |
| 4. | Dr. Alfred Jimin           | Member           |
| 5. | Dr. A.O. Adaikwu           | Member/Secretary |

**TECHNICAL/INFORMATION AND COMMUNICATION**

- |    |                            |                  |
|----|----------------------------|------------------|
| 1. | Professor Shaibu Idoga     | Chairman         |
| 2. | Professor M.I. Ogbaji      | Member           |
| 3. | Professor James O. Jayeoba | Member           |
| 4. | Dr. Aminu Ali              | Member           |
| 5. | Dr. N.B. Ibrahim           | Member/Secretary |
| 6. | Dr. P.I. Agber             | Member           |
| 7. | Dr. Alfred Jimin           | Member           |

**OPENNING CEREMONY/OTHER SESSIONS**

- |    |                     |                  |
|----|---------------------|------------------|
| 1. | Dr. Aminu Ali       | Chairman         |
| 2. | Dr. Matthew Adamu   | Member           |
| 3. | Dr. S.A. Ejembi     | Member           |
| 4. | Mrs. Blessing Agada | Member           |
| 5. | Dr. Abraham Ajon    | Member           |
| 6. | Dr. Alfred Jimin    | Member           |
| 7. | Mr. Friday Ibrahim  | Member/Secretary |

**ACCOMODATION**

- |    |                        |                  |
|----|------------------------|------------------|
| 1. | Rev. Dr. S.O. Odoemena | Chairman         |
| 2. | Dr. Samuel Terhide     | Member           |
| 3. | Dr. S.T. Wuese         | Member/Secretary |
| 4. | Dr. A.I. Sengev        | Member           |
| 5. | Mr. Philemon D. Adaaku | Member           |



### PROTOCOLS/ENTERTAINMENT

- |    |                            |                  |
|----|----------------------------|------------------|
| 1. | Professor James O. Jayeoba | Chairman         |
| 2. | Dr. N.B. Ibrahim           | Member           |
| 3. | Mrs. Blessing Agada        | Member/Secretary |
| 4. | Mrs. Grace Daehung         | Member           |
| 5. | Dr. A.I. Sengey            | Member           |
| 6. | Mrs. Maureen Kajo          | Member           |
| 7. | Mrs. Esther Akor           | Member           |

### PUBLICITY/REGISTRATION

- |    |                          |                  |
|----|--------------------------|------------------|
| 1. | Dr. P.I. Agber           | Chairman         |
| 2. | Dr. S.T. Wuese           | Member           |
| 3. | Dr. A.I. Sengey          | Member           |
| 4. | Mrs. Judith Chia Chiluan | Member/Secretary |
| 5. | Mr. Philip Nzua          | Member           |

### FIELD TRIP/EXCURSION/EXHIBITION

- |    |                        |                  |
|----|------------------------|------------------|
| 1. | Professor Shaibu Idoga | Chairman         |
| 2. | Dr. Aminu Ali          | Member           |
| 3. | Dr. P.I. Agber         | Member           |
| 4. | Dr. S.O.I Abagyeh      | Member/Secretary |
| 5. | Mr. M.S. Adegoye       | Member           |

### SECURITY

- |    |  |                  |
|----|--|------------------|
| 1. | University Chief Security Officer        | Chairman         |
| 2. | University Deputy Chief Security Officer | Member           |
| 3. | Dr. Bemgba Anjembe                       | Member           |
| 4. | Dr. Samuel Terhide                       | Member           |
| 5. | Mr. E.T. Ali                             | Member/Secretary |



## FOREWORD

The 43<sup>rd</sup> Annual Conference of the Soil Science Society of Nigeria held on the 15<sup>th</sup> to 19<sup>th</sup> July, 2019 at the College of Agronomy Complex, Federal University of Agriculture, Makurdi with the theme: "Understanding Nigerian Soils for Sustainable Food and Nutrition Security and Healthy Environment". The choice of the theme was premised on the fact that Soil is a non-renewable resource and thus a sustained production of crops that would guarantee food and nutrition security while preserving and protecting the environment in Nigeria would require a clear understanding of the nature, dynamics and properties of these soils.

Papers were presented in the areas of Pedology, Soil survey and Climatology, Land use and Land Use Planning, Soil fertility and Plant nutrition, Fertilizer use and Management, Soil chemistry and Microbiology, Soil biology and Biochemistry, Biotechnology and Bioremediation studies, Soil physics, Soil and water management, Land degradation and Soil conservation.

The modern environmental movement that has evolved from a traditional emphasis on chemical reactions affecting plant growth to a focus on soil contaminant reactions was clearly demonstrated at this conference in terms of the number and quality of papers presented in this area. It was evident that the sources of these contaminants include fertilizers, pesticides, acid deposition, agricultural and industrial waste materials as well as oil spillage in the delta areas of Nigeria. Discussions on these contaminants and the soil chemical reactions that these contaminants undergo and how a knowledge of these reactions is critical in predicting their effect on the environment and quality of produce is well reported in this book of proceedings.

There were no field trips as a result of the prevailing security situation in the state. However, a presentation of the school laboratory project by the OCP Africa fertilizer company came in handy. This is a modern mobile Laboratory that conducts on site analysis of soil samples, instantly calibrates and provides field specific fertilizer recommendations. This ensures that fertilizers are applied at rates required by crop plants thus avoiding excesses that are bound to be applied when blanket recommendations are made with its attendant effect on the environment.

The papers presented at the conference which are published in this book of proceedings were painstakingly subjected to a peer review process. It is our hope that the information contained herein will be found helpful to environmentalists, Agriculturalists, policy makers, the academia and indeed a wide range of professionals.

**Bemgba Anjembe Ph.D.**  
Chairman, Local Organizing Committee.  
14<sup>th</sup> February, 2020.



## ACKNOWLEDGEMENTS

The Local Organizing Committee (LOC) of the 43<sup>rd</sup> Annual Conference of Soil Science Society of Nigeria (SSSN) wishes to express her profound appreciation to the Vice Chancellor, Federal University of Agriculture, Makurdi for providing venues and an enabling environment for the successful hosting of the conference. Apart from providing seed money as loan, his contributions towards provision of accommodation for the Leadership of our society is deeply appreciated.

His excellency, the executive Governor of Benue State Dr. Samuel Ibori apart from supporting the LOC financially, was represented at the opening ceremony by his deputy. He graciously granted the Society a state banquet, personally decorated our President and presented him with an honorary citizenship of the state.

The notable contributions of OCP Africa, Indorama, IFDC, Value Seeds Nigeria Limited, Teryima Nigeria Limited and other bodies significantly contributed to the success of the conference.

The chairmen of technical sessions and rapporteurs contributed greatly to the success of the conference. We sincerely appreciate all reviewers of manuscripts presented at the conference. We are also thankful to the Management and Staff of ASOFAD Printing Ltd. who worked tirelessly to ensure that our book of abstracts, programme of events, customized tags and certificates of attendance as well as the conference proceedings were delivered on time.

We are very highly indebted to the National Executive Committee of SSSN for their support and encouragement through this tasking period. The long conversations, arguments and constructive criticisms from the National President Prof. B. A. Raji largely improved the quality of our decisions and have tremendously contributed to our successful hosting of the 43<sup>rd</sup> Annual Conference.

**Bemgba Anjembe Ph.D**

Chairman, LOC

January, 2020



**TABLE OF CONTENTS**

	Page
Citation	ii
National Executive Committee of SSSN (2019)	iii
Members of Council Nigeria Institute of Soil Science	iii
Fellows of the Soil Science Society of Nigeria	iv
Food Basket 2019 Central Local Organizing Committee	vi
Local Organizing Sub-Committee Food Basket 2019	vi
Forward	ix
Acknowledgements	x
Table of Contents	xi
 <b>Opening Ceremony</b>	
Opening Remark by the Vice Chancellor	xix
Presidential Address	xx
Governors Speech	xxii
 <b>Conference Papers</b>	
<b>GROUP A: PEDOLOGY, SOIL SURVEY AND CLIMATOLOGY, LAND USE AND LAND USE PLANNING</b>	1
 Characterization and Land Suitability Evaluation of Soils on Diverse Land Forms in Ishiagu, South Eastern Nigeria for Yam ( <i>Discorea Spp</i> ) Cultivation <b>Nsor, M.E</b>	 2
 Fertility Assessment of Alluvial Soils of Different River Plains in Bayelsa State, Southern Nigeria, for Sustained Food Security <b>Achimota A. Dickson, Funso R. Kutu, and Joseph Aruleba</b>	 14
 Characterization and Classification of Soils Affected by Erosion in Igbudu Ikwo Southeastern Nigeria. <b>Obasi S. N., Obasi C. C., Nweke V., Nsoanya L. N.</b>	 32
 Characterization and Mapping of upland Soils formed from Sand Stone and Micaceous Schist in Koko/Besse LGA, Kebbi State. <b>Ande O.T., Are K.S., Adeyolanu O.D., Ojo A.O., Oke A.O., Denton O., Adelabu Lucas, Omodele Taiwo, Oluwatosin G.A. and Adediran J.A.</b>	 43
 Suitability Ratings of a Teak ( <i>Tectonagrandis</i> ) Plantation Soils for Rubber ( <i>Heveabrsiliensis</i> ) Production in Edo State, Nigeria <b>Oghomieje, L.A., S.O. Idoko, T.U. Esekade And S. Ighedosa</b>	 60
 Assessment of some Selected Soil Chemical Properties of Irrigated Farmlands of Jere Bowl Near Maiduguri, Nigeria <b>Adam Lawan Ngala, Ismaila Mohammed, and Dimhe James</b>	 70
 Assessment of the Productivity Status of Vertisols in Sudan Savanna Zone of Nigeria <b>S. Idris, H.M. Lawal, M.M Maiyaki, A.T Mahmud and M. Adamu.</b>	 79



Evaluation of Soils and Irrigation Water Quality along Meme River Lokoja, Kogi State, Nigeria for Maize ( <i>Zea Mays</i> ) Production Adava, I.O., Otene, I.J.J., Adeyanju, O.E., and Obafunmi E.O.-	88
Effects of Biochar Amendments on Nitrate Leaching and ZPC in Four Soil orders from Nigeria's Savanna Zones Solomon Rejoice Ibrahim, Abdu Nafiu, Yusuf Ado Adamu and Bello Muktar	97
Characterization, Classification and Management of Soils formed under different Landscape Positions in Osina, Imo State, Southeastern Nigeria for Agricultural Production Egboka*, N.T., Onweremadu, E.U., Aririguzo, B.N., Ahukaemere, C.M., Okoli, N.H. And Afangide, A.L.	115
Evaluation of Soil Nutrient Status of Selected Upland Rice Farms in Omi Yagbawest Local Government Area, Kogi State Nigeria Otene, I.J.J., Adava, I., Muhammed, B., and Bashir S.O.	128
Evaluation of Profile Distribution of forms of Aluminium, Iron, and Manganese in Selected Agro Ecological Zones of Nigeria Muhammed, B., Samndi, A.M., Otene, I.J.J., and Abubakar, M.A.	135
Evaluation and Mapping of Soil Salinity and Sodicity in Sector One of Watari Irrigation Scheme, Kano State. Bashir A.Y, Samndi A.M, Adam M.A, Moji K.T and Auwal M.	144
Suitability Assessment of some Soils along a Toposequence in Ambrose Alli University Teaching and Research Farm for the Cultivation of Rice ( <i>Oryza Sativa</i> ), Edo State Southern Nigeria Eghaghara, O. O. and Suleiman, J.A.-	153
Spatial Variability of Selected Soil Properties in a Cultivated Gravelly Landscape in Minna, Niger State, Nigeria Lawal, B.A., Tsado, P.A., Afolabi, S.G., Oguntoye, G.P. and Ogbodo, C.N.	163
<b>GROUP B: SOIL FERTILITY AND PLANT NUTRITION, FERTILIZER USE AND MANAGEMENT-</b>	170
Growth and Yield Response of Rice ( <i>Oryza Sativa</i> ) to Application of some Microminerals (Cu, Fe, Mn And Zn) in Doma, Nasarawa State Collins Maichibi, Bemgba Anjembe and O. Olatunji	171
Effect of Fertilizer, Rhizobium Inoculum and Manure Application on Production of Soybean [ <i>Glycine Max</i> (L.) Merr.] in Northern Nigeria Savanna B. Ukem, N. Kamai, E. Sangodele, J. Abikoye and E. Chinedu	183
Effect Of Fertiplus and Crystallizer on Soil Properties and Yield of White Maize in Makurdi, Southern Guinea Savanna Zone of Nigeria Ali, A., Anjembe, B.C. and Usman, M.	198



Growth, Yield, and Nutrient uptake of edible Amaranth ( <i>Amaranthus Hybridus</i> L.) in response to application of Composted and Stacked Pig Manure	
Oderinde O. L., Adejuyigbe, C. O., Azeez J. O. and Olowu V. I. O.	258
Effects of Aqueous Ethanol Moringa ( <i>Moringa Oleifera</i> Lam.) Leaf extracts on growth and yield of Soybean ( <i>Glycine Max</i> L. Merrill)	
Mustapha, Y., Isah, L., Hayastuddeen, A.M. and Oghovwa, M.	268
Impact of Biochar fortified with Urea on Soil properties, nutrient uptake and dry matter yield of <i>Amaranthus Cruentus</i> in a Rainforest Ultisol	
Otobong B. Iren and Mmadara I. Umanah	275
Effects of Fertiplus and Crystallizer on the Growth and Yield of White Maize in Lafia, Southern Guinea Savanna Zone of Nigeria	
Ali, A., Usman, M. and Maku, A.A.	285
Growth and Yield of Tomato ( <i>Lycopersicon Lycopersicum</i> ) on Soils contaminated with Diesel and Engine Oil	
Audu, A., Okoh, M.O., Otene, I.J.J., Ambakhian, S.O, Muhammed, B., and Ayuba, S.A.	247
Influence of fertilizer types on soil physical properties, growth and yield of Maize ( <i>Zea Mays</i> L.) in Southwest Nigeria	
Ewetola E. A. and Isola J. O.	261
Effect of Biochar on available Phosphorus Status in Alfisols Derived from Charnockite in Ekiti State, Nigeria.	
Ilori, A. O. A., Shittu, O. S., Ayodele, O. J., Awe, G.O. and Oso, A. O.	269
Effects of Integrated Nutrients Management on Weed Infestation, Growth and Yield of Cowpea varieties ( <i>Vigna Unguiculata</i> L.) in moist Savanna of Nigeria.	
M.A. Yusuf, A.O. Uzoma, A.Y. Mamudu and M.T. Salaudsen	279
Effects of different Organic amendment sources on the Physico-Chemical Properties of a nutrient depleted Vertisol	
Unagwu, B. O., Ene, J., Azuka, C.V.	287
Effects of Poultry Manure and Urea Fertilizer on some Soil Chemical and Microbial characteristics, in a Rubber Nursery of an Ultisol in Southern Nigeria.	
Idehen, C.N., S.O. Idoko, T.U. Esekhade, R.O. Okundia, U. Esemuede And E. Unabor	294
Maize Growth and Performance in different Landform Units of Coastal Plains Sand, Imo State, Nigeria	
Ukaegbu, E.P., Asadu, C.L.A. Ezeaku. P.I., Nwaigwe, M.O.	302
Evaluation of Single Super Phosphate fertilizer on growth and yield of Sesame ( <i>Sesamum Indicum</i> L.) in Wukari, Southern Guinea Savanna of Nigeria.	
Onwu A. C. Waizah, Y and Apagi, J. N.	309



Effects of Fertiplus and Crystallizer on the growth and yield of Yellow Maize in Makurdi <b>Ali, A., Anjembe, B.C. and Usman, M.</b>	314
Soil Chemical Properties, Nutrient uptake and performance of Tomato ( <i>Lycopersicon Lycopersicum</i> ) and Cucumber ( <i>Cucumis Sativus</i> ) on Saline Soils amended with Gypsum and NPK fertilizer <b>Okoh, M.O., Audu, A., Otene, I.J.J., Ayuba, S.A., and Adebayo, A.A.-</b>	326
Calibration of Poultry Manure for fresh yield of Fluted Pumpkin ( <i>Telfairia Occidentalis</i> ) and it's effect on Soil Chemical properties at Ibadan, Nigeria <b>Kayode, C.O., Ayanfeoluwa O.E., Komolafe A.F., Ezekiel-Adewoyin D.T., Ogunleti D.O., and Akinlabi A.E.</b>	343
Evaluation of Heavy Metal Status, quality and safety of Vegetables sold in Markets located in Makurdi Metropolis, Nigeria for Human Consumption <b>Bemgba Anjembe and Nguveren Tyavkase</b>	349
Influence of some Soil properties on the availability of some Nutrients from the application of Urea Super Granule to a Lowland Rice Soil <b>Girei, A. H., Nabayi, A., Amapu, I.Y., Mudiare, O. J. and Abdulkadir, A.,</b>	357
<b>GROUP C: SOIL CHEMISTRY AND MICROBIOLOGY, SOIL BIOLOGY AND BIOCHEMISTRY, BIOTECHNOLOGY AND BIOREMEDIATION STUDIES</b>	365
Use of Lime and Biochar for Acidic Soil Remediation, Growth and Macro-Nutrients uptake of Maize ( <i>Zea Mays L.</i> ) <b>Ndor E, Jayeoba O.J and Azagaku E.D</b>	366
Comparative Studies of Microbial Properties in Wetland and Coastal Plain Soils of Calabar Agro-Ecological Zone <b>Akpan, J. F., Isong, I. A. and Solomon, M. G.</b>	372
Evaluation of the Phytoremediation Potentials of some Vegetables for Remediation of Heavy Metal contaminated Soils <b>Ndor E, Jayeoba O.J and Azagaku, E.D</b>	383
Effect of Treated Refinery Effluent on Bacterial and Fungal Population in Soil. <b>Ogboghodo, I. A., Okwerekwu, Y.A., Ogbemudia, I. And Edosa, V. I. O</b>	390
Status and Distribution of Exchangeable Potassium in aggregate size fractions of Coastal Plain Sands under Five Land-Use types in Umudike, Abia State, Nigeria. <b>Akinmutimi, A.L. and Onyetorochukwu, D.F.</b>	399
Interactive Antagocism of Three Biological Control Agents against <i>Fusarium Oxysporum</i> Causal Organism of Fusarium Wilt of Cucumber <b>Edosa, V.I.O., Osarhiemen, O. V., Ogbemudia, I., Akaeze, O.O.-</b>	407



Acidity Indices Response to Biochar Application on Coastal Plain Sand at Umudike Southeast, Nigeria <b>Onwuka, B.M. and Onwuka, M.I.</b>	413
Effect of Herbicides on Soil Microbial Biomass, Biological Nitrogen Fixation and Yield of Cowpea in Makurdi, Nigeria <b>Ibrahim, M.M and Agbo, M.S.</b>	420
Nematode Population as Influenced by Different Tillage Practices in Maize-Cowpea Cropping System. <b>Adesina Abudlkabir Wahab, Afe Isaac Ade, Ismail Adeshina Sanusi</b>	429
Assessment of Copper, Lead and Nickel Contamination in Soils and Cocoyam ( <i>Xanthosoma Sagittifolia</i> ) Grown Near Waste Dumpsites in the outskirts of Uyo Metropolis, Akwa Ibom State <b>Umoren, A.S., Effiong, G.S. and Udosen, C.I.</b>	444
Effects of Tannery Wastewater on Soil Physico-Chemical Properties and Heavy Metals Concentration in Challawa Industrial Estate of Kano State. <b>Salihu Ahmad Pantami</b>	457
Soil Physicochemical properties, nutrient uptake, growth and yield of Fluted Pumpkin ( <i>Telfairia Occidentalis</i> ) under combined application of Cattle Dung and NPK fertilizer in Anyigba, Kogi State, Nigeria. <b>Frank, O.O., Paul, O.J., Amhakhian, S.O, and Otene, I.J.J.</b>	468
Tolerance of <i>Jatropha Curcas</i> L.to different levels of Crude Oil Pollution in the Niger Delta region of Nigeria. <b>Christiana J. Ijah and Otobong B. Iren</b>	488
Dynamics of Soil pH, Phosphorus availability and Cowpea Growth in simulated Aluminum Toxic Alfisols amended with Phosphorus Fertilizers <b>Oyeyiola Y.B., Omueti J.A.I., Kolawole G.O., Olatunji O.O.</b>	495
A Review of the Status of Soil Pollution in Nigeria with specific reference to Eleme Community in Rivers State of the Niger-Delta Area <b>Ugadu, M.E. and Chude, V.O.</b>	508
Identification and Isolation of Microorganisms in Wetland Soils of Niger State, Nigeria <b>Zakari, S.I, Bello, O.S., Adegbite, M.A., Mohammed, A. and Sanda, A.R.</b>	519
Vertical Distribution of Cadmium in Coastal Soils affected by Crude Oil Spillage at Bodocity, Rivers State. <b>Afangide A. I., E.U. Onweremadu., J. U. Amaechi., Okon M. A., Okoli N. H And Egboka, N. I.</b>	526
Impact of Soil Microorganisms and Water Availability Management for Sustainable Agricultural Productivity <b>Ojobo J.O. and Ezendu C.O.</b>	537



- Response of Soil, Rubber Seedlings to the Application of Mycorrhizae, NPK and Poultry Manure in Southern Nigeria  
**Izevbogie F.C, Idoko S, Okundia R.O, Esemuide U, Idehen C, Idika R, and Unabor E,** - - - 547
- Phosphorus Sorption in Soils Amended with Rice Husk Biochar  
**Ayodele, T.O And Thomas E.Y.** - - - 560
- Dynamics of Soil pH, Phosphorus availability and Cowpea Growth in simulated Aluminum Toxic Alfisols Amended with Phosphorus Fertilizers  
**Oyeyiola Y.B., Omueti J.A.I., Kolawole G.O., Olatunji O.O.** - - - 574
- Effects of Irrigation Water Quality on Soil Microbial Enzyme activities and Crop Quality in Minna, Niger State.  
**Awele O. Osa-Afiana, Abdullahi Bala, Anthony O. Uzoma, Abubakar Bello.** - - - 587
- Symbiotic Properties of Rhizobia Strains Isolated from Soils of Birnin-Gwari and Shika In Kaduna  
**Obioha, N. J., Bolarinwa, T. S., Uzoma, O.A., Bala, A. and Osunde, A.O.** - - - 600
- Response of Four Cowpea Varieties to Bradyrhizobial Inoculation in some Soils of Northern Guinea Savanna Zone of Nigeria  
**Tanko, F., D.T Ezekiel-Adewoyin., A. O. Osunde, And A. Bala** - - - 608
- Biological Nitrogen Fixation and Productivity of Soybean in Maize-Soybean Intercrop under Tillage and Bradyrhizobium Inoculation on Alfisols of Northern Guinea Savanna Nigeria  
**Omeke, J. O. Yusuf, A. A., Uyovbisere, O. E. and Aliyu, I. A.** - - - 617
- Seasonal and Irrigation Reuse Quality Assessment of Effluents from Experimental Vertical-Flow Wetlands Treating Domestic Wastewater polluted with Hydrocarbons  
**Abdulkadir, S., Adamu, U.K., and Scholz, M.** - - - 629
- Short-Term effect of Tillage and Nitrogen Fertilization in Bradyrhizobium Inoculation Soybean-Maize-Based Cropping Systems on selected Physical Soil Productivity Indicators  
**Omeke J. O., Nnaji, G. U. and Ihem E.E.** - - - 645
- Growth and Yield Response of Inoculated Soybean (*Bradyrhizobium Japonicum*) to Inorganic Fertilizer (Soil and Foliar Application) in the Northern Savannah Zone of Ghana.  
**Ezekiel-Adewoyin, D. T., Tanko, F., Ayanfeoluwa, A.O., Komolafe, O., Kayode, C.O. and Aklinlabi A.E.** - - - 656



<b>GROUP D: SOIL PHYSICS, SOIL AND WATER MANAGEMENT, LAND DEGRADATION AND SOIL CONSERVATION AND OTHERS</b>	<b>668</b>
Food Security in Nigeria: Challenges and Wayforward. Hejimala Augustine Chukwumeka	669
Impact of Land Use on Soil Quality in Ijebu-Igbo, Ogun State, South West Nigeria Amosa S.A.	680
Geo-Spatial variability of Soil Physico-Chemical properties for Sorghum based cropping system of Sudano-Sahelian zone of Nigeria. Amin, M.A, Uzaiifa, U.K, Abdulkadir, S., Adam, M.A, Adam, I.A, Ahmad, U.B	686
Evaluation of Water resources potentials of part of River Ogbadigbo catchment area for agricultural uses in Okpokwu LGA of Benue State. Bodinga T. Garpiya, M. S. Adegoye, Kyat M. Makama, Patrick Kefas, and S. A Gisilanbe -	698
Effects of Tillage Practices on selected Soil Properties in Sudan Savannah Agro Ecological Zone; A case study of Federal University Dutse Teaching and Research Farm. Nabayi, A., Girei, A.H., Hayatu, N.G., Mahmud, A.T., Usman, S., Santuraki, A.H., Haruna, F.D., Yusif, S.A., Nkereuwem, M.E and V.O. Onokebhagbe. -	704
Estimation of Soil loss due to Cassava Harvesting in Ibadan, Nigeria S.O. Oshunsanya, V.M. Samson	713
Influence of Organic Matter on some selected Physical and Hydraulic Properties of a Sudan Savannah Entisols Nabayi, A., Danmowa, N.M., Hayatu, N.G., Girei, A.H., Santuraki, H.A., Haruna, F.D., Yusif, S.A., Onokebhagbe, V.O., Maunde, M.M., Nkereuwem, M.E. and A.T. Mahmud.	719
Effects of Different Tillage practices on Soil strength and selected Agronomic characteristics of Cowpea ( <i>Vigna Unguiculata</i> [L.] Walp) Shittu, K. A; Oyedele D. J; Nwoke, O. C.; Lamidi, W. A and Babatunde K. M.	726
Effect of Tin-Mine Residue on Soil Physico-Chemical Properties and Agricultural Production in Jos Plateau of Nigeria Gyang, B.D., Ezeaku, P.I., Asadu, C.L.A., Umoren, A.S. and Uchechukwu, K.	735
Assessment of Land Degradation in Three Locations and Sampling Periods in Physiographic Units under Cultivated Crops in Sub- Humid Tropics of Nigeria. Onyishi I.F. and P.I. Ezeaku	745



Impact of Land Degradation on Cultivated Arable Crop Soils along Three Toposequences in Nsukka Sub Humid Tropical Area of Nigeria. <b>Onyishi I.F. and P.I. Ezeaku</b>	764
Effect of Continuous Cultivation on some Physical and Chemical Properties of the Soils of Bankanu and Balkore Area of Kware Local Government, Sokoto State <b>N.G. Hayatu, S.S. Noma, S.A. Yusuf, M.R. Sharif, S.A. Lukman, M. Haliru, M.M. Sauwa, N. Shehu and G.A. Abubakar</b>	782
Assessment of Planting date and variety on Water use efficiency of Potato ( <i>Solanum Tuberosum L.</i> ) in Kuru, Jos, Nigeria, <b>Dayok, S.T., Agber, P.I., Tetok, N.I. and Gani, A.T.</b>	793
Integrated Soil and Water Management strategies and performance of Maize at Danbatta and Jangwano Villages in Kano and Jigawa States Nigeria <b>Sanda, Ahmad R. and Ramatu Idris</b>	802
Assessment of Soil loss tolerance limits for Makurdi, Benue State Nigeria <b>Agada B.I.(Mrs), Dr. Agber, P.I., Professor Idoga S., Dr. Ugesi F.D.</b>	805
Spatial Variability of Soil Loss in the Ban Dan Na Kham Watershed of Northern Thailand. <b>Madueke, C.O., Shrestha, D.B.P., Nyktas, P. and Orlowsky, K.</b>	813
Seasonal and Irrigation Reuse quality assessment of Effluents from experimental Vertical-Flow Wetlands treating domestic Wastewater polluted with Hydrocarbons <b>Abdulkadir, S., Adamu, U.K., and Scholz, M.</b>	824
Physical and Hydraulic properties of five Alfisol Delineated for Dry Season Vegetable production in Ekiti State <b>Ayodele A.A, O.O Nurudeen and Ajayi A.S.</b>	841
Evaluation of Saxton and Rawls Pedotransfer functioni in predicting Soil Hydraulic properties of Sandy Loam Soil in Lafia, Nasarawa State, Nigeria <b>V. B. Ogbe, S. M. Amana, O. J. Jayeoba, M. A. Abubakar and T. O. Negedu</b>	848
Effects of Maize-Cowpea Intercrop on A typic Plinthustalfs Soil of Southern Guinea Savannah Zone, Nigeria. <b>Alabi, K.O., Afe, I.A., and Wahab, A.A.</b>	857
Combating Tropical Soil Degradation: The Role of Nitrogen Fertilizer as a Climate-Smart Strategy towards Maize ( <i>Zea Mays L.</i> ) productive Cultivation <b>M. S. Adiaha and A. A. Onwukwe</b>	865
Communique -	883



## RESPONSE OF FOUR COWPEA VARIETIES TO BRADYRHIZOBIAL INOCULATION IN SOME SOILS OF NORTHERN GUINEA SAVANNA ZONE OF NIGERIA

Tanko, F., D.T Ezekiel-Adewoyin., A. O. Osunde, and A. Bala

Department of Soil Science and Land Management, School of Agriculture and Agricultural Technology, Federal University of Technology Minna, PMB 65, Minna Niger State.

### ABSTRACT

Yield of cowpea in savanna soils of sub-sahara Africa has remained low as a result of less effective native rhizobia, necessitating the use of elite inoculant strains to improve Biological Nitrogen Fixation. The main objective of this study was to observe the response of four varieties of cowpea to inoculation by two elite strains of bradyrhizobia in some soils of Northern Guinea Savanna (NGS) zone of Nigeria. Two greenhouse experiments were carried out at the Teaching and Research Farm of the Federal University of Technology Minna. To determine (1) the size of the native rhizobial population in the soils using the Most Propable Number (MPN) technique and (2) The Need to Inoculate cowpea in the soils under study. The treatments for the MPN experiment consisted of (i) soils in area of close proximity to homestead (< 50 m), and those far from homestead (> 250 m) and (ii) cowpea varieties (Kanannado, IT93K-452-1, IT97K-499-35, and IT90K-277-2). The experiment was laid out in a 2 x 4 factorial in a completely randomised design (CRD) replicated four times. Treatment for the need to inoculate experiment consisted of (i) five locations (Rijau, Kontagora, Magama, Mashegu, Mariga,) in the NGS zone of Nigeria (ii) four nitrogen sources (plants inoculated with *Bradyrhizobium* sp. strain BR 3262, *Bradyrhizobium* sp. strain BR 3267, control (neither N nor inoculant rhizobia applied), and plants treated with N in form of urea at the rate of 100 kg N ha<sup>-1</sup>, (iii) four cowpea varieties, (iv) two levels of proximities to homestead. Proximity to homestead and cowpea varieties were the same as in the MPN experiment. The experiment was laid out in a 5 x 4 x 4 x 2 factorial fitted to a completely Randomised block design (CRD) replicated four times. In general the MPN value ranged from 4.61 x 10<sup>6</sup> to 7.19 x 10<sup>7</sup> cells g<sup>-1</sup> of soil. MPN value for IT90K-277-2 variety (9.27 x 10<sup>6</sup> cells g<sup>-1</sup>) was significantly lower than that estimated by the other varieties Kanannado (3.31 x 10<sup>7</sup> cells g<sup>-1</sup>), IT93K-452-1 (1.76 x 10<sup>7</sup> cells g<sup>-1</sup>), and IT97K-499-35 (1.42 x 10<sup>7</sup> cells g<sup>-1</sup>). In spite of the high population of the native rhizobia in the soils, the cowpea varieties responded to inoculation suggesting the effectiveness of the elite bradyrhizobia strains used. Proximity of sampling site close to homestead had significantly (P < 005) higher Shoot dry weight than those away from homestead. The results from this experiment showed that the main effects of Nitrogen sources and varieties had significant effect on shoot dry weight of cowpea in the all locations. The shoot dry weight of plants that received 100 kg N ha<sup>-1</sup> was marginally higher than those of the inoculated plants, indicating that elite rhizobia strain only partially met the N need of the cowpea crops. Further studies are suggested to evaluate larger number of bradyrhizobial on this environment.

### INTRODUCTION

The most frequently limiting nutrient in soils of the Nigerian Savanna is nitrogen, due to its continuous depletion from the soil pool by processes such as volatilization, leaching and removal of nitrogen containing crop residues from the soil (Aliyu *et al.*, 2013). Legumes such as cowpea that biologically fix nitrogen are integrated as part of the

cropping systems to optimize the supply of this nutrient in agricultural systems. Legumes are able to meet part of their nitrogen requirement through biological nitrogen fixation (BNF) (Bagayoko *et al.*, 2000).

This is however not the case when the native rhizobia are absent, small in number, or the presence of incompatible or less effective



indigenous rhizobia strains in soil. Under this situation, BNF can be improved through inoculation of soils with effective and compatible elite rhizobial strains (Abaidoo *et al.*, 2007). The presence of large population density of compatible rhizobia does not however, impede the possibility that responses to inoculation can be obtained if competitive and highly effective strains are introduced in high quality inoculants (Giller, 2001). Inoculating cowpea with rhizobia has been used to achieve substantial increases in legume nodulation, grain and biomass yield, nitrogen fixation and post-crop soil nitrate levels over this period. These gains are usually highest when the inoculated legumes are grown in devoid of rhizobia or low-rhizobia soils, but marginal in soils already containing high number of compatible rhizobia (GRDC, 2013).

Inoculation is not a common practice in Nigeria, although results from field trials involving cowpea often varied, with some showing significant increases in yield Giller *et al.* (2005) and others showing little or marginal differences between inoculated and uninoculated crops. Results from Southern Africa, shows variability in soil fertility as farmers preferably apply nutrient resources to fields closest to homesteads leading to gradients of decreasing soil fertility with proximity away from homesteads (Zingore *et al.*, 2007). Information on weather soils in the Nigerian Savanna exhibit similar gradients in fertility is lacking. Additionally, few studies have been conducted on the varietal response of cowpea to inoculation in soils of Nigerian Savanna. In view of this, a need to inoculate trial was carried out in the teaching and research farm of the federal university of Technology Minna to determine the following

- (i) Population of native rhizobia present in some soils of Northern Guinea Savanna.
- (ii) Response of four cowpea varieties (Kanannado, IT93K-452-1, IT97K-499-35, and IT90K-277-2) to *Bradyrhizobium* inoculation.

- (iii) To investigate if there exist a relationship between cowpea response to inoculation and proximity of field to homestead.

## MATERIALS AND METHODS

### Description of Study Area

The study was conducted in the screen house of the Federal University of Technology Minna. The study area is located in the Southern Guinea Savanna zone of Nigeria which lies between longitudes 90 30' and 90 40' E and latitudes 6° 30' and 6° 35' N at an elevation of about 258.5 m above sea level.

### Soil Sampling and Analysis

Soil samples were collected from 5 locations in the Northern Guinea Savanna (NGS) (Rijau, Kontagora, Magama, Mashegu, and Mariga,) of Niger state. At each location, soils were collected from sites close to homesteads (0-50 m) and soils sampled further away from homesteads (at least 250 m away) and bulked to form composites sample using sterilized shovel before and after sampling. In order to characterize the soil, part of the soil samples was taken to the laboratory, air-dried gently crushed and passed through a 2 mm sieve for physical and chemical analyses by standard methods (IITA, 1989). The remaining samples were prepared moist for greenhouse study.

### Treatments and Experimental Design

The treatments for the MPN experiment were (i) 2 proximity of field to homestead soils in area of close proximity to homestead (< 50 m), and those far from homestead (> 250 m) (ii) 4 cowpea varieties (Kanannando, IT93K-452-1, IT97K-499-35, IT90K-277-2). The experiment was arranged in a 2 x 4 factorial fitted to a completely randomized design (CRD) replicated four times. While for the need to inoculate trial the treatments were (i) 5 locations (Rijau, Kontagora, Magama, Mashegu, and Mariga). (ii) 4 nitrogen sources (urea at the rate of 100 kg ha<sup>-1</sup>, plant inoculated with *Bradyrhizobium* sp. strain BR 3262 or *Bradyrhizobium* sp. strain BR 3267, and control (Neither N nor inoculant rhizobia applied), (iii) 4 cowpea varieties (iv) 2 proximity (varieties and



proximity to homestead was the same as that of MPN). The experiment was a 5 x 4 x 4 x 2 factorial, fitted to a completely randomised design (CRD) and replicated four times.

#### Greenhouse study

##### MPN Determination

The assay was conducted using modified Leonard jar method in Southern Australia (Howieson *et al.*, 2014) using coarse sand as the potting medium. Coarse sand was washed thoroughly with water to remove all traces of dissolved nutrients before sun drying. 2.5 kg of the sand was transferred to a 3 litre pot and 200 ml of calcium solution was added directly to the sand. Cowpea seeds were sown at the rate of 4 seeds per pot and thinned to 1 per pot a week after planting. Plants were watered daily using sandsman's nitrogen free nutrient solution (Sandsman, 1970). Inoculated was done a week after planting with 1 ml aliquots of the soil suspensions made from serial dilution of soils sampled. Each soil sample was initially diluted 20-fold by adding 10 g of soil in 190 ml of sterile distilled water. The resulting suspension was thoroughly mixed and then used to prepare a 10-fold serial dilution (Woomer, 1994). Each dilution level was replicated four times resulting in a total of 24 pots per each soil sample. Harvesting was done in a period of seven weeks after planting to observe the patterns of nodule appearances. The (+) sign was used to indicate presence of at least one nodule, while (-) denoted absence of nodules (Vincent, 1970).

##### Need to Inoculate Trial

Soil samples were added to 2.5 litre pot at 2 kg per pot. To each pot was added 372.6 mg of P, K, Mg, Zn, Mo, and B fertilizer and thoroughly mixed. Cowpea seeds were planted at 2 per pot and thinned to one seedling per pot one week after emergence. N treated plants were supplied with nitrogen and was split applied to the N treatment at first week (81.60 mg) and second week (244.80 mg) after planting. Inoculants were applied to the seeds before planting in inoculated treatment. The plants were

watered daily with sandsman's N free nutrient solution 200 ml per pot for the first four weeks and later sterile distill water was used, plants were harvested 7 weeks after planting.

##### Data Analysis

The data was subjected to statistical analysis using MINITAB 17.0. Analysis of variance (ANOVA) of the general linear model was used to check for significant effects and significant means were separated using Fisher least significant difference (L.S.D) method.

## RESULTS

### Physical and chemical characteristics of the experimental soil

The physical and chemical properties of soils used for this study are presented in (Table 1). Soils of the experimental site from both close and far proximities to homestead were either sandy loamy or loamy sandy, except for Mashegu close which was sandy soil. Soil pH for both proximities to homestead in the 5 locations was slightly acidic to near neutral ranging from 5.31-6.79 but a moderate acidity of 4.77 was observed for soils sampled near homestead in Kontagora. Total nitrogen was low for all locations and proximities to homestead ( $0.06 \text{ g kg}^{-1}$  -  $0.22 \text{ g kg}^{-1}$ ), also organic carbon was low in the 5 locations but a moderately high value was observed with Kontagora close to homestead ( $12.45 \text{ g kg}^{-1}$ ). Available P ranges from low to moderately low ( $18.0 \text{ mg kg}^{-1}$ ) in Kontagora close to homestead and very high ( $31.0 \text{ mg kg}^{-1}$ ) in Mariga away from homestead. The result of Effective cation exchange capacity (ECEC) ranges between  $5.79 \text{ cmol kg}^{-1}$  -  $8.25 \text{ cmol kg}^{-1}$  respectively.

### Estimate for rhizobial population of four varieties of cowpea in NGS of Niger State

The rhizobial populations as estimated using four (4) cowpea varieties as trap host are presented in (Table 2). The soils of the study sites had high numbers of indigenous rhizobia ( $>10^3$  rhizobia cells  $\text{g}^{-1}$  of soil). However, soils sampled close to homestead in Mashegu, and for those away from



homestead in Mariga and Magama had the highest population ( $71.84 \times 10^6$  cell  $g^{-1}$  of soil) using Kanannado variety while the least native populations ( $4.61 \times 10^6$  cell  $g^{-1}$  of soil) were obtained using IT93K-452-1 variety for Kontagora and Rijau using soils sampled close to homestead.

**Effect of nitrogen sources, proximity to homestead and varieties on shoot dry weight of cowpea**

The main effect of nitrogen sources significantly ( $P < 0.05$ ) affected shoot dry weight of cowpea in the 5 locations (Table 3). The Urea treated plants produced the highest shoot dry weight of cowpea in all locations while the control plants had the smallest shoot biomass. There was a significant response of cowpea to rhizobia inoculation, plant inoculated with either or both of the inoculant strains produced the greater shoot dry weight than the control in all soils.

The main effect of proximity to homestead on shoot dry weight of cowpea was significant ( $P < 0.05$ ) for all locations except in Magama, and Rijau (Table 3). Soils sampled close to homestead supported higher cowpea shoot dry matter than soils further away from homestead.

The varietal effect on shoot dry weight was significant in the five locations (Table 3). The Kanannado variety produced the highest shoot dry weight while IT90K-277-2 variety had the smallest dry weight but the reverse was the case with IT93K-452-1 in Mashegu. IT93K-452-1 and IT97K-499-35 produced similar shoot biomass at four out of the five locations. The interactive effect among nitrogen source, proximity to homestead, and varieties significantly affected the shoot dry weight of cowpea in all location.

**Table 1. Physico-chemical properties of soils used for this study**

Location	Proximity	pH CaCl	OC g kg <sup>-1</sup>	N	Avail.P (mg kg <sup>-1</sup> )	Textura l Class	Ca (cmol kg <sup>-1</sup> )	ECEC
Kontagora	Close	4.77	12.45	0.20	18.0	LS	5.20	7.64
	Far	5.31	2.55	0.10	6.0	LS	4.40	6.13
Mashegu	Close	5.84	3.75	0.10	6.0	S	4.64	6.34
	Far	5.97	4.95	0.08	5.0	SL	5.60	7.43
Rijau	Close	6.57	4.80	0.06	8.0	LS	4.64	6.17
	Far	5.99	6.15	0.15	9.0	SL	4.88	6.49
Mariga	Close	5.78	6.45	0.22	6.0	LS	6.40	8.25
	Far	5.85	8.25	0.10	31.0	LS	5.20	7.53
Magama	Close	5.47	2.70	0.14	6.0	SL	4.56	5.79
	Far	5.57	3.75	0.20	7.0	LS	5.44	7.18

\*SL-sandy loamy \*LS-loamy sand \*S-sandy



**Table 2** Estimate of rhizobial population for four varieties of cowpea at the 5 locations in ( $\times 10^6$  cell  $g^{-1}$  of soil)

Location	Proximity	Kanannado	IT93K-452-1	IT97K-499-35	IT90K-277-2	Means
Kontagora	Close	10.94	4.61	11.24	7.19	8.50
	Far	22.48	31.85	12.27	11.24	19.46
Mashegu	Close	71.89	22.48	11.24	7.19	28.2
	Far	12.27	22.48	11.24	7.19	13.30
Rijau	Close	12.27	4.61	9.26	12.27	9.60
	Far	12.27	12.27	9.26	10.94	11.19
Mariga	Close	22.48	42.59	11.24	7.19	20.88
	Far	71.89	12.27	12.22	11.24	26.91
Magama	Close	22.48	10.94	42.59	7.19	20.80
	Far	71.89	12.27	11.24	11.24	26.66
<b>Means</b>		<b>33.06</b>	<b>17.64</b>	<b>14.18</b>	<b>9.29</b>	

**Table 3** Effect of nitrogen sources, proximity to homestead and varieties on shoot dry weight ( $g\ plant^{-1}$ ) of cowpea

Treatment	Kontagora	Mashegu	Rijau	Mariga	Magama
<b>Nitrogen sources (N)</b>					
Control	1.27c	1.16c	1.39c	1.52b	1.13c
Urea	1.87a	1.86a	2.13a	2.06a	1.75a
BR 3262	1.54b	1.48b	1.36c	1.50b	1.27bc
BR 3267	1.22c	1.80a	1.65b	1.91a	1.30b
Significance	**	**	**	**	**
LSD (0.05)	0.17	0.17	0.23	0.25	0.16
<b>Proximity (P)</b>					
Close	1.55a	1.70a	1.63a	1.85a	1.33a
Far	1.40b	1.45b	1.63a	1.64b	1.40a
Significance	*	*	NS	*	NS
LSD (0.05)	0.12	0.12	0.16	0.17	0.11
<b>Varieties (V)</b>					
Kanannado	1.95a	1.88a	2.01a	2.02a	1.67a
IT93K-452-1	1.39b	1.34c	1.65b	1.89ab	1.17c
IT97K-499-35	1.42b	1.49bc	1.72b	1.71b	1.38b
IT90K-277-2	1.14c	1.58b	1.16c	1.37c	1.23bc
Significance	**	**	**	**	**
LSD (0.05)	0.17	0.17	0.23	0.25	0.16
<b>Interaction</b>					
N X P	NS	*	*	NS	NS
N X V	**	**	**	**	NS
P X V	NS	*	NS	**	NS
N X P X V	*	*	*	*	*

Means with the same letters in a column under a treatment are not statistically different ( $P > 0.05$ ). Significant at ( $P < 0.05$ ), \*\* highly Significant at ( $P < 0.01$ ), NS (not significant)



**Interactive effect of nitrogen sources, proximity to homestead and varieties on shoot dry weight of cowpea in soils from Mashegu**

Kanannado variety treated with nitrogen in soils sampled further away from homestead had the heaviest shoot dry weight. These were however similar to the shoot dry weight produced by the same variety treated with BR 3262 or BR 3267 for soils close to homestead using IT90K-277-2 (Fig 1).

The smallest shoot biomass was produced by IT93K-452-1 plant in soils further away from homestead treated with BR 3262. A similar trend was observed using the same inoculant and proximity to homestead for IT90K-277-2 also IT93K-452-1, IT90K-277-2, and IT97K-499-35 that were either not treated with N or inoculated or those treated with either inoculant strains in soils sampled further away from homestead.

**Interaction of nitrogen sources, proximity to homestead and varieties on shoot dry weight of cowpea in soils from Mariga**

Fig 2 shows that nitrogen sources, proximity to homestead, and varieties of cowpea had a significant ( $P < 0.05$ ) positive effect on shoot dry weight of cowpea. The N – treated

kanannado plants grown on soils sampled close to homestead produced the largest biomass yield. The N – treated kanannado plants in soils further away from homestead had similar yield to those of BR 3267 inoculated IT97K-499-35 plant using soils sampled away from homestead. The least biomass was produced generally by the various varieties with no amendments.

**Interactive effect of nitrogen sources, proximity to homestead and varieties on shoot dry weight of cowpea in soils from Kontagora**

Kanannado plant irrespective of proximity to homestead treated with nitrogen had the heaviest shoot dry weight (Fig 3). These were however similar to the yield produced by the N-treated IT90K-277-2 plant using soils sampled close to homestead and BR 3262 inoculated Kanannado plant in soils sampled from both proximities to homestead. The least biomass was produced generally by IT90K-277-2 plant in soils of both proximities to homestead that were treated with inoculants or the unamendment plant. These was however not the case for the urea treated plant where IT97K-499-35 plant had the lowest shoot dry weight.




Fig 1

Interaction effect of nitrogen sources, proximity to homestead and varieties on shoot dry weight of cowpea in soils from Mashegu






Fig 2 Interaction effect of nitrogen sources, proximity to homestead and varieties on shoot dry weight of cowpea in soils from Mariga




Fig 3 Interaction effect of nitrogen sources, proximity to homestead and varieties on shoot dry weight of cowpea in soils from Kontagora



## DISCUSSION

Organic carbon of the experimental soil was low also the nitrogen content was much lower than the critical level of 15 g Kg<sup>-1</sup> (Okalebo *et al.*, 1993). The available phosphorus result indicate that Kontagora far away from homestead and Mariga close to homestead soils fall within the optimum available P concentration that would help enhance nodulation and Nitrogen fixation in the grain legume (Enwezor, 1990). Results of ECEC confirm that the experimental soil is low in fertility.

In this study, proximity of fields close to homesteads had significantly higher shoot dry weight than those away from homestead similar reports was made by Zingore *et al.* (2007) whose study shows that the fertile plots is often closest to homesteads, as a result of continuous accumulation of organic ammendment including all kinds of manure and household waste applied directly surrounding the villages.

The success of rhizobia inoculation primarily depends on rhizobial strain, the legume genotype, environmental conditions, and the crop management (Woomer *et al.*, 2014). Although, there are two main situations where there is likely to be a response to rhizobia inoculation; where compatible rhizobia of the host legume are absent and where native rhizobia population is low. In this study, counts of native rhizobia population were very high (>10 cells g<sup>-1</sup> soil). Suggesting that there may likely be no response to inoculation in these soils. Contrary to this observation, response to inoculation was observed in this study. This result agrees with the findings of Giller (2001) who report in his findings that the presence of large population density of compatible rhizobia does not, however, preclude the possibility that responses to inoculation can be obtained if competitive and highly effective strains are introduced in high quality inoculants. Similar results were obtained which could be due to the effectiveness of the elite strain introduced. This study shows that Kanannado variety had better response to inoculation than the other

varieties used. Also, Interaction between nitrogen sources, proximity to homestead and varieties significantly affect the shoot dry weight.

Result from this study suggests that the two elite inoculant strains used for this study could establish effective symbiosis with cowpea in soils of the study area. Thereby, increasing the yield of cowpea plants. However, in spite of the beneficial effect of elite rhizobia strains, shoot dry weight of the N-treated plant was marginally higher than those of the elite strains thus unable to meet the optimum nitrogen requirement of the plant. Further study could however be carried out to evaluate greater number of inoculant strains to further get to the conclusion on the efficacy of inoculation on cowpea which could serve as a cheaper alternative to inorganic N-fertilizer application.

## REFERENCES

- Abaidoo, R.C., Keyser, H.H., Singleton, P.W., Dashiell, K.E. & Sanginga, N. (2007). Population size, distribution, and symbiotic characteristics of indigenous Bradyrhizobium spp. that nodulate TGx soybean genotypes in Africa. *Applied Soil Ecology* 35: 57-67.
- Aliyu, I., Yusuf, A. & Abaidoo, R. (2013). Response of Grain Legumes to Rhizobial Inoculation in Two Savanna Soils of Nigeria. *African Journal of Microbiology Research*, 7, 1332-1342.
- Bagayoko M, Buerkert A, Lung G, Bationo A, & Romheld V. (2000) Cereal/legume rotation effects on cereal growth in Sudano-Sahelian West Africa: soil mineral nitrogen, mycorrhizae and nematodes. *Plant Soil*, 2000; 218: 103-116.
- Enwezor, W.O., Udo, E.J. Ayotade K.A., Adepetu J.A and. Chude. V.O (1990). and Natural resources, Lagos, pp. 241-279.
- Giller, K.E. (2001). Nitrogen fixation in tropical cropping systems. CAB International, Walliford, UK.



- Giller, K.E., Rowe, E., de Ridder, N., & van Keulen, H., (2005). Resource use dynamics and interactions in the tropics: scaling up in space in time. Agricultural System, in press.
- GRDC (Grains Research and Development Corporation), Australia. 2013. Rhizobial inoculants factsheet. Available at [www.coretext.com.au](http://www.coretext.com.au). January, 2013. <http://www.grdc.com.au/~media/B943F697AF9A406ABBA20E136FDB7DC4.ashx> Accessed on 24/9/2013.
- Howieson, J. & Ballard, R. (2004). Optimizing the legume symbiosis in stressful and competitive environments within southern Australia - some contemporary thoughts. *Soil Biology and Biochemistry* 36, 1261-1273.
- IITA (International Institute of Tropical Agriculture). (1989). Selected methods for soil and plant analysis; Manual Series, No. 1 Revised Edition, 1979. IITA, Ibadan, Nigeria. 69.
- Okalebo, J.R., Gathua, K.W., & Woomer, P.L. (1993). Laboratory methods of Soil and Plant Analyses: A Working Manual TSBF UNESCO-ROSTA. pp 22-25.
- Sandman, W. P. L. (1970). Practical aspects of Rhizobium bacteriology. Rhodesian Ministry of Agriculture. 185.
- Vincent, J. M. (1970). A manual for the practical study of root nodule bacteria. International Biological programme handbook no 15 Blackwell Scientific Publ Ltd, Oxford, United Kingdom.
- Woomer, P. L. (1994). Most Probable Counts. In Methods of soil analysis, Part 2. Microbiological and Biochemical Properties.
- Woomer, P. L., Huising, J. & Giller, K.E. (2014). *N2Africa Final Report of the First Phase 2009-2013*. Available at: [www.N2Africa.org](http://www.N2Africa.org) 138.