

## BACTERIOLOGICAL ASSESMENT OF BEEF CATTLE SLAUGHTERED IN MINNA METROPOLIS.

J.Y.ADAMA, I.C.ALEMEDE, A.MISAN AND S. OGUNBAJO

Department of Animal Production, Federal University of Technology P.M.B. 65 Minna, Nigeria.

E-mail: adama.live@yahoo.com

### ABSTRACT

This study was conducted to establish the bacterial status of beef cattle slaughtered in minna metropolis. Five (5) abattoir/slaughter slabs were selected for the study, they include; Bosso, maitumbi, Chanchaga, Tunga and Maikunkele. A total of fifty (50) fresh beef sample were collected for a period of ten (10) weeks and analysed for total viable bacterial counts. Four (4) bacteria that were characterized and identified are *Bacillus subtilis*, *staphylococcus aureus*, *streptococcus pneumonia* and *streptococcus faecalis*. The total viable counts ranged from  $9.0 \times 10^7$  -  $4.8 \times 10^7$  CFU/g, the mean viable bacterial counts for the various locations includes;  $1.03 \times 10^6 \pm 2.36 \times 10^6$ ,  $1.55 \times 10^5$ ,  $0.08 \times 10^5$ ,  $5.11 \times 10^5 \pm 1.21 \times 10^5$ ,  $1.48 \times 10^6 \pm 3.12 \times 10^6$  and  $7.19 \times 10^5 \pm 2.19 \times 10^5$  for Bosso, Maitumbi, Chanchaga, Tunga and maikunkele respectively. The results revealed high bacterial counts and significant differences. Tunga was revealed to have the highest mean bacterial count of  $1.48 \times 10^6$  CFU/g, while Maitumbi had the least mean bacterial count of  $1.55 \times 10^5$  CFU/g, it is recommended that the quality control system in use at the various abattoir/slaughter slabs be improved upon.

### INTRODUCTION

Beef is consumed in almost every part of the world. It generally has no religious taboos attached to it and so it is accepted by both the young and the old, the rich and the poor, and since it is a major source of protein, it is very important to consumers.

However, the high nutritive nature of fresh beef, along with a favourable water activity and appropriate PH, allows a vast variety of bacteria to grow on it (Kibicki, 1995). In Nigeria, there are so many problems associated with bacterial content of meat. These problems include: pre-harvest and harvest source of meat contamination, transport system of meat and meat products. Processing and storage of meat and meat products in most cases, these meats after slaughter are transported on motorcycles with the rider carrying some of the meat on his body and some dropping on the ground.

Meat quality is dependent on the entire meat production chain, from where the animals are processed to the consumer. It covers sensory and microbiological properties (monin 1991). Therefore, this study is designed to establish the bacteriological status of beef cattle slaughtered in Minna metropolis.

### MATERIALS AND METHODS

#### Location of Study Area

This was carried out in Minna environs. Minna has a land mass of 228.5km square and lies between longitude  $6^{\circ}29'E$  and latitude  $9^{\circ}31'N$  of

the equator. The average temperature ranges between  $18-39^{\circ}C$  with an average monthly rainfall of 120mm (student handbook, 2008).

#### Methods of Collection of Samples

Five (5) abattoir/slaughter slabs were selected for this study. They include; Bosso modern abattoir, Mainkunkele, Tunga, Maitumbi and Chanchaga slaughter slabs. These abattoir/slaughter slabs were selected because they are the major abattoir/slaughter slabs where cattle are slaughtered in Minna on daily basis. A sample of fresh beef was collected from each of the abattoir/slaughter slabs once a week making a total of five (5) samples a week. They were collected in the morning and wrapped with foil papers to prevent contamination from external factors. The duration for the study was ten (10) weeks, making a total of fifty (50) sample used for the study. The sample collected were taken to the Department of Microbiology Laboratory, Federal University of Technology, Minna for bacteriological examination using Gram stain technique and enumeration of bacteria through the use of colouring counting plate.

The results obtained were subjected to statistical analysis using the SAS system involving the Analysis of Variance (ANOVA) procedure.

### RESULTS AND DISCUSSION

Table 1.0 indicates the bacterial counts obtained from the agar plate counts. The total viable bacteria count in the sample were analysed and expressed as Log 10 CFU/g (colony forming

units per gram). The result of the total viable count of bacteria load of the fresh beef samples revealed that the samples had high bacteria counts ranging from  $9.0 \times 10^5 \pm 0.08 \times 10^5$  to  $4.8 \times 10^6 \pm 0.08 \times 10^6$  CFU/g. These values exceed the bacteria infective dose level of  $10^5$  CFU/g as stated by South African Department of Health (2000). Similarly, Nortje et al. (1995) from the outcome of their study, suggested that total viable count in the regions of  $10^7$  CFU/g in meat is regarded as the upper limit of acceptability and that such result suggest a deterioration of hygienic standards.

The result also revealed that the location having the highest bacteria count was Tunga, with a mean bacteria count of  $1.48 \times 10^6$  CFU/g, while Maitumbi had the least count of  $1.55 \times 10^5$  CFU/g.

The bacteria isolate were fully characterized and identified as species of staphylococcus aureus, Bacillus subtilis, strephylococcus pneumonia, and staphylococcus faecalis as shown in table 2.0. The presence of this bacteria isolate could suggest po

Hygienic and working practices of the meat handlers during the processing stage as well as lack of sterilization of utensils and working surfaces which confirms earlier report (Atanassova et al. 2001) that humans have often been reported to be carrier of staphylococci. According to (Jay, 1996), the first contamination is usually caused by non-sterilized knives and surface at slaughtering. Streptococcus faecalis as found in some of the sample analysed is an indication of faecal pollution of some of the fresh beef samples. The bacteria are of zoonotic importance, in that they have the ability to cause food borne infection when consumed by

humans. For instance, staphylococcus aureus has ability to cause boil (skin infection) as well as respiratory and urinary tract infections. It can also cause infertility. Infection caused by Bacillus subtilis in characterized by abdominal pains, profuse watery diarrhea and rectal tenesmus.

#### CONCLUSION

The results of this study has shown that beef cattle slaughtered in various abattoir/slaughter slabs in Minna metropolis had high bacterial counts exceeding the bacterial infective dose level, indicating that the sanitary procedures might have been compromised

#### ACKNOWLEDGEMENT

I wish to acknowledge the technical assistance given by the technical staff of the microbiology laboratory of the Department of Microbiology, Federal University of Technology, Minna.

#### REFERENCES

- Atanassova, R.N., Scheller, K.K and Arp, S.C. (2001). Bacterial growth in ground beef patties. *Food protection* 31: 36-40.
- Jay, C.M. (1996). Detection of the microbial spoilage of fresh beef. *Food science technol.* 12: 414-424
- Krockel, J.M. (1996). Microbial contamination of meat and cured meat products. *International journal for food microbial* 33: 103-113
- Monin, E.T. and Quali, I.M. (1991). Sensory analysis and microbiological properties of fresh beef. *Meat science* 10: 345-355
- Nortje, R.T., Hill, D.J and Dodd, C.E. (1999). Bacterial Community Structure and Location in Fresh meat. *Meat science* 73: 103-114
- South African Department of Health (2000). Bacteriological levels in fresh meat. *International meat science*. 61: 542-549

Table 1. Total Viable Count of Bacteria Isolate from Fresh Beef Samples from Selected Abattoir/slaughter Slabs in Minna Metropolis (CFU/g).

Variable	Location			level of Significance	RS <sup>a</sup> / DL
	Chanchaga	Tunga	Makunkele		
1.5 x 10 <sup>5</sup>	4.2 x 10 <sup>5</sup> ± 0.14 x 10 <sup>5</sup>	5.2 x 10 <sup>5</sup> ± 1.08 x 10 <sup>5</sup>	2.5 x 10 <sup>5</sup> ± 0.05 x 10 <sup>5</sup>	*	10 <sup>5</sup>
2 x 10 <sup>5</sup>	5.3 x 10 <sup>5</sup> ± 0.13 x 10 <sup>5</sup>	4.8 x 10 <sup>5</sup> ± 0.06 x 10 <sup>5</sup>	2.9 x 10 <sup>5</sup> ± 0.04 x 10 <sup>5</sup>	*	10 <sup>5</sup>
3 x 10 <sup>5</sup>	6.4 x 10 <sup>5</sup> ± 0.14 x 10 <sup>5</sup>	6.0 x 10 <sup>5</sup> ± 0.09 x 10 <sup>5</sup>	1.2 x 10 <sup>5</sup> ± 0.07 x 10 <sup>5</sup>	*	10 <sup>5</sup>
4 x 10 <sup>5</sup>	7.5 x 10 <sup>5</sup> ± 0.20 x 10 <sup>5</sup>	1.4 x 10 <sup>5</sup> ± 0.08 x 10 <sup>5</sup>	5.2 x 10 <sup>5</sup> ± 0.08 x 10 <sup>5</sup>	*	10 <sup>5</sup>
5 x 10 <sup>5</sup>	8.6 x 10 <sup>5</sup> ± 0.21 x 10 <sup>5</sup>	8.0 x 10 <sup>5</sup> ± 0.07 x 10 <sup>5</sup>	9.3 x 10 <sup>5</sup> ± 1.01 x 10 <sup>5</sup>	*	10 <sup>5</sup>
6 x 10 <sup>5</sup>	9.7 x 10 <sup>5</sup> ± 0.22 x 10 <sup>5</sup>	1.2 x 10 <sup>5</sup> ± 0.08 x 10 <sup>5</sup>	1.5 x 10 <sup>5</sup> ± 0.13 x 10 <sup>5</sup>	*	10 <sup>5</sup>
7 x 10 <sup>5</sup>	1.08 x 10 <sup>5</sup> ± 0.23 x 10 <sup>5</sup>	2.0 x 10 <sup>5</sup> ± 1.08 x 10 <sup>5</sup>	1.8 x 10 <sup>5</sup> ± 0.10 x 10 <sup>5</sup>	*	10 <sup>5</sup>
8 x 10 <sup>5</sup>	1.19 x 10 <sup>5</sup> ± 0.24 x 10 <sup>5</sup>	3.0 x 10 <sup>5</sup> ± 0.10 x 10 <sup>5</sup>	4.7 x 10 <sup>5</sup> ± 0.06 x 10 <sup>5</sup>	*	10 <sup>5</sup>
9 x 10 <sup>5</sup>	1.30 x 10 <sup>5</sup> ± 0.25 x 10 <sup>5</sup>	1.3 x 10 <sup>5</sup> ± 0.09 x 10 <sup>5</sup>	1.3 x 10 <sup>5</sup> ± 0.55 x 10 <sup>5</sup>	*	10 <sup>5</sup>
10 x 10 <sup>5</sup>	1.41 x 10 <sup>5</sup> ± 0.26 x 10 <sup>5</sup>	4.8 x 10 <sup>5</sup> ± 0.19 x 10 <sup>5</sup>	1.7 x 10 <sup>5</sup> ± 0.75 x 10 <sup>5</sup>	*	10 <sup>5</sup>
11 x 10 <sup>5</sup>	1.52 x 10 <sup>5</sup> ± 0.27 x 10 <sup>5</sup>	1.4 x 10 <sup>5</sup> ± 0.10 x 10 <sup>5</sup>	7.9 x 10 <sup>5</sup> ± 2.19 x 10 <sup>5</sup>	*	10 <sup>5</sup>

Significant superscript letters (P < 0.05) different

RS<sup>a</sup> = Resistant Significance (P < 0.05)

Table 2.0 Characteristics and Identification of Bacteria Isolated from Fresh Beef Samples Collected from the Various Abattoir slaughter Slabs in Minna Metropolis.

Location	Gram's reaction	Catalase test	Coagulase test	Starch Hydrolysis test	Hydrogen sulphide	Methyl red test	Voges proskauer test	Oxidase test	Motility test	Indole test	Organism
Iosso	+c	+	+	-	-	-	-	-	-	-	<i>Staphylococcus aureus</i>
	+c	-	-	-	-	-	-	-	-	-	<i>Streptococcus faecalis</i>
	+c	-	-	-	-	-	-	-	-	-	<i>Streptococcus pneumoniae</i>
Iaitumbi	+R	+	-	+	-	-	+	-	+	-	<i>Bacillus subtilis</i>
	+c	-	-	-	-	-	-	-	-	-	<i>Streptococcus faecalis</i>
	+c	-	+	-	-	-	-	-	-	-	<i>Streptococcus aureus</i>
Ihanchaga	+c	+	+	-	-	-	-	-	-	-	<i>Staphylococcus aureus</i>
	-R	+	-	+	-	-	+	-	+	-	<i>Bacillus subtilis</i>
	+c	+	+	-	-	-	-	-	-	-	<i>Staphylococcus aureus</i>
Iunga	+c	-	-	-	-	-	-	-	-	-	<i>Streptococcus faecalis</i>
	+c	-	-	-	-	-	-	-	-	-	<i>Streptococcus pneumoniae</i>
	+R	+	-	+	-	-	+	-	+	-	<i>Bacillus subtilis</i>
Iaikunkele	+c	+	+	-	-	-	-	-	-	-	<i>Staphylococcus aureus</i>
	+c	-	-	-	-	-	-	-	-	-	<i>Streptococcus faecalis</i>

+R = positive rod shape      - = Negative Result

+C = positive cocci shape      += Positive Result



# GUINEA FEED... Pelletize

For poultry farmers. Our Layers are assured for better performance

## PELLETIZED GUINEA FEED



- Special formulated and Technically produced for Higher overall performance.
- Highly Palatable.
- Maximum Profit on investment.

## FOR BEST RESULT



### FEED TO LAYERS FROM 18TH WEEK OF AGE

- Feed on bag to 200 Birds per day.
- Give adequate hygienic water.
- Ensure ideal ventilation.
- Avoid Bird overcrowding
- Provide good farm management

## GUARANTEED



- Faster growth rate
- Improve feed conversion
- High production depending on age Birds
- Eliminate wet Litter problem resulting in a healthier environment
- Bigger egg size and more weight more money

*Benefit from our scientific know - how!!!*



# FISH FEED... Pelletize

For fingerlings, Juvenile and adult

RC. No 57274

**Bendel Feed & Flour Mill Limited**

Factory: Km 100, Benin - Auchu Road, Ewu, Edo State. Tel: 08035357474