



# INTERNATIONAL JOURNAL OF APPLIED BIOLOGICAL RESEARCH

VOLUME 4, NUMBER, 1 & 2,  
DECEMBER, 2012.



**IJABR**

ISSN 2141 - 1441

Editor-in-Chief

# INTERNATIONAL JOURNAL OF APPLIED BIOLOGICAL RESEARCH

## EDITORIAL BOARD

Editor - in - Chief	Prof. H. A. Suberu	Dean, School of Natural and Applied Sciences, Federal University of Technology, Minna
Managing Editor	Dr. I. C. J. Omalu	Head, Department of Biological Sciences, Federal University of Technology, Minna
Secretary	Dr. I. K. Olayemi	Department of Biological Sciences, Federal University of Technology, Minna
Associate Editor	Prof. O. A. Falusi	Department of Biological Sciences, Federal University of Technology, Minna
Associate Editor	Prof. S. Alonge	Department of Biological Sciences, Ahamadu Bello University Zaria
Associate Editor	Prof. G. I. Imande	Department of Zoology, University of Jos
Associate Editor	Ass. Prof. O. T. Mustapha	Department of Plant Biology, University of Ilorin.
Associate Editor	S. O. Abolarinwa	Department of Biological Sciences, Federal University of Technology, Minna
Associate Editor	Dr. C. Mesmerembwa	African Institute of Biomedical Science and Technology, Harare, Zimbabwe
Associate Editor	Prof. M. Galadima	Department of Micrbiology, Federal University of Technology, Minna
Consultant	Prof. A. A. Oladimeji	Department of Biological Sciences, IBB University, Lapai

A full list of editors will be published later.

	PAGES
The Role of Hep-2 Cell Adherent <i>Escherichia coli</i> Strains on Diarrheic Children in Imo State of Nigeria. Ene, P. C., Abbey, S. D., Madubuattah, C. G. and Ugwu, K. C.....	8
Potentials of Stem Cuttings of Four Selected Tree Species for the Establishment of Shelterbelt in the Semi-Arid Zone, Kano Nigeria. Mustapha, O. T. and Abubakar, A. A.....	15
Mutagenic Effects of Fast Neutron Irradiation on Selected Morphological Characters and Yield of the African Long Pepper ( <i>Capsicum annum</i> var. <i>accuminatum</i> ). Daudu O. A. Y., Falusi O. A., Dangana M. C., Thomas, T., Bello, I. M. and Muhammad, L. M.....	19
Proximate Composition and Post-Production Stability of Poultry Waste Fertilizer Pellets. Babatope, A., Musa, J. J. and Farouk, F.....	25
Influence of Initial Moisture Content on Some Proximate Quality Attributes of Packaged Gari in Storage. Adejumo, B. A.....	32
Essential amino acid composition of freshly fermented seeds and powdered condiments of <i>Prosopis africana</i> , <i>Parkia biglobosa</i> and <i>Glycine max</i> with organism's consortia and natural fermentation. Gberikon, G. M., Ameh, J. B., Ado, S. A. and Umoh, V. J.....	39
Effects of Different Feeds on the Chromosome Behaviour of <i>Zonocerus variegatus</i> . Ogueji, E. O.....	45
Influence of Temperature and Relative Humidity on the Culture of Fungi Isolated from Rotting of Sweet Potato ( <i>Ipomoea batatas</i> (L.) Lam.). Bello, I. M., Suberu, H. A., Kuta, F. A. and Daudu, O. A. Y.....	51
Spatial Distribution of Lake-water Insects in Minna, North central Nigeria: Bio-indication of Water Quality. Mohammed, A. Z., Sidiq, O. L., Olayemi, I. K. and Ayanwale, A. V.....	57
Influence of Different Rates of Application of Poultry Dropping on the Growth and Yield of Tomato ( <i>Lycopersicum esculentum</i> L.) cultivars. Saidu . A., Bello, L. Y., Tsado, E. K. and Sani, A.....	65
Allometric Relationships in Field-Grown Annual Plant Species with Different Architectures as a Measure of Intraspecific Competition. Japhet, W. S., Aliyu, N. and Abdullahi, P. A.....	71
Screening Five Fungal Isolates as Potential Antagonists to <i>Phytophthora palmivora</i> on Cocoa Pod Pericarp. Adebola , M. O. and Amadi, J. E.....	78
Environmental Appraisal of Drainage System in Minna, Niger State-Nigeria. Musa, J. J.....	85
Effect of Sub-Lethal Doses of Lambda-Cyhalothrin on Leukocyte Sub-Population (Differential Count) of African Catfish <i>Clarias gariepinus</i> (Burchell, 1822). Ogueji, E. O. and Ibrahim, B. U.....	94

**Mutagenic Effects of Fast Neutron Irradiation on Selected Morphological Characters and Yield of the African Long Pepper (*Capsicum annum* var. *accuminatum*)**

Daudu O. A. Y\*, Falusi O. A., Dangana M. C., Thomas, T., Bello, I. M. and Muhammad, L. M.

Department of Biological Sciences, Federal University of Technology, Minna, Niger State, Nigeria.

Submitted: April 7, 2012; Accepted: Nov. 12, 2012; Published: Dec. 17, 2012.

**ABSTRACT**

The effects of fast neutron irradiation using an Americium-Beryllium source with a flux of  $1.5 \times 10^4 \text{ n cm}^{-2} \text{ s}^{-1}$  on the morphological and yield traits of African long pepper (*Capsicum annum* var. *accuminatum*) were studied. The pepper seeds were irradiated for 0, 30, 60, 90 and 120 minutes before they were sown, with their respective controls, in order to assess the effects of the different irradiation treatments on percentage survival, days to maturity, plant height, number of fruits per plant, number of seeds per fruits, height of fruit, weight of fruit and width of fruit in the  $M_1$  generations. All irradiation periods caused leaf abnormalities such as leaves with invaginated or inverted margins, or with a blunt or bifurcated apex, or leaves that turned bifoliage when compared with Control plants. There was, also, pronounced variation in germination percentage, plant height, number of leaves/plant, fruits/plant, seeds/fruit, weight of fruit, length and width of fruits; all essentially may affect productivity negatively. Moreover, 120 min was an effective irradiation period to induce viable and useful mutations for yield parameters in pepper. These results suggest the possibility of evolving higher yield variants of pepper through proper selection.

**Key words:** Americium-Beryllium,  $M_1$  generations, Leaf Abnormalities, Selection

\*Corresponding Author: dauduoladipupoyusuf@yahoo.com

**INTRODUCTION**

*Capsicum* belongs to the nightshade Family, Solanaceae (GRIN, 2009). The genus consists of over 100 species and even more botanical varieties (Ado, 1999; Falusi, 2007), including five domesticated species (i.e. *C. annum*, *C. frutescens*, *C. baccatum*, *C. chinense* and *C. pubescens*) all believed to have originated from the New World (Bosland, 1994). *C. annum* and *C. frutescens* are the most recognized species grown in commercial quantities all over Nigeria (Falusi and Morakinyo, 2001; Mady *et al.*, 2005). These two species form an important ingredient in people's diet around the world (GRIN, 2009), due to the pungency properties of the fruits resulting from their high concentration of capsaicinoid alkaloid (Bosland and Vostava, 2000). In addition, *Capsicum* is a

rich source of vitamins A and C (Gill, 1992; Ado, 1999). *Capsicum* fruits are also popular as food spices, as a colouring agent and serve as pharmaceutical ingredients (Bosland, 1996). In African medicine, it is used to treat sore throats (Abdullahi *et al.*, 2003).

Mutations, which result in alteration of the genetic information of a cell or living creature are the tools used to study the nature and function of genes which are the building blocks and basis of plant growth and development, thereby producing raw materials for genetic improvement of economic crops (Adamu and Aliyu, 2007). Mutation technology has been used to produce many cultivars with improved economic value and to advance the study of genetics and plant developmental phenomena (Bertagne-Sagnard *et al.*, 1996;

Poornananda and Hosakatte, 2009). Genetic variability for desired characters can be induced successfully through mutations, with high practical value in plant improvement programs (Fahad and Salim, 2009).

Induced mutations have been used to generate genetic variability for the improvement of yield components of various crops like *Oryza sativa* (Awan *et al.*, 1980; Singh *et al.*, 1998), *Hordeum vulgare* (Ramesh *et al.*, 2001), *Triticum durum* (Sakin and Yildirim, 2004), *Cicer arietinum* (Wani and Anis, 2001), *Vigna mungo* (Misra *et al.*, 2001), *Helianthus annuus* (Elangovan, 2001) *Cajanus cajan* (Ravikesavan *et al.*, 2001), *Sesame indicum* (Mensah *et al.*, 2007), *Guizotia abyssinica* (Misra, 2001). These reports show that mutagenesis is a potential tool to be employed for crop improvement.

The FAO (2009) reported that 2008 marked the 80<sup>th</sup> anniversary of mutation induction in plants. The application of gamma rays and other physical mutagens such as fast neutrons has generated a vast amount of genetic variability and has played a significant role in plant breeding and genetic studies (David, 2010). The widespread use of induced mutants in plant breeding programmes throughout the world has led to the official release of more than 2700 plant mutant varieties (FAO 2009). The present work aimed to investigate the growth responses and the yield of the African Long Pepper (ALP) to different doses of Fast Neutron Irradiation (FNI).

## MATERIALS AND METHODS

### Description of African Long Pepper

Medium-sized annual plant with a long pointed and pendant fruits with hot taste; it produces one pedicel per node.

Fifty fresh fruits of ALP accessions were bought from a local farmer in Minna (North Central geopolitical zone of Nigeria, found within longitude 6°34' East and latitude 9°36' North), Niger State, Nigeria. The fruits were kept in a clean polythene bag. The

accessions were identified as *Capsicum annum* var. *abbreviatum* Fingerh using taxonomic aid provided by Simmond (1976), as well as morphological descriptions of Hutchinson and Dalziel (1963), Schippers (2000) and Abdullahi *et al.*, (2003). Each fruit of the ALP accessions was cut open; the seeds removed, and separately sun-dried for 8 h. The sun-dried seeds were tested for viability using the floatation method before FNI treatment. The dry seeds of *Capsicum* were irradiated at the Centre for Energy and Research Training (CERT), Ahmadu Bello University, Zaria with FNI using an Americium-Beryllium source with a flux of  $1.5 \times 10^4$  n cm<sup>-2</sup> s<sup>-1</sup> for five different irradiation exposure periods (IEPs): 0, 30, 60, 90, and 120 min. The equipment used was a Miniature Neutron Source Reactor (MNSR) designed by the China Institute of Atomic Energy (CIAE) and licensed to operate at a maximum power of 31 kW (Sar, 2005).

Treated seeds (100 from each treatment) were then planted in nursery trays to obtain seedlings, which were transplanted into 3.5 litres plastic pots containing garden soil, at a rate of three seedlings/pot, after 4 weeks in the nursery. No fertilizer was applied although, when the crop began to flower, an insecticide was applied to prevent insect-borne diseases. The planted seeds were watered once daily between 5.00-6.30 pm using bore-hole water. Each treatment was replicated four times, using a completely randomized design (CRD).

Data were collected from 15 plants for each accession for germination percentage, number of leaves/plant at maturity, height of plant at maturity (number of days to 50% flowering) and yield/plant in each M1 (M = first filial mutant) generation. Data was analyzed using analysis of variance (ANOVA) and Least Significant Difference test was used to separate the means with significant differences detected at  $P = 0.05$ . Pearson's correlation analysis was used to find the relationship between treatments and selected parameters.

## RESULTS AND DISCUSSION

There was a negative strong coefficient of correlation between irradiation period and seed survival percentage in MN/SH/001 (Table 1), implying that as irradiation time increased, percentage survival decreased. Wide variation was observed in plant height, number of leaves/plant, fruits/plant, and seeds/fruit, and weight, length and width of fruits for the ALP accessions (Table 2). FNI is capable of producing significant changes in the morphological and yield parameters of the pepper plant. Similar effects of ionizing radiation on reproductive and other yield parameters has been reported for tomato exposed to Sodium azide with a concentration between 1 and 4 mM (Adamu and Aliyu, 2007) and also for Okra exposed to Gamma irradiation doses between 300 and 500 Gray (Hegazi and Hamideldin, 2010). Asmahan and Nada (2006), Fahad (2009) and Hegazi and Hamideldin (2010) reported that an increase in irradiation dose tended to increase certain morphological traits such as plant height.

FNI for 120 min could effectively induce viable and useful mutations in pepper yield parameters. A similar result was reported by Reddy and Smith (1978) and Adamu *et al.* (2004), who reported that obtaining

mutations in the crops they studied (*Sorghum bicolor* and *Lycopersicon esculentum*, respectively) using ionizing radiation were dose dependent.

All the plants produced from the non-irradiated seeds which served as the control produced normal leaves which had pointed apices and entire margin (Fig. A). However, this was not the case among the ALP plants whose seeds were irradiated. Leaf irregularities such as leaves with invaginated or inverted margins, or with a blunt or bifurcated apex, or leaves that turned bifoliage were observed. These observed leaf morphological abnormalities are indications that FNI affected the plants negatively. Shah *et al.* (2008), Khan and Aslam (1982) and Islam *et al.* (1994) reported similar leaf morphological abnormalities in *Crotalaria saltiana* and *C. juncea* treated with ionizing radiation. Low genetic variability for major characters is a limiting factor for the improvement of a crop but induced mutations can provide an additional source of variability for quantitative and qualitatively inherited traits in a number of crop plants. FNI can be used to increase selected growth and yield characters and induce beneficial variability in peppers which can be incorporated in conventional breeding for the improvement of the crop.

Table 1: Correlation (r) between the treatments and percentage of seeds that germinated per 100 seeds sown for three pepper (*Capsicum* spp.) accessions.

Treatments	ALP/%
Control	100
30 min	50
60 min	51
90 min	51
120 min	37
R	-0.8156

ALP: African Long Pepper

Table 2: LSD of the effects of Fast neutron irradiation on morphological and yield traits of ALP accessions.

Characters	Productivity in ALP during the Irradiation periods (min)				
	0	30	60	90	120
ALP	44.20 ± 5.19 a	45.27 ± 11.52 a	52.86 ± 4.93 ab	50.23 ± 14.13 ab	59.93 ± 6.00 b
Plant height (cm)	129.93 ± 10.37 a	132.70 ± 44.63 a	173.00 ± 36.45 a	161.80 ± 62.46 a	230.80 ± 32.78 b
Number of leaves/plant	8.00 ± 2.48 a	15.30 ± 4.90 b	14.80 ± 3.79 b	14.30 ± 4.99 b	15.80 ± 4.94 b
Number of fruits/plant	102.80 ± 18.21 a	105.00 ± 21.65 ab	108.10 ± 22.20 ab	110.30 ± 23.77 ab	125.80 ± 0.45 b
Number of seeds/fruit	7.26 ± 1.41 a	7.54 ± 1.70 a	6.94 ± 1.50 a	7.67 ± 1.49 a	8.20 ± 1.71 a
Length of fruit (cm)	1.93 ± 0.29 a	1.97 ± 0.21 abc	1.90 ± 0.26 ab	2.15 ± 0.18 bc	2.28 ± 0.45 c
Width of fruit (cm)	7.7 ± 0.67 a	8.40 ± 2.01 a	8.70 ± 1.84 a	9.10 ± 1.10 ab	10.2 ± 1.48 b
Weight of fruit (g)					

Values are mean ± SD. Values followed by the same letter(s) within the same row do not statistically differ at the 5% level according to LSD




Plate 1: A. Normal leaf of *C. annuum* var *accuminatum* B. A leaf with another leafy outgrow at the petiole (90 min IEP) C. Leaves showing invaginated margins (120 min IEP) D. The leaf that turned bifoliage (30 min IEP) E. Leaf with bifurcated apex (60 min IEP).

REFERENCES

Abdullahi, M., Muhammad, G. and Abdulkadir, N. U. (2003). *Medicinal and economic plants of Nupeland*. Jube-Evans Books and Publications, Bida, Nigeria, 276 pp.

Adamu, A. K., Clung, S. S. and Abubakar, S. (2004). Effects of ionizing radiation (gamma-rays) on tomato (*Lycopersicon esculentum* L.). *Nigerian Journal of Experimental and Applied Biology*, 5(2): 185-193.

Adamu, A. K. and Aliyu, H. (2007). Morphological effects of Sodium azide on tomato (*Lycopersicon esculentum* Mill.). *Science World Journal*. 2(4): 9-12.

- Ado, S. G. (1999). Potentials of native and exotic pepper germplasm in Nigeria: An exploitable resource in the next millennium. *Commemorative publication on the Silver Jubilee of the Genetic Society of Nigeria*. pp. 22-36.
- Asmahan, A. M. and Nada, A. (2006). Effect of gamma irradiation and Sodium azide on some economic traits in Tomato. *Saudi Journal of Biological Science*, 13(1): 44-49.
- Awan, M. A., Konzak, C. F., Rutger, J. N. and Nilan, A. (1980). Mutagenic effects of Sodium azide in rice. *Crop Science*, 20: 663-668.
- Bertagne-Sagnard, B., Fouilloux, G. and Chupeau, Y. (1996). Induced albino mutations as a tool for genetic and cell biology in flax (*Linum usitatissimum*). *Journal of Experimental Botany*, 47: 189-194.
- Bosland, P. W. (1996). *Capsicum*: Innovative uses of an ancient crop. In: *Progress in New Crops*, Janick, J. (ed.) ASHS Press, Arlington, VA, pp. 479-87.
- Bosland, P. W. (1994). Chiles: history, cultivation, and uses. In: Charalambous, G. (ed.), (*spices, herbs and edible fungi*). Elsevier Publ., New York, pp. 347-366.
- Bosland, P. W. and Vostava, E. J. (2000). *Peppers: Vegetable and Spice Capsicum*. CABI Publishing, Wallingford, UK, UK. 204 pp.
- David, T. (2010). All you wanted to know about induced mutations in crop breeding. *Bulletin of Biofortified*. <http://www.webster@ext.msstate.edu>
- Elangovan, M. (2001). Gamma radiation induced mutant for improved yield components in sunflower. *Mutat. Breed. Newslett.* 45: 28-29.
- Fahad, A. (2009). Effects of Sodium azide on growth and yield traits of *Eruca sativa* (L.). *World Applied Science Journal*, 7(2): 220-226.
- Fahad, A. and Salim K. (2009). Mutagenic effects of Sodium azide and its application in crop improvement. *World Applied Science Journal*, 6 (12): 1589-1601.
- Falusi, O. A. (2007). Germplasm collection of peppers (*Capsicum* spp.) in Nigeria. *Res. Crops* 8(3), 765-768.
- Falusi, O. A. and Morakinyo, J. A. (2001). Pollen and hybridization studies in some Nigerian species of peppers. *Nigerian Journal of Technology & Education*, 1 & 2: 40-43.
- Food and Agricultural Organization. (2009). *Induced plant mutations in the genomic era*. Publication of the Food and Agricultural Organization of the United Nations (FAO) Rome.
- Germplasm Resources Information Network (GRIN). (2009). *Capsicum* L. United States Department of Agriculture. <http://www.ars.grin.gov/cgi> (last accessed 21 August, 2011).
- Gill, L. S. (1992). *Ethnomedicinal uses of plants in Nigeria*. Uni. Ben. Press Benin City, 276pp.
- Hegazi, A. Z. and Hamildeldin, N. (2010). The effect of gamma irradiation on enhancement of growth and seed yield of okra [*Abelmoschus esculentus* (L.) Moench]] and associated molecular changes. *Journal of Horticulture and Forestry*, 2(3), 38-51.
- Hutchinson, J. and Dalziel, J. M. (1963). *Floral of West Tropical Africa II* Crown Agents, London, 533 pp.
- Islam, S. M. S., Kabir, G., Shahiquzzaman, G. M. and Khan, M. R. (1994). Effects of gamma rays and temperature treatment on germination, plant height and morphological abnormalities of *Crotalaria juncea* L. *Journal of Biological Science*, 2: 31-37.



Khan, M. A. and Aslam, S. (1982). Effects of gamma rays on the morphology and growth of *Crotalaria juncea* and *C. sericea*. *Journal of Asiatic Society Bangladesh, (SC)*: 8: 71-77.

Mady, E. A., Uguru, M. I. and Ugwoke, K. I. (2005). Interrelations of growth and disease expression in pepper using principal component analysis. *Proc. 30<sup>th</sup> Annual National Conference of Genetic Society of Nigeria*. pp. 55-59.

Mensah J. K., Obadoni, B. O., Akomeah, P. A., Lkhajiagbe, B., and Ajibolu J. (2007). The effect of sodium azide and colchicine treatments on morphological and yield traits of sesame seed (*Sesame indicum* L.). *African Journal of Biotechnology*, 6(5): 534-538.

Misra, R. C., Mohapatra, B. D., and Panda, B. S. (2001). High yielding mutants of blackgram variety PH-25. *Mutat. Breed. Newslett.*, 45: p. 39.

Misra, R. C. (2001). Productive mutants of niger. *Mutation Breeding Newsletter*, 45: 42-43.

Poornananda, M. N. and Hosakatte, N. M. (2009). The effect of Gamma and Ethylmethyl sulphonate treatments on agronomical traits of niger (*Guizotia abyssinica* Cass.). *African J. Biotech.*, 8(18): 4459-4464.

Ramesh, B., Prasad, B. K., and Singh, V. P. (2001). Semi dwarf, high yielding and high protein mutants in barley. *Mutation Breeding Newsletter*, 45: 26-27.

Ravikesavan, R., Kalaimagal, T., Rathnaswamy, R. (2001). An extra early mutant of pigeonpea. *Mutation Breeding Newsletter*, 45: 19-20.

Sakin, M. A., Yildirim, A. (2004). Induced mutations for yield and its component in durum wheat (*Triticum durum* Defs.). *Food Agric. Environ.* 2(1): 285-290.

SAR. (2005). Final Safety Analysis Report of Nigeria Research Reactor-1. CERT Technical Report-CERT/NIRR-1/FSAR-01.

Schippers, R. R. (2000). African indigenous vegetables: An overview of the cultivated species. Natural Resources Institute/ACP-EU Technical Centre for Agricultural and Rural Cooperation, Chathan, UK, pp. 122-133.

Shah, A. B. S., Ud-Deen, M. M., Naz, A., Sarker, J. K. and Kabir, G. (2008). Post-irradiation ageing effect on morphological characters of *Crotalaria saltiana*. *Journal of Biological Sciences*, 16: 89-93.

Simmond, N.W. (1976). *Evolution of Crop Plants*. Longman Co. Ltd, London. pp. 265-68.

Singh, S., Richharia, A. K., Joshi, A. K. (1998). An assessment of Gamma ray induced mutations in rice (*Oryza sativa* L.). *Indian J. Genet.*, 58(4): 455-463.

Wani, A. A. and Anis, M. (2001). Gamma rays induced bold seeded high yielding mutant in chickpea. *Mutation Breeding Newsletter*, 45: 20-21