

VIABILITY AND VIGOUR OF COWPEA (*Vigna unguiculata* L. Walp) VARIETIES SEEDS OBTAINED FROM PLANT SOWN IN SEQUENTIAL CROPPING SYSTEM

O. A. ADEDIRAN^{*1}, A. M. MUSTAPHA¹, H. IBRAHIM¹, O. O. OLUFAJO² AND J. A. OLADIRAN¹

¹Department of Crop Production, Federal University of Technology, Minna, Nigeria.

²Department of Agronomy, Ahmadu Bello University, Zaria, Nigeria.

*Corresponding author

E-mail- o.adediran@futminna.edu.ng Department of Crop Production, Federal University of Technology, P.M.B. 65, Minna, Nigeria.

ABSTRACT

The poor quality of seeds used by farmers is a limiting factor contributing to low yields obtained in cowpea. This study aimed to determine the viability and vigour of cowpea seeds produced in sequential cropping system. The seeds obtained from an earlier field experiment on cowpea varieties planted in sequential cropping system were used for this study. The treatments were factorial combination of five varieties of cowpea viz: IT90K-76, IT93K-452-1, IT99K-573-1-1, TVX 3236, Oloyin and two sequence of planting. They were replicated three times. Data were collected on 100 seed weight, germination percentage and seedling vigour index. Results showed that all the varieties produced significantly heavier seeds in the second sequence than the first except IT93K-452-1 and TVX 3236 plants which produced similar seed weight in the two sequence. Cowpea seeds obtained in the second sequence of planting had significantly higher ($P<0.05$) germination percentage and seedling vigour index than seeds obtained in the first sequence except in TVX 3236 variety. Significant variation existed in the viability and seedling vigour of the varieties in storage. Variety IT93K-452-1 stored better and maintained significantly higher ($P<0.05$) germination percentage and seedling vigour than the other varieties assessed.

Key words: Cowpea, Varieties, Germination percentage, Vigour.

INTRODUCTION

Cowpea (*Vigna unguiculata* L. Walp) is an important grain legume in the dry savannah of the tropics covering 12.5 million hectares with annual production of about 3 million tons (FAO, 2005). The grains are utilized in a wide variety of local dishes and have great potential to fortify food. Yield of cowpea in sub-Saharan Africa has been very low with just about 22% of the potential yield obtained on farmers' fields. The poor quality of seeds used by farmers have been

identified as a limiting factor contributing to low yields in cowpea (Odindo, 2007). Availability of quality seeds of improved, high yielding varieties to local farmers is still a bottleneck for increasing the grain yield of cowpea.

Seed is an important input for agricultural production. Improving the availability of high-quality seeds of improved varieties is important to boost agricultural productivity, leading to higher farmers' income, reduced poverty and improved food security (Abdoulaye *et al.*, 2009). Seed quality includes genetic quality, physical quality, physiological quality and seed health (Louwaars, 2007). Physiological quality refers to the seeds' ability to germinate, emerge from the soil and form a vigorous seedling, especially under the stressful conditions prevailing in the fields of most smallholder farmers. High physiological quality is essential to create an optimal and uniform plant density with minimum seeding rates. Physiological quality is influenced by environmental conditions during crop growth, harvesting and storage conditions (Ghassemi-Golezani and Mazloomi-Oskooyi, 2008).

The only way to increase agricultural production is to increase the yield of individual crop per unit area of land, as food requirements have increased over the years while land availability has become less due to increase in population and urban development. Sequential cropping system refers to growing two or more crops in sequence on the same field per year; the succeeding crop is planted after the preceding crop has been harvested. It is an option to increase the productivity of crop from a unit of land. Its adoption may substitute for expanding cropland acreage which may not always be available and sustainable due to increase in population and urbanization. In an attempt to increase the productivity of crops using the sequential cropping system, it is equally important to evaluate the quality of the seeds obtained in the system as majority of smallholder farmers in developing countries depend on seed-producing farmers or on their own seed saved

from their food harvest; the informal seed system. In such a system seed quality remains unknown, or the perception of the quality is merely based on the reputation of the producer (Van Gastel *et al.*, 2002). Farmers are not fully aware of the agronomic value of high-quality seed.

Hence, this study aimed at evaluating the quality of cowpea seeds produced in cowpea sequential cropping system using the germination percentage and seedling vigour index as indices.

MATERIALS AND METHODS

The experiment was carried out in the Department of Crop Production laboratory, Federal University of Technology Minna, Nigeria. The treatments were five varieties of cowpea (IT93K-452-1 (SAMPEA 8), IT99K-573-1-1 (SAMPEA 14), TVX 3236, IT90K-76 and Oloyin) and two sequence of planting. These were factorially combined and arranged in completely randomized design with three replicates. The seeds obtained from cowpea varieties planted in sequential cropping system were used for the experiment. The plants that produced the seeds harvested in the first sequence were planted on May 5th, 2017 and the second sequence was planted immediately after the harvest of the first planting (in different dates in August, 2017 depending on the variety).

The seeds from all the treatment combinations were collected in small plastic containers stored at initial moisture content of 12% for two weeks after each harvest before testing the quality. The containers were then stored in the incubator at 41.56°C and relative humidity of 81.2% for 10 weeks to accelerate the ageing process.

Data were collected on 100 seed weight in which three replicates of hundred seeds were counted and weighed on a Mettler balance. Seed viability was tested using the seed germination test following the procedures of ISTA (2003). Twenty five seeds in three replicates were taken from

each of the lot stored in the incubator every 2 weeks. They were placed in paper towels moistened with 10 ml of distilled water. The paper towel were placed inside a germinator and seedlings evaluated at 1 to 7 days after sowing. Germination percentage (GP) was then expressed

as:
$$\frac{\text{Number of normal seedlings}}{\text{No of total seeds tested}} \times 100$$

At seven days, ten normal seedlings were randomly selected from each replicate and the plumule length was measured with the aid of a meter rule and the average found. Seedling vigour index (SVI) was then calculated as follows:

$$\text{SVI} = \frac{\text{seedling plumule length} \times \text{germination percentage}}{100}$$

Data collected were subjected to analysis of variance (ANOVA) using SAS version 9.1. Means were separated using Duncan multiple range test at P=0.05. Were significant interaction exists throughout the sampling period, interaction means at 0, 6 and 10 weeks are presented.

RESULTS

Plants sown in the second sequence produced significantly heavier seeds (15.37 g) than plant sown in the first sequence (13.89 g). There was significant difference (P<0.01) among the seed weight of the varieties; the order of performance in respect of seed weight was Oloyin > IT99K-573-1-1 > IT90K-76 > IT93K-452-1 > TVX 3236. However, there was no significant difference between the values recorded in IT99K-573-1-1 and IT90K-76 (Table 1). The interaction effect of sequence of planting and variety on seed weight was highly significant (P<0.01). All the varieties produced significantly heavier seeds in the second sequence than the first except IT93K-452-1 and TVX 3236 plants which produced similar seed weight in the two sequence (Figure 1).

Seeds obtained from plants sown in the second sequence had significantly higher germination percentage than those obtained in the first sequence of planting throughout the storage period.

Among the varieties, Oloyin had significantly highest germination percentage (68.11%) before storage, followed by IT99K-573-1-1 similar to IT90K-76 and the least was observed in TVX 3236 (38%). There was variation in the germination response of the varieties throughout the storage period and GP reduced as the storage period increased from 2 to 10 Weeks in storage (WIS). At late storage however (10 WIS), only seeds of IT93K-452-1 remained viable as other varieties had lost their viability at this stage.

Highly significant interactions ($P < 0.001$) existed between the sequence of planting and variety on germination percentage throughout the sampling period. Before storage (0 WIS), there was no significant difference between the germination percentage of seeds obtained in the two sequence of planting in IT93K-452-1 and TVX 3236 varieties. However, in the other varieties, seeds harvested in the second sequence of planting had significantly higher GP than those harvested in the first sequence. At mid storage (6 WIS), all the seeds of the varieties harvested in the second sequence of planting had significantly higher GP than those harvested in the first sequence except IT99K-573-1-1 which had similar GP in both sequence. IT93K-452-1 sown in the second sequence maintained significantly higher GP (100%) than the other varieties whose GP ranged between 57-85% in the second sequence. The least GP in this sequence was obtained in IT99K-573-1-1 (12%). At late storage, there was no significant difference between the GP obtained at 1st and 2nd sequence of planting among all the varieties as most have lost their viability except IT93K-452-1 in which seeds obtained in the second sequence of planting had significantly higher GP (28%) than those obtained in the first sequence (1.33%) (Figure 2).

Seeds obtained in the second sequence had significantly higher seedling vigour index than those obtained in the first sequence throughout the sampling period (Table 2). Among the varieties, TVX 3236 had significantly highest vigour (12.30) before storage followed by IT93K-452-1

(10.15). The least was recorded in IT90K-76 (1.00) similar to Oloyin (2.60). Significant variation existed between the vigour of the varieties in storage, however, IT93K-452-1 maintained significantly higher vigour than the remaining varieties in storage. At late storage (9 WIS), IT99K-573-1-1 and Oloyin had lost their vigour completely and by 10 WIS, all the varieties except IT93K-452-1 had completely lost vigour.


The interaction between sequence of planting and variety on seedling vigour index was significant ($P < 0.05$) throughout the sampling period. Before storage (0 WIS), IT99K-573-1-1 and IT90K-76 planted in the first sequence had similar vigour with those planted in the second sequence. For the remaining varieties however, seeds obtained in the second sequence of planting had significantly higher SVI than those obtained in the first planting. Among the plants sown in the first sequence, TVX 3236 had significantly highest SVI (10.72) followed by IT93K-452-1 (6.16) similar to IT99K-573-1-1 (4.80) and the least was recorded in IT90K-76 (0.90) similar to Oloyin variety (1.16). Among the seeds obtained in the second sequence, IT93K-452-1 had the significantly highest SVI (14.13) similar to TVX 3236 (13.87). The least was recorded in IT90K-76 (1.10).

At 6 WIS, there was no significant difference between the vigour recorded in seeds obtained in the two sequence except in IT93K-452-1 and IT90K-76 in which seeds obtained in the second sequence had significantly higher SVI than the first. IT93K-452-1 maintained significantly higher SVI in 1st and 2nd sequence of planting at mid storage (3.43 and 16.73 respectively). At this storage period, seeds of IT99K-573-1-1 harvested in both sequence had completely lost vigour similar to the seeds of IT90K-76 and Oloyin harvested in the first sequence of planting. At late storage (10 WIS), all the varieties planted in both sequence had lost their viability except IT93k-452-1 obtained in the second sequence.

Table 1: Effect of sequential cropping system on 100 seed weight (g) and germination percentage (%) of cowpea varieties.

Treatments		Germination percentage Weeks in storage					
Sequence of planting (S)	100-seed wt	0	2	4	6	8	10
1 st planting	13.89b	55.55b	29.87b	12.53b	6.93b	7.73b	0.27b
2 nd planting	15.37a	61.47a	80.00a	64.27a	33.33a	26.13a	6.67a
SE \pm	0.12	0.49	2.79	1.46	1.85	0.20	0.73
Varieties (V)							
IT93K-452-1	14.20c	56.80c	76.67a	68.67a	61.33a	50.67a	14.67a
IT99K-573-1-1	16.24b	64.94b	30.67c	8.00d	2.00d	0.67d	0.00b
TVX 3236	9.68d	38.71d	78.00a	44.00b	16.67b	16.67b	0.00b
IT90K-76	16.00b	64.01b	50.67b	42.67b	13.33bc	10.67bc	2.00b
Oloyin	17.03a	68.11a	38.67c	28.67c	7.33c	6.67c	0.00b
SE \pm	\pm 0.20	0.78	3.75	2.31	2.93	2.37	1.16
Interaction							
S x V	**	**	**	**	**	**	**

Means followed by similar alphabets within the same factor and column are not significantly different using Duncan multiple range test at P=0.05



Bars carrying similar alphabets are not significantly different using DMRT at P =0.05




Figure 2: Interaction effect of sequence of planting and variety on germination percentage of cowpea

Bars carrying similar alphabets are not significantly different using DMRT at P =0.05

Table 2: Effect of sequential cropping system on seedling vigour index of cowpea varieties.

Treatments	Weeks in storage					
	0	2	4	6	8	10
Sequence of planting (S)						
1 st planting	4.75b	2.94b	1.57b	1.00b	0.64b	0.00b
2 nd planting	7.86a	6.04a	8.72a	4.68a	2.44a	0.54a
SE \pm	0.44	0.40	0.47	0.24	0.20	0.07
Varieties (V)						
IT93K-452-1	10.15b	8.38a	11.50a	10.10a	5.36a	1.36a
IT99K-573-1-1	5.54c	3.02c	0.81c	0.00d	0.00c	0.00b
TVX 3236	12.30a	5.03b	4.76b	2.15b	1.38b	0.00b
IT90K-76	1.00d	3.27bc	4.74b	1.27c	0.96b	0.00b
Oloyin	2.60d	2.76c	3.89b	0.71cd	0.00c	0.00b
SE \pm	0.78	0.62	0.74	0.39	0.32	0.11
Interaction						
S x V	**	**	*	**	**	**

Means followed by similar alphabets within the same factor and column are not significantly different using Duncan multiple range test at P=0.05




Figure 3: Interaction effect of sequence of planting and variety on seedling vigour index of cowpea

Bars carrying similar alphabets are not significantly different using DMRT at P = 0.05

DISCUSSION

The difference in the germination percentage and vigour of seeds recorded in the two sequence could be attributed to the different environmental conditions prevalent at the period of production of the mother plants. Temperature, photoperiod, water deficits and soil fertility are known for their effects on seed mass, number and quality (Gutterman, 1992; Bewley and Black, 1994; Dornbos, 1995; Wulf, 1995). Temperature was reported to influence seed germination and vigour. High temperatures during seed development was reported to cause early pod ripening and rapid seed maturation and result in small, poor quality seeds in common beans (Siddique and Goodwin, 1980) and annual rye grass (Steadman *et al.* , 2004). In seeds of bean, wheat and barley, lower temperatures during and after seed filling significantly influenced the development of seed quality by extending the post maturation phase, resulting in seeds with higher potential longevity (Ellis and Filho, 1992; Sanhewe and Ellis, 1996).

The decline in the GP and vigour of the seeds with increase in storage period is as a result of ageing. Seed ageing and deterioration affects both seed germination and seed vigor (Mohammadi *et al.*, 2011). Seed deterioration is inevitable but the rate can be reduced by paying due attention to the temperature and relative humidity during storage (Adetumbi *et al.*, 2009).

The significantly lower GP recorded in TVX 3236 compared to the remaining varieties could be attributed to the smaller seed size obtained in this variety. Similar results was reported by Samreen and Shahid (2000) who reported that GP depended on seed size with large seeds having highest GP. Genetic variation is the major cause of variation in size of seed between varieties. The trend of GP observed in this study followed similar trend with the 100 seed weight.

The insignificant difference between the GP (at early storage) of TVX-3236 seeds obtained in the two sequence, and the highest value of seedling vigour obtained in the first sequence in this

variety could indicate that viable and vigorous seeds could be obtained from this variety irrespective of the production period provided the seeds will not be stored for long. This implies that farmers can multiply these seed early in the season (as soon as rain is established) to produce seeds for the major planting in August to September which is around the time local farmers plant cowpea in the southern guinea savanna of Nigeria. However, if the farmer is to store the seeds for long, seeds harvested in the second sequence of planting (from plants sown in August) is preferable. The significant variation that existed in the varieties in respect of viability, vigour and longevity could be attributed to the variation in the genetic make-up of the varieties.

The higher GP obtained in IT93K-452-1 seeds at 2-6 WIS (100%) compared to the 55% obtained before storage (0 WIS) could be an indication that the variety exhibits some dormancy. Seed dormancy is a physiological adaptation to environmental heterogeneity (Bewly *et al.*, 2013). Dormancy provides a strategy for seed to spread germination in time to reduce the risk of plant death and possible species extinction in an unfavourable environment. Previous study on this variety revealed that the rate of imbibition was slower in this variety than the other five varieties tested. The authors attributed the slow imbibition rate to slow permeability of the testa and seed composition (Wada and Abubakar, 2013). Seed testa exerts its role to protect the embryo against adverse environmental condition by controlling germination through dormancy imposition by limiting the detrimental activity of physical and biological agents during seed storage. This could have contributed to the significantly higher viability and vigour recorded in IT93K-452-1 at late storage. This implies that this variety could store longer than the remaining varieties and it is a reflection of the genetic quality of the variety.

CONCLUSION

Variety IT93K-452-1 was superior in quality parameters (germination percentage, seedling vigour index and longevity) than the other varieties assessed. Also cowpea seeds obtained in the second sequence had better quality than those obtained in the first sequence. However viable and vigorous seeds can be obtained in TVX-3236 produced in two sequence within a growing season in the southern guinea savanna of Nigeria.

RECOMMENDATION

Cowpea seeds harvested in the first sequence should be used as grains except for TVX-3236 variety while those harvested in the second sequence can be used as seeds. IT93K-452-1 can be recommended as the best variety with the highest seed quality among those tested.

REFERENCES

- Abdoulaye T., Sanogo D., Langyintuo A., Bamire S. A. & Olanrewaju A. 2009. Assessing the constraints affecting production and development of maize seed in DTMA countries of West Africa. *International Institute of Tropical Agriculture*, Ibadan, Nigeria.
- Adetumbi J. A., Odiyi A. C., Olakojo S. A. & Adebisi M. A. 2009. Effect of storage environments and storage materials on drying and germination of maize (*Zea mays* L.) seed. *Electronic Journal of Environmental, Agricultural and Food Chemistry* 8 (11), 1140-1149
- Bewley J. D. & Black M. 1994. Seed physiology of development and germination. New York and London: Plenum press. 86-90.
- Bewley, J. D., Bradford, K., Hilhorst H and Nonogaki H. 2013. Seeds: Physiology of development, germination and dormancy, 3rd Edn. New York: Springer Verlerg.
- Dornbos D. L. 1995. Production environment and seed quality. In: Basra A. S., ED. *Seed Quality Basic Mechanisms and Agricultural Implications*. . New York, London, Australia. Food Products Press. 116-134.
- Ellis R. H & Filho C. P. 1992. The development of seed quality in spring and winter cultivars of barley and wheat. *Seed Science Research*, 9, 373-409.
- FAO, 2005. Cowpea production database for Nigeria, (1990-2004). *Food and Agricultural Organization*. <http://www.faostat.fao.org/>.
- Ghassemi-Golezani K. & Mazloomi-Oskooyi R. 2008. Effect of water supply on seed quality development in common bean (*Phaseolus vulgaris* var.). *International Journal of Plant Production*, 2, 117-24.

- Gutterman Y, 1992. Maternal effects on seeds during development: Fenner M. *The Ecology of Regeneration in Plant Communities*. Wallingford, UK: CAB International. 115-135.
- ISTA (International seed testing Association) Hampton J. G. & Tekrony B. M. 2003. *Hand book of Vigor test method 3rd Edition*. Zurich, Switzerland: 3, 68-91.
- Louwaars N. 2007. *Seeds of confusion: the impact of policies on seed systems. Wageningen University, PhD Thesis*. 191.-230.
- Mohammadi H, Soltani A, Sadeghipour H. R & Zeinali E. 2011 Effects of seed aging on subsequent seed reserve utilization and seedling growth in soybean. *International Journal of Plant Production* 5 (1), 65-70.
- Odindo A. O. 2007. *Cowpea seed quality in response to production site and water stress, PhD Thesis, University of KwaZulu-Natal, Pietermaritzburg, South Africa*. 75-130.
- Samreen S. and Shahid S. S. 2000. Effect of seed size on germination, emergence, growth and seedling survival of occidentalis, *Pakistan Journal of Biological Sciences*, 3: 292-295.
- Sanhewe A. J & Ellis R. H. 1996. Seed development and maturation in *Phaseolus vulgaris* .II. Post-harvest longevity in air-dry storage. *Journal of Experimental Botany*, 47, 959-965.
- Siddique M. A & Goodwin P. B. 1980. Seed vigour in bean (*Phaseolus vulgaris* L.cv. Apollo) as influenced by temperature and water regime du ring development and maturation. *Journal of Experimental Botany*, 31, 313-323.
- Steadman K. J, Ellery A. J, Chapman R., Moore A. & Turner N. C. 2004. Maturation temperature and rainfall influence seed dormancy characteristics of annual ryegrass (*Lolium rigidum*). *Australian Journal of Agricultural Research*, 55, 1047-1058.
- Van Gastel T. J. G, Gregg B. R & Asiedu E. A. 2002. Seed Quality Control in Developing Countries. *Journal of New Seeds*, 4, 117-30.
- Wada B. Y. and Abubakar B. Y. 2013. Germination studies in some varieties of *Vigna unguiculata* L. Walp (Cowpea) from Northern Nigeria. *Pakistan Journal of Biological Sciences*, 161220-1222
- Wulf R. D. (1995). Environmental maternal effects on seed quality and germination. In: Kigel J, Galili G, eds. New York: Marcel Dekker. *Seed development and germination*, 4, 117-30