



وزارة البيئة والمياه والزراعة  
Ministry of Environment Water & Agriculture  
Kingdom of Saudi Arabia المملكة العربية السعودية



## **ABSTRACT BOOK**

### **MENA Poultry Conference**

### **"Poultry Production under High Environmental Temperature"**

**14-16 March 2022**

**Riyadh International Convention and Exhibition center**

**Riyadh, Saudi Arabia**

**Welcome from Chairman of the  
MENA Poultry Conference  
"Poultry Production under High Environmental Temperature"**

Dear all attendance

On behalf of the Organizing Committee and the WPSA Saudi Branch it gives me a great pleasure to welcome you all to the MENA Poultry Conference "Poultry Production under High Environmental Temperature". This conference will provide a platform for the exchange of new ideas, information on the latest advances and for building up and strengthening professional relationships.

I would like to express my sincere gratitude to the esteemed scientists who will make this conference more meaningful with their presentations and the sponsors who made this conference possible.

I wish you all an enjoyable and productive time at the conference and a wonderful stay in Riyadh.

**Prof. Ibrahim H. Al-Homidan**

President of World's Poultry Science Association- Saudi Branch  
Chairman of The Middle East and North Africa Poultry Conference

## Scientific Committee

1. Prof. Ibrahim H. Al-Homidan (President of WPSA- Saudi Branch)
2. Prof. Moataz M. Fathi (Secretary of WPSA- Saudi Branch)
3. Prof. Tarek A. Ebeid
4. Prof. Mostafa M. Zeitoun
5. Prof. Mohamed F. El-Zarei
6. Dr. Mohamed A. Shehab-El-Deen
7. Dr. Osama Abou-Emera
8. Mr. Saleh Alwaily
9. Mr. Abdulaziz Al-Dokhail
10. Mr. Abulrahman Almoshwah
11. Mr. Mohamed Al-Muzaini
12. Mr. Malik Alshobramy

## Table of Contents

No	Title	Page
S1.1	Evaluation of production performance and humoral immune response of two Saudi chicken lines and their crosses with foreign breeds under Kingdom of Saudi Arabia environment <i>M. Shawky, Y. Alyousef, A. O. Abbas, and H. Najib</i>	
S1.2	Virulence and antimicrobial resistance genes of <i>Campylobacter jejuni</i> in broiler <i>M. M. Sobhy and Khadijah A. Y. Altammar and Ghaneem, H. E.</i>	
S1.3	Repeatability estimates for feed efficiency of laying hens from local Egyptian strains and there hybrids <i>M. E. Soltan and Eman A. Hussein</i>	
S1.4	Genetic polymorphism of the prolactin promoter region among Red Jungle fowl, Egyptian, and other native fowl populations <i>I. Elkhayat, K. Kawabe, W. Rerkamnuaychoke, T. Shimogiri, K. Saleh, H. Younis, R. Nofal, S. Masuda, S. Taura and S. Okamoto</i>	
S1.5	Crossbreeding effects for heat tolerance traits of Sinai, Gimmizah and Sliver Montazah strains <i>M. E. Soltan, S. omer , Amal S. Zharan and S. A. Farrag</i>	
S1.6	Immunocompetence profile of Saudi native chickens compared to exotic breeds under high environmental temperature <i>A. M. Al-Moshawah, M. M. Fathi, I. H. Al-Homidan, T. A. Ebeid, and O. K. Abou-Emera</i>	
S2.1	Modulating laying hens productivity and immune performance in response to oxidative stress induced by <i>E. coli</i> challenge using dietary propolis supplementation <i>A. O. Abbas, A. A. Alaqil, H. S. El-Beltagi, H. K. Abd El-Aaty, and N. N. Kamel</i>	
S2.2	Ochratoxin-A induced pathological alterations in IgA and IgY of broilers <i>S. A. Khan, and E. N. Itano</i>	
S2.3	Effects of maternal dietary selenium sources on growth performance, meat quality, antioxidant status and immune responses of broiler progeny <i>S. Ashraf, S. A. Bhatti, T. Mahmood, M. Yousaf, N. Us Sahar, and M. Shoaib</i>	
S2.4	<i>Citrullus colocynthis</i> Seeds: A Potential Natural Immune Modulator Source for Broiler Reared under Chronic Heat Stress <i>M. I. Alzahrah, F. Althobiati, A. O. Abbas, G. M. Mehaisen, and N. N. Kamel</i>	
S2.5	Protective effect of <i>Curcuma Longa</i> against Newcastle Disease (ND) and Marek's Disease (MD) viruses <i>H. M Madbouly, M.A.Saif, Fadwa Amin, SM.Tamam, AS husein, and Samar S. Eweis</i>	
S2.6	Phytogenic feed additives made of <i>Artemisia annua</i> and thyme in broiler diet to enhance performance, carcass characteristics, and intestinal parameters <i>A. Al-Zahrani and H. Najib</i>	
S3.1	Effect of probiotics and organic acids supplementation on growth performance, carcass characteristics, microbiota, antioxidative status, and immune response in broiler chickens <i>R. E. Al-Jamaan, T. A. Ebeid, I. H. Al-Homidan, O. K. Abou-Emera, and M. M. Fathi</i>	
S3.2	Effect of crate type on egg quality and shelf life of eggs from Isa-brown layers reared on the deep litter system	

	<i>A, Usman, K. E. Akande, F. G. Adegboyega, and H.A. Dikko</i>	
S3.3	Potential of a host specific multispecies synbiotic on broiler performance and gut health <i>S. Basharat, I. Mueller, M. Mohnl</i>	
S3.4	Effect of dietary fibre and fat levels on growth, carcass, digestive enzymes, intestinal microbiota and economics in growing Egyptian geese <i>M.E. Abd El-Hack, M.S. El-Kholy, E.A. Ashour, S. Mahgoub, A.M.T. Mohamed and M. Alagawany</i>	
S3.5	Influence of probiotic inclusion and stocking density on body weight and bacterial count profile of broiler chickens <i>B. Al-Habob, M. M. Fathi, I. H. Al-Homidan, T. A. Ebeid, and O. K. Abou-Emera</i>	
S3.6	Inclusion of Citrullus colocynthis seed extract into diets induced a hypolipidemic effect and improved layer performance <i>M. I. Alzahrah, A. A. Alaqil, A. Abbas, F. S. Nassar, G. M. Mehaisen, G. F. Gouda, H. K. Abd El-Aty, and E. S. Moustafa</i>	
S4.1	Effects of dietary probiotic (Bacillus subtilis) supplementation on productive performance, immune response and egg quality characteristics in laying hens under high ambient temperature <i>A. A, Al-Dokhail, M. M. Fathi, I. H. Al-Homidan, T. A. Ebeid, and O. K. Abou-Emera</i>	
S4.2	Effects of humate substances on blood haematology and iron profile of broiler chicks reared under Egyptian summer conditions <i>S. S. Elnesr, A.M. Abdelsalam and H. A. M. Elwan</i>	
S4.3	The effect of heat stress on broiler and layer performance according the house environment <i>H. A. Fahmy and V. A.ELsayaad</i>	
S4.4	Alleviation of chronic heat stress in broilers by dietary supplementation of novel feed additive combinations <i>S. I. Al-Sultan, S. M. Abdel-Rahman, S. S. Abd-Allah and A. M. Edris</i>	
S4.5	Effect of using yeast in water and waste date palm meal supplementation on productive performance and physiological status of Japanese quail during summer season <i>W. Fouad</i>	
S4.6	Dietary supplementation of ginger (Zingiberofficinale) powder to alleviate heat stress in broilers during summer months <i>M.A.A. Abdelmageed and I. H. Al-Homidan</i>	
S5.1	Use of Exogenous Feed Pigments for Egg Yolk Pigmentation – A New Innovation Tool <i>S. Ashraf and I. Hashmi</i>	
S5.2	Productive performance, ovarian follicular development, lipid peroxidation, antioxidative status, and egg quality in laying hens fed diets supplemented with salvia officinalis and origanummajorana powder levels <i>A. Saleh, S. Hamed, A. M. Hassan 2, K. Amber, W. Awad 3, M. H. Alzawqari, M. Shukry, and A. Y. Desouky</i>	
S5.3	Genetically modified feeds and their effects on poultry carcass and meat quality <i>M. Bakhsh and M. K.Yar</i>	
S5.4	The role of dietary yeast extracts in improving the performance and health of quail breeders reared under high stocking density	

### S3.2

#### **Effect of crate type on egg quality and shelf life of eggs from ISA-brown layers reared on the deep litter system**

*Usman, A.<sup>1\*</sup>, K. E. Akande<sup>1</sup>, F. G. Adegboyega<sup>1</sup>, H.A. Dikko<sup>1</sup>*

*<sup>1</sup> Department of Animal Production, School of Agriculture and Agricultural Technology, Federal University of Technology, Minna, Niger State, Nigeria*

\*Presenter: usman.abdul@futminna.edu.ng

#### **Abstract**

**Implications:** Measurement of egg quality is important for direct consumption or setting in the breeder industry. Poor quality eggs from storage are a great source of concern in microbial contamination and hatchability of the eggs. Eggs are collected in crates and stored before consumption or setting in the hatcheries. Information on comparative properties of eggs stored in different crate types are few, which this study has sought to address.

**Introduction:** Poultry is for most Nigerians, can be depended upon to ameliorate the deficit in protein supply of the diet in the growing population as an egg is reported to provide up to 6g of protein while the egg-white protein has a biological value of 100; the highest biological value of any single protein ever reported. An egg can easily get spoiled and quickly lose its quality if not properly handled. This is because egg shells permits passage of air and fluid, hence moisture and carbon dioxide could escape through the shell under poor storage conditions. There has been growing research interest in studying egg quality during storage which forms the basis for the study in seeking the effect of paper crates, glass crates and plastic crates on egg quality over different storage periods.

**Materials and methods:** A total of 180 freshly laid eggs were used for the study and the eggs were randomly and equally allotted to 3 treatments of the crate types (paper, glass and plastic) in a Completely Randomized Design (CRD) with each treatment having 60 eggs. Six eggs were selected at random from each egg crate type (paper, glass and plastic). In each crate type, 6 eggs were picked on a weekly interval of up to 8 weeks (144 eggs) for the determination of internal and external parameters such as the Haugh Unit (HU) and Yolk Index (YI) were measured.

**Results:** The results obtained showed that at week three of egg storage, egg quality indices were significantly ( $P < 0.05$ ) affected by storage duration and crate types. Based on grading of stored eggs using HU as a standard for egg acceptability to the consumer, eggs stored using paper crate had grade A (78%), while eggs stored in a plastic crate (68.7%) and eggs in a glass crate (63.2%) which were both grade B. Whereas the YI was not affected significantly ( $P > 0.05$ ) by storage duration in the different crate types.

**Conclusion:** Based on the observations of results from this study, it is therefore recommended that eggs should be stored in paper crates. This is because both the internal and external quality parameters were better preserved by using the paper crate and this also resulted in the maintenance of grade A egg quality even at three weeks of storage.