



Contents lists available at ScienceDirect

Journal of Acute Disease

journal homepage: www.jadweb.org



Document heading doi: 10.1016/S2221-6189(13)60072-4

Inhibitory activity of xanthine oxidase by fractions *Crateva adansonii*

Abdullahi A^{1*}, Hamzah RU¹, Jigam AA¹, Yahya A¹, Kabiru AY¹, Muhammad H¹, Sakpe S¹, Adefolalu FS¹, Isah MC², Kolo MZ¹

¹Department of Biochemistry, Federal University of Technology, Minna, Niger state, Nigeria

²Department Biological Sciences, IBB University, Lapai, Niger State, Nigeria

ARTICLE INFO

Article history:

Received 10 January 2012

Received in revised form 15 March 2012

Accepted 15 May 2012

Available online 20 November 2012

Keywords:

Crateva adansonii

Inhibitory

Xanthine oxidase

ABSTRACT

Objective: To study the inhibitory effect of various extracts from *Crateva adansonii* (*C. adansonii*) used traditionally against several inflammatory diseases such as rheumatism, arthritis, and gout, was investigated on purified bovine milk xanthine oxidase (XO) activity. **Methods:** Xanthine oxidase inhibitory activity was assayed spectrophotometrically and the degree of enzyme inhibition was determined by measuring the increase in absorbance at 295 nm associated with uric acid formation. Enzyme kinetics was carried out using Lineweaver–Burk plots using xanthine as the substrate. **Results:** Among the fractions tested, the chloroform fraction exhibited highest potency (IC₅₀ 20.2±1.6 μg/mL) followed by the petroleum ether (IC₅₀ 30.1±2.2 μg/mL), ethyl acetate (IC₅₀ 43.9±1.4 μg/mL) and residual (IC₅₀ 98.0±3.3 μg/mL) fractions. The IC₅₀ value of allopurinol used, as the standard was 5.7±0.3 μg/mL. **Conclusions:** Enzyme inhibition mechanism indicated that the mode of inhibition was of a mixed type. Our findings suggest that the therapeutic use of these plants may be due to the observed Xanthine oxidase inhibition, thereby supporting their use in traditional folk medicine against inflammatory-related diseases, in particular, gout.

1. Introduction

Gout and hyperuricemia are the common metabolic disorders in human, associated with an elevated uric acid level in the blood^[1], leading to the deposition of urate crystals in the joints and kidneys, leading to gouty arthritis and uric acid nephrolithiasis. Xanthine oxidase (XO) and xanthine dehydrogenase (XDH) are the two interconvertible forms of same gene product known as xanthine oxidoreductase (XOR). It catalyses the oxidation of hypoxanthine to xanthine and then to uric acid, the final reactions in the metabolism of purine bases^[2,3]. The treatment for gout is either increasing the excretion of uric acid or reducing the uric acid production. Xanthine oxidase inhibitors (XOIs) are much useful, since they possess lesser side effects compared to uricosuric and anti-inflammatory agents. Allopurinol is the only clinically used XOI, which also suffers from many side effects such as hypersensitivity syndrome, Stevens Johnson syndrome and renal toxicity^[4].

Thus, there is a need to develop compounds with XOI activity which are devoid of the undesirable side effects of allopurinol. A potential source of such compounds can be obtained from medicinal plants^[5,6]. Flavonoids and polyphenolic crude extracts have been reported to possess xanthine oxidase inhibitory activity^[1,7,8]. *Crateva adansonii* (*C. adansonii*) belonging to the family *Capparaceae* and phylum *Magnoliophyta* is small tree of forest and savanna woodland, often on river-banks, widely distributed in Nigeria, and Across Africa^[9,10]. The leaves are applied externally to relieve pain in joints; the fresh juice of leaves is used for the relief of ear ache, eye infection and anodyne in toothache. Powder of bark is used in rheumatism, itch, epilepsy, and asthma^[11]. Many medicinal plants have been used for the prevention and treatment of gout and related inflammatory disorders, but they lack sufficient scientific evidence. Hence, the objective of the present study is to determine the *in vitro* xanthine oxidase inhibitory activity of the fractions of the leaves *C. adansonii* and its enzyme inhibition mechanism.

*Corresponding author: Abdullahi A, Department of Biochemistry, Federal University of Technology, Minna, Niger state, Nigeria.
E-mail: abkad2008@gmail.com

2. Materials and methods

2.1. Plant material

The plant material consists of dried powdered leaves of *C. adansonii* belonging to the family *Capparaceae* and phylum *Magnoliophyta*. The leaves were collected in and around Minna, Niger State of Nigeria and was authenticated by the staff of Biological Science Department of Federal University of Technology, Minna.

2.2. Preparation of the extract and fractionation

The air-dried powdered leaves of *C. adansonii* (500 g) were extracted with methanol–water (7:3) mixture using a mechanical shaker for 4 h. The resultant extract was concentrated under reduced pressure to yield a residue. The hydromethanolic extract was then extracted successively with equal volumes of petroleum–ether, chloroform and ethyl acetate. Each fraction was then concentrated under reduced pressure to obtain petroleum–ether fraction (PEF), chloroform fraction (CF), ethyl acetate fraction (EAF) and residual fraction (RF).

2.3. Chemicals

XO (source: Bovine Milk) XO was purified from bovine milk using the method of Atmani et al^[12] and XO was kept frozen and thawed just before use. The buffer used in our study was 0.1 M monopotassium phosphate–disodium phosphate ($\text{KH}_2\text{PO}_4/\text{Na}_2\text{HPO}_4$), and xanthines were obtained from Department of Biochemistry Laboratory Federal University of Technology, Minna. Allopurinol was obtained from Zagbayi pharmaceutical Minna. All other drugs and chemicals used in the study were obtained commercially and were of analytical grade.

2.4. Phytochemical screening

Preliminary phytochemical screening of the plant extract was performed for the presence of alkaloids, phenolics, tannins, flavonoids, saponins, carotenoids, carbohydrates and glycosides^[13].

2.5. In vitro xanthine oxidase inhibitory activity

All the fractions of *C. adansonii* were assayed for their *in vitro* xanthine oxidase inhibitory activity. The XO activity was assayed spectrophotometrically using xanthine as the substrate^[14]. The assay mixture consisted of 1 mL of the fraction (5–100 $\mu\text{g/mL}$), 2.9 mL of phosphate buffer (pH 7.5) and 0.1 mL of xanthine oxidase enzyme solution (0.1 units/mL in phosphate buffer, pH 7.5), which was prepared immediately before use.

After preincubation at 25 °C for 15 min, the reaction was

initiated by the addition of 2 mL of the substrate solution (150 mM xanthine in the same buffer). The assay mixture was incubated at 25 °C for 30 min. The reaction was then stopped by the addition of 1 mL of 1 N hydrochloric acid and the absorbance was measured at 290 nm using a UV spectrophotometer^[15,16]. Different concentrations of the fractions (5–100 $\mu\text{g/mL}$) were dissolved in dimethyl sulfoxide (DMSO) and the final concentration of DMSO was 5%, which did not affect the enzyme assay.

Proper controls with DMSO were carried out. Allopurinol (5–100 $\mu\text{g/mL}$), a known inhibitor of XO, was used as the positive control. One unit of XO is defined as the amount of enzyme required to produce 1mmol of uric acid/min at 25 °C. XO activity was expressed as the percentage inhibition of XO in the above assay system calculated as

Inhibition (%) = $(1 - [B/A]) \times 100$, where A represents the activity of the enzyme without plant extract and B is the activity of XO in the presence of plant extract.

The enzymatic inhibitory mode of action of different extracts from both plants that exhibited the highest inhibition of XO activity was determined by the Lineweaver–Burk plot using varying concentrations of xanthine.

2.6. Enzyme inhibition mechanism

To determine the mode of inhibition by the most active chloroform fraction, Lineweaver–Burk plot analysis was performed. The assay was carried out in the absence and presence of the chloroform fraction with varying concentrations of xanthine as the substrate, using the XO assay methodology. The mode of inhibition was compared with that of the positive control allopurinol. The Lineweaver–Burk transformed values were plotted to determine the mode of inhibition^[17,18].

3. Results

Preliminary phytochemical screening of the plant extract revealed the presence of phenolics, alkaloids, flavonoids and saponins (Table 1). All the fractions of *C. adansonii* elicited a dose dependent inhibition of xanthine oxidase enzyme activity. Inhibition of xanthine oxidase resulted in a decreased production of uric acid, which was measured spectrophotometrically.

At a concentration of 100 $\mu\text{g/mL}$, highest activity was observed in the chloroform fraction followed by the petroleum–ether, ethyl acetate and residual fractions. The concentration of the fractions at which 50% of the xanthine oxidase inhibitory activity (IC_{50}) was calculated.

These results were compared with the standard drug allopurinol, which showed (92.25±0.05)% inhibition at 100 $\mu\text{g/mL}$ concentration with IC_{50} value (5.7±0.3) $\mu\text{g/mL}$ (Table 2).

Table 1Phytochemical screening of *C. adansonii*.

Ingredient	Petroleum–ether	Chloroform	Ethyl acetate
Alkaloids	+	+	+
Phenolics	+	+	+
Tannins	–	–	–
Flavonoids	+	+	+
Saponins	+	+	+
Carotenoids	–	–	–
Carbohydrates	–	–	–
Glycosides	–	–	–
Balsams	–	–	–

Key: (+) present, (–) – absent.

Table 2Percentage yield and xanthine oxidase inhibitory activity of the fractions of *C. adansonii*.

Fraction	% Yield fraction (w/w)	Percentage xanthine oxidase inhibition					IC ₅₀ (μg/mL)
		5 μg/mL	10 μg/mL	25 μg/mL	50 μg/mL	100 μg/mL	
Petroleum–ether	1.8	28.35 ± 0.90	39.53 ± 1.20	48.32 ± 0.90	66.73 ± 1.57	78.59 ± 1.60	30.10 ± 2.20
Chloroform	1.7	19.55 ± 1.30	36.00 ± 0.80	55.00 ± 1.00	76.70 ± 1.10	84.75 ± 0.50	20.20 ± 1.60
Ethyl acetate	0.9	11.85 ± 0.30	23.75 ± 0.70	39.50 ± 1.00	56.35 ± 1.20	64.55 ± 1.50	43.90 ± 1.40
Residual	5.1	5.90 ± 1.10	28.80 ± 1.20	35.70 ± 0.50	41.50 ± 1.20	50.05 ± 1.80	98.00 ± 3.30
Allopurinol		44.91 ± 1.80	66.91 ± 0.80	75.00 ± 1.40	85.46 ± 0.80	92.25 ± 0.50	5.70 ± 0.30

The inhibition mechanism of the chloroform fraction of *C. adansonii*, which is a potent inhibitor of xanthine oxidase, was studied by kinetic analysis using double–reciprocal plotting. Lineweaver–Burk plots of reactions in the presence and absence of the chloroform fraction in a xanthine oxidase reaction mixture is shown in Figure 1.

The mode of inhibition was investigated and compared with the standard drug, allopurinol. The data indicates that the mode of xanthine oxidase inhibition for the chloroform fraction is of mixed type (between uncompetitive and non–competitive type of inhibition).

Therefore, the fraction inhibited XO by binding either with the free enzyme or the enzyme–substrate complex.




Figure 1. Lineweaver–Burk plot of inhibition of xanthine oxidase by chloroform fraction of *C. adansonii* (●) and allopurinol (▲). The symbol (◆) represents negative control.

4. Discussion

Recent findings show that the occurrence of gout is increasing worldwide, possibly due to the changes in dietary habits like intake of food rich in nucleic acids, such as meat and seafoods. Hypouricemic agents are commonly employed for the treatment of chronic gouty arthritis, which includes xanthine oxidase inhibitors and uricosuric agents[19]. In general, allopurinol is the drug of choice; however it has serious side effects. Thus, new alternatives with increased therapeutic activity and lesser side effects are desired. We thus began our program to look for xanthine oxidase inhibitors of phytochemical origin from the various fractions of the hydromethanolic extract of the leaves of *C. adansonii*. The leaves of *C. adansonii* are being frequently used in Nigeria traditional medicinal system for the treatment joint pain and related inflammatory disorders.

All the fractions of *C. adansonii* inhibited xanthine oxidase in a concentration–dependent manner. The *in vitro* inhibition of xanthine oxidase by the chloroform fraction is moderate when compared to allopurinol. However, at higher doses of the fraction, xanthine oxidase would be significantly inhibited. Flavonoids are a group of polyphenolic compounds, which have been reported to possess xanthine oxidase inhibitory activity[20].

Hence, the presence of phenolic and flavonoid content in the extract would have contributed towards xanthine oxidase inhibition.

In conclusion, the study suggests that *C. adansonii*

leaves possess xanthine oxidase inhibitory activity that might be helpful in preventing or slowing the progress of gout. Further *in vivo* experiments and the isolation and identification of active compounds present in the leaves should be carried out to identify a potential chemical entity for clinical use in the prevention and treatment of gout and related inflammatory disorders.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgement

We acknowledge Mal. Shuaibu. M and all other supporting staff of the department of Biochemistry.

References

- [1] Meriem B, Karima A, Dina A, Djebbar A, Mustapha B. Kinetic Study on the Inhibition of Xanthine Oxidase by Extracts from Two Selected Algerian Plants Traditionally Used for the Treatment of Inflammatory Diseases. *J Med Food* 2010; **13**(4): 896–904.
- [2] Cos P, Ying L, Calomme M, Hu JP, Cimanga K, Poel BV, et al. Structure–activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and super oxide scavengers. *J Nat Prod* 1998; **61**: 71–6.
- [3] Borges F, Feranandes E, Roleira F. Progress towards the discovery of xanthine oxidase inhibitors. *Curr Med Chem* 2002; **9**: 195–217.
- [4] Burke A, Smyth E, FitzGerald GA. Analgesic–antipyretic agents: pharmacotherapy of gout. In: Brunton LL, Lazo JS, Parker KL. (Eds.). *Goodman & Gilman's The Pharmacological Basis of Therapeutics*. New York: McGraw–Hill Medical Publishing Division; 2006, p. 706–10.
- [5] Kong LD, Cai Y, Huang WW, Cheng CHK, Tan RX. Inhibition of xanthine oxidase by some Chinese medicinal plants used to treat gout. *J Ethnopharmacol* 2000; **73**: 199–207.
- [6] Zhou CX, Kong LD, Ye WC, Cheng CHK, Tan RX. Inhibition of xanthine and monoamine oxidases by stillbenoids from *Veratrum taliense*. *Planta Medica* 2001; **67**: 158–61
- [7] Lio M, Moriyama A, Matsumoto Y, Takaki N, Fukumoto M. Inhibition of xanthine oxidase by flavonoids. *J Agric Biol Chem* 1985; **49**: 2173–6.
- [8] Chang WS, Lee YJ, Lu FJ, Chaing HC. Inhibitory effects of flavonoids on xanthine oxidase. *Anticancer Res* 1993; **13**: 2165–70.
- [9] Abdullahi M, Muhammed G, Abdulkadir U. Medicinal and economic plants of Nupeland. 2003, p. 61.
- [10] Abdullahi Mann, Kolo Ibrahim, Adebayo O Oyewale, Joseph O Amupitan, Joseph I Okogun. Antimycobacterial activity of some medicinal plants in niger state, nigeria. *Afr J Infect Dis* 2009; **3**(2): 44–8.
- [11] Sivarajan VV, Balachandran I. *Ayurvedic Drugs and their Plant Sources*. Delhi: Oxford and IBH Publishing Company Pvt. Ltd; 1994.
- [12] Atmani D, Benboubetra M, Harrison R. Goats' milk xanthine oxidoreductase is grossly deficient in molybdenum. *J Dairy Res* 2004; **71**: 7–13.
- [13] Harborne JB. *Phytochemical Methods. A Guide to Modern techniques of Plant Analysis*. London: Chapman and Hall Ltd; 1998.
- [14] Owen PL, Johns T. Xanthine oxidase inhibitory activity of northeastern North American plant remedies used for gout. *J Ethnopharmacol* 1999; **64**: 149–160.
- [15] Nguyen MTT, Awale S, Tezuka Y, Tran LQ, Watanabe H, Kadota S. Xanthine oxidase inhibitory activity of Vietnamese medicinal plants. *Biol Pharm Bull* 2004; **27**: 1414–21.
- [16] Umamaheswari M, Asokkumar K, Somasundaram A, Sivashanmugam T, Subhadradevi V, Ravi TK. Xanthine oxidase inhibitory activity of some Indian medical plants. *J Ethnopharmacol* 2007; **109**: 247–551.
- [17] Yu Z, Fong WP, Cheng CHK. The dual action of morin (3, 5, 7, 4, pentahydroxyflavone) as a hypouricemic agent, uricosuric effect and xanthine oxidase inhibitory activity. *J Pharmacol Exp Ther* 2005; **316**: 169–75.
- [18] Nguyen MTT, Awale S, Tezuka Y, Tran LQ, Watanabe H, Kadota S. Xanthine oxidase inhibitors from the heart wood of Vietnamese *Caesalpinia sappan*. *Chem Pharm Bull* 2005; **53**: 984–8.
- [19] Pacher P, Nivorozhkin A, Szabo C. Therapeutic effect of xanthine oxidase inhibitors: renaissance half a century after the discovery of allopurinol. *Pharmacol Rev* 2006; **58**: 87–114.
- [20] Costantino L, Albasini A, Rastelli G, Benvenuti S. Activity of polyphenolic crude extracts as scavengers of superoxide radicals and inhibitors of xanthine oxidase. *Planta Medica* 1992; **58**: 342–4.