ation.

entive

tigate rmine

C13-

ck or

UBT.

I, 4th,

ori to

e was

and

atory

n the

pri to

astric

city.

Amino acid profile, protein digestibility, antioxidant and functional properties of protein concentrate of local varieties (Kwandala, Yardass, Jeep and Jamila) of rice bran from Nigeria

C. E. Chinma¹, J. C. Anuonye¹, S. O. Azeez¹, O. B. Ocheme¹, C. M. Yakubu¹, S. James¹, E. U. Ohuoba¹ and I. A. Baba¹

Department of Food Science and Technology, Federal university of Technology Minna, Nigeria

There is growing interest in the use of rice bran protein in food formulation due to its hypoallergenic protein, nutritional value and health promoting potentials. For the first time, amino acid profile, protein antioxidant and functional properties of protein concentrate of some local varieties of rice bran Nigeria were studied. It was found that the protein content of Kwandala, Yardass, Jeep and Jamila 69.24, 69.97, 68.73 and 71.62%, respectively while total essential amino acid composition were 52.71, 53.03, 51.86 and 55.75 g/100 g protein, respectively. Protein digestibility of protein concentrate from Is a separate of the separate immunition of proteins from Kwandala, Yardass, Jeep and Jamila were 48.15, 48.90, 47.56 and 53.29%, while ferric reducing antioxidant power were 0.52, 0.55, 0.47 and 0.67mmol TE per gram, espectively. Jamila rice bran protein had higher values of onset (92.57°C), denaturation temperature and enthalpy (0.72J/g) than Jeep (91.46°C, 101.76°C and 0.68J/g, respectively), Kwandala 200.54°C and 0.57 J/g, respectively) and *Yardass* (88.94°C, 99.45°C and 0.51J/g, respectively). digestibility of protein concentrate from Kwandala, Yardas, Jeep and Jamila were 90.70, 91.39, 91.63% respectively. Oil absorption capacity of Kwandala, Yardass, Jeep and Jamila were 3.61, 3.40 and 4.23g oil/g sample respectively, while water absorption capacity were 4.19, 4. 32, 3.55 and water/g sample, respectively. The protein concentrates had low bulk density (0.37-0.43g/ml). Protein expectate from Jamila had the highest foam capacity (37.25%), followed by Yardass (34.20%), Kwandala [32] [4%] and Jeep (28.90%). The protein concentrates showed low emulsifying and gelling capacities. Rice protein concentrate from these local rice varieties from Nigeria could serve as functional ingredients in food formulations and for enriching low protein foods.