
Mathematics, Game Theory and Algebra Compendium





MATHEMATICS RESEARCH DEVELOPMENTS

MATHEMATICS, GAME THEORY AND ALGEBRA COMPENDIUM

VOLUME 3

No part of this digital document may be reproduced, stored in a retrieval system or transmitted in any form or by any means. The publisher has taken reasonable care in the preparation of this digital document, but makes no expressed or implied warranty of any kind and assumes no responsibility for any errors or omissions. No liability is assumed for incidental or consequential damages in connection with or arising out of information contained herein. This digital document is sold with the clear understanding that the publisher is not engaged in rendering legal, medical or any other professional services.

MATHEMATICS RESEARCH DEVELOPMENTS

Additional books in this series can be found on Nova's website under the Series tab.

Additional e-books in this series can be found on Nova's website under the e-book tab.

MATHEMATICS RESEARCH DEVELOPMENTS

MATHEMATICS, GAME THEORY AND ALGEBRA COMPENDIUM

VOLUME 3

JACOB H. MATHIAS EDITOR



Copyright © 2013 by Nova Science Publishers, Inc.

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means: electronic, electrostatic, magnetic, tape, mechanical photocopying, recording or otherwise without the written permission of the Publisher.

For permission to use material from this book please contact us: Telephone 631-231-7269; Fax 631-231-8175 Web Site: http://www.novapublishers.com

NOTICE TO THE READER

The Publisher has taken reasonable care in the preparation of this book, but makes no expressed or implied warranty of any kind and assumes no responsibility for any errors or omissions. No liability is assumed for incidental or consequential damages in connection with or arising out of information contained in this book. The Publisher shall not be liable for any special, consequential, or exemplary damages resulting, in whole or in part, from the readers' use of, or reliance upon, this material. Any parts of this book based on government reports are so indicated and copyright is claimed for those parts to the extent applicable to compilations of such works.

Independent verification should be sought for any data, advice or recommendations contained in this book. In addition, no responsibility is assumed by the publisher for any injury and/or damage to persons or property arising from any methods, products, instructions, ideas or otherwise contained in this publication.

This publication is designed to provide accurate and authoritative information with regard to the subject matter covered herein. It is sold with the clear understanding that the Publisher is not engaged in rendering legal or any other professional services. If legal or any other expert assistance is required, the services of a competent person should be sought. FROM A DECLARATION OF PARTICIPANTS JOINTLY ADOPTED BY A COMMITTEE OF THE AMERICAN BAR ASSOCIATION AND A COMMITTEE OF PUBLISHERS.

Additional color graphics may be available in the e-book version of this book.

LIBRARY OF CONGRESS CATALOGING-IN-PUBLICATION DATA

ISBN: ; 9: /3/84479/972/9 (eBook)

ISSN: 2325-9329

Published by Nova Science Publishers, Inc. † New York

CONTENTS

Preface		vii
Chapter 1	Mathematical Formulation of NMR Experimental Parameters for Diffusion Magnetic Resonance Imaging – Part I (Cylindrical Geometry) <i>M. Dada, O.P. Faromika, O.B. Awojoyogbe, M.A. Aweda</i> <i>and I.A. Fuwape</i>	1
Chapter 2	Mathematical Formulation of NMR Experimental Parameters for Diffusion Magnetic Resonance Imaging – Part II (Cylindrical Geometry) <i>O.B. Awojoyogbe, M. Dada, O.P. Faromika, M.A. Aweda</i> <i>and I.A. Fuwape</i>	21
Chapter 3	Optimization of Kalman Filtering Performance in Received Signal Strength Based Mobile Positioning J.G. Markoulidakis	41
Chapter 4	On Ore Extensions Over near Pseudo Valuation Rings V.K. Bhat and Neetu Kumari	67
Chapter 5	Subset Selection of Treatments & - Better than the Best Control <i>Amar Nath Gill, Anju Goyal and Parminder Singh</i>	77
Chapter 6	Rigorous Kinetic Analysis of the Racket Flick-Motion in Tennis for Generating Topspin and Backspin Katica (Stevanović) Hedrih and Tijana Ivancevic	95
Chapter 7	Linear versus Nonlinear Human Operator Modeling Tijana T. Ivancevic, Bojan N. Jovanovic, Sasha A. Jovanovic, Leon Lukman, Alexandar Lukman and Milka Djukic	121
Chapter 8	Derivations of a Linear Lie Algebra Over a Commutative Ring Xinxing Lai, Shikun Ou and Shuzhen Luo	151

Chapter 9	Investigating the Development of Arithmetic and Algebra in Vedic India: Tribute to Swami Dayananda Saraswati <i>Gurudeo Anand Tularam</i>	163
Chapter 10	Regular Fuzzy Biclosure Spaces U. D. Tapi and R. Navalakhe	185
Chapter 11	On Matrix Near Ring Necat Görentaş and Tevfik Bilgin	193
Chapter 12	Remarks on Contra-αĝ-Continuous Function S. Rose Mary and M. Lellis Thivagar	199
Chapter 13	Decompositions of $(1,2)^*\alpha$ -Continuous Function in Bitopological Spaces S. Rose Mary and M. Lellis Thivagar	207
Chapter 14	Exact Taylor Series with Constant Coefficients Moawia Alghalith	217
Chapter 15	Hedging and Production Decisions under Uncertainty: A Survey <i>Moawia Alghalith</i>	221
Chapter 16	On δ_M - Lifting Modules Ali Omer Alattass	231
Chapter 17	An Application of Ghoussoub-Preiss's Generalized MPL to Periodic Solutions of Hamiltonian Systems with a Fixed Period <i>Li Bingyu and Zhang Shiqing</i>	241
Chapter 18	Partial Actions of Inductive Groupoids on Rings Dirceu Bagio	247
Index		263

PREFACE

This book is devoted to new advances in all branches of mathematics, game theory and applications, and pure and applied algebra and geometry including mathematical formulation of NMR experimental parameters for diffusion magnetic resonance imaging; optimization of Kalman Filtering performance in received signal strength based mobile positioning; ORE extensions over near pseudo valuation rings; subset selection of treatments; rigorous kinetic analysis of the racket flick-motion in tennis for generating topspin and backspin and linear versus nonlinear human operator modeling.

Magnetic resonance sequences have been developed to label the position of a spin, albeit indirectly, through the Larmore frequency. This provides the basis for measuring diffusion. However, the interpretation of the data is complicated by the effects of restricted geometries and the mathematical modeling required to account for this becomes nontrivial especially in cylindrical and spherical geometries. Generally one have to resort to numerical methods and/or approximations to model diffusion within restricted geometries depending on the experimental conditions. Based on the new NMR diffusion equation, parts I and II of Chapter 1 intend to provide a very straightforward theoretical background for measuring diffusion of water protons and specific chemicals encountered in most common advanced MRI methods including diffusion MRI, functional MRI.

During the past decade, major breakthroughs in magnetic resonance imaging (MRI) quality were made by means of great improvement in scanner hardware and pulse sequences. Some advanced MRI techniques have truly revolutionized the detection of disease states and MRI can now-within a few minutes-acquire important quantitative information non-invasively from an individual in any plane or volume at comparatively high resolution. However, the very basic physics of this promising technological breakthrough is not well understood. In Chapter 2, parameters that are measured from time to time in advanced MRI seem to be logically and functionally related but the theoretical facility to optimally explore them is still missing. In a single experimental investigation, for example, few of huge amount of information available are effectively used. Parts I and II of this study intend to provide a very straightforward theoretical background for measuring diffusion of water protons and specific chemicals encountered in most common advanced MRI methods including diffusion MRI, perfusion MRI, functional MRI.

Chapter 3 considers the problem of Mobile Terminal (MT) positioning based on a time series of Received Signal Strength (RSS) measurements and provides statistical estimators of the MT- Base Station (BS) distance which substantially improve the MT positioning

accuracy. The set of MT-BS distance estimators defined for optimizing RSS based positioning when a single RSS measurement sample is available [1] is used as a basis. The problem that is analyzed refers to the adaptation of these estimators to the presence of Kalman filtering in the MT position calculation process. Three different Kalman filtering options can be applied at different stages of the MT position calculation process [2]: (a) RSS Kalman filtering, (b) MT-BS distance Kalman filtering and (c) MT position coordinates Kalman filtering. To identify the optimal MT-BS distance estimators at the presence of Kalman filtering a method exploiting the characteristics of the steady state Kalman filter performance is developed. This novel method allows for the definition of refined MT-BS distance that the resulting estimators provide good efficiency for all Kalman Filter options.

Let *R* be a commutative Noetherian ring which is also an algebra over \Box , \Box is the field of rational numbers). Let σ be an automorphism of *R* and δ a σ -derivation of *R* such that $\sigma(\delta(a)) = \delta(\sigma(a))$, for all $a \in R$. Chapter 4 concerns Ore extensions over near pseudo valuation rings (NPVR) and almost δ -divided rings. Towards this the authors prove:

- 1. If R is a near pseudo valuation ring, then $O(R) = R[x; \sigma, \delta]$ is a near pseudo valuation ring.
- 2. If R is an almost δ -divided ring, then $O(R) = R[x; \sigma, \delta]$ is an almost δ -divided ring.

In Chapter 5, the authors consider $p (\ge 2)$ independent treatment populations characterized by the unknown location parameters $\mu_1, ..., \mu_p$ and $q(\ge 2)$ independent control populations characterized by the unknown location parameters $\xi_1, ..., \xi_q$. The authors assume that θ is the common scale parameter of all the p+q populations. Let $\xi_{[q]}$ be the largest of ξs and define a treatment population to be "good" if its location parameter exceeds $\xi_{[q]} - \varepsilon$ for a given $\varepsilon > 0$. A selection procedure is proposed to select a subset of the *p* treatment populations which includes all the good treatment populations with probability at least P^* , where P^* is a pre-assigned value. Moreover, for given values of ε_1 and ε_2 ($\varepsilon_2 > \varepsilon_1 > 0$), the authors define the ith treatment population as "bad" if $\mu_i < \xi_q - \varepsilon_2$, i =1, ..., p. In this case a selection procedure is proposed and a sample size is determined so that the probability of omitting a "good" treatment population or selecting a "bad" treatment population is at most $1 - P^*$. Finally, the implementation of the proposed methodology is demonstrated through numerical example based on real life data.

In modern tennis, the essential part of executing high-performance shots is generating topspin and backspin on the ball. The purpose of Chapter 6 is to present a rigorous kinetic analysis of the flick-motion of the tennis racket, based on the original concept of the vector of mass/inertia moments coupled to the pole and for the corresponding axis. The forward racket flick-motion generates topspin on the ball, while the backward racket flick-motion generates backspin (also known as slice) on the ball. To describe both kinds of the racket flick-motion the authors use rigorous kinetic analysis. This advanced rigid-body analysis includes the six degrees-of-freedom (DOF) Newton-Euler dynamics, a new sophisticated form of

vectors/tensors of the racket mass inertia moments, impact forces during the racket-ball contact, and mass-deviational moment vectors of the racket and ball rotation before, during and after contact.

The motivation behind mathematically modeling the *human operator* is to help explain the response characteristics of the complex dynamical system including the human manual controller. In Chapter 7, the authors present two approaches to human operator modeling: classical linear control approach and modern nonlinear control approach. The latter one is formalized using both fixed and adaptive Lie-Derivative based controllers.

Suppose that $m \ge 4$ and that R is a commutative ring with identity in which 2 is invertible. Let N(m,R) be the nilpotent subalgebra of the orthogonal Lie algebra o(2m,R). In Chapter 8, the authors give an explicit description of the derivation algebra of N(m,R).

Historical development of algebra occurred in three stages; rhetorical or prose algebra, syncopated or abbreviated algebra and symbolic algebra - known as "school algebra". The analysis of Chapter 9 suggests that the first civilization to develop symbolic algebra was the Vedic Indians. The philosophical and religious ideas influenced the development of the decimal system and arithmetic and that led to algebra. Symbolic algebra appears to be deep rooted in Vedic philosophy. The Vedic mathematic were of a high level at an early period. The Hindus applied algebra freely creating formulas that simplified calculations. In geometry and trigonometry they developed formulas useful to understand the physical world satisfying the needs of religion (apara and para vidya). Geometrical focus, logic and proof type are features of Greek mathematics "boldness of conception, abstraction, symbolism" are evident in Indian mathematics. From history, a number of implications can be drawn. Real life, imaginative and creative problems that encourage risk should be the focus in student learning; allowing students freely move between symbols, numbers and magnitudes rather than taking a static unchanging view. Concrete, pictorial and symbolic modes are present in ancient learning. Real life practical, philosophical and religious needs in concert motivated progress to symbolic algebra. The historical analysis supports the use of rich context based problems that stimulate and motivate students to raise levels higher to transfer knowledge. The road from arithmetic to algebra was clearly in line with current emphasis in mathematics education but at an early stage in human history.

The purpose of Chapter 10 is to introduce the concept of regular fuzzy biclosure spaces and investigate some of their characterizations.

In Chapter 11 it is given the basic properties of a near-ring, which involve a set of matrix units and matrix near-rings and it is investigated the answer of the question "when the inclusion $Mat_n(R;R) \subseteq M_F(R^n)$ becomes an equality?" which posed in [1].

In Chapter 12 the authors introduce a new class of functions called contra- $\alpha \hat{g}$ -continuous functions and study some of their basic properties in topological spaces.

The purpose of Chapter 13 is to introduce a new class of functions called (1,2)*generalized α -continuous functions, (1,2)* α -generalized-continuous functions and (1,2)* α -strongly semi-continuous functions in bitopological spaces. Also the authors obtain some decompositions of (1,2)* α -continuous function in bitopological spaces.

Chapter 14 provides a simple exact expansion of a general function, while using constant coefficients. In addition, the point of expansion is not arbitrarily chosen.

Chapter 15 synthesizes and analyzes some important current and recent contributions to the theory of the firm under uncertainty. In so doing, it examines the production and hedging decisions of the competitive firm under a single source and multiple sources of uncertainty.

In Chapter 16 several properties and a characterization of δ_M -lifting modules are proved. Also the authors investigate the interconnections between δ_M -lifting, δ_M - supplemented modules and δ_M - semiperfect modules.

In Chapter 17, the authors apply Ghoussoub-Preiss's generalized Mountain Pass Lemma with Cerami-Palais-Smale type condition to study the existence of new periodic solutions with a given period for some second order Hamiltonian systems.

Chapter 18 deals with partial actions of inductive groupoids on rings. The authors establish a one-to-one correspondence between partial actions of an inductive groupoid G on a ring R, in which the domain of each partial bijection is an ideal of R, and meet-preserving global actions of the Birget-Rhodes expansion G^{BR} of G on R. Using this correspondence the authors obtain that the Birget-Rhodes expansion of the action groupoid $R \times_{\alpha} G$ and the action groupoid $R \times_{\beta} G^{BR}$ are isomorphic, where α is a partial action of G on a ring R and β is the correspondent action of G^{BR} on R. In particular, they get the equivalence of two suitable functors from the category of the partial actions of inductive groupoids into that of the ordered groupoids.

Chapter 1

MATHEMATICAL FORMULATION OF NMR EXPERIMENTAL PARAMETERS FOR DIFFUSION MAGNETIC RESONANCE IMAGING – PART I (CYLINDRICAL GEOMETRY)

M. Dada¹, O.P. Faromika², O.B. Awojoyogbe^{1,*}, M.A. Aweda³ and I.A. Fuwape²

¹Department of Physics, Federal University of Technology, Minna, Niger State, Nigeria ²Department of Physics, Federal University of Technology, Akure, Ondo State, Nigeria ³Department of Radiation Biology and Radiotherapy,

College of Medicine of the University of Lagos, Idi-Araba, Lagos State, Nigeria

Abstract

Magnetic resonance sequences have been developed to label the position of a spin, albeit indirectly, through the Larmore frequency. This provides the basis for measuring diffusion. However, the interpretation of the data is complicated by the effects of restricted geometries and the mathematical modeling required to account for this becomes nontrivial especially in cylindrical and spherical geometries. Generally one have to resort to numerical methods and/or approximations to model diffusion within restricted geometries depending on the experimental conditions. Based on the new NMR diffusion equation, parts I and II of this study intend to provide a very straightforward theoretical background for measuring diffusion of water protons and specific chemicals encountered in most common advanced MRI methods including diffusion MRI, functional MRI.

Keywords: Bloch NMR flow equations, Diffusion, Brownian motion, Perfusion, fMRI, Biological flow

^{*} E-mail address: awojoyogbe@yahoo.com

Chapter 2

MATHEMATICAL FORMULATION OF NMR EXPERIMENTAL PARAMETERS FOR DIFFUSION MAGNETIC RESONANCE IMAGING – PART II (CYLINDRICAL GEOMETRY)

O.B. Awojoyogbe^{1,*}, M. Dada¹, O.P. Faromika², M.A. Aweda³ and I.A. Fuwape²

¹Department of Physics, Federal University of Technology, Minna, Niger State, Nigeria ²Department of Physics, Federal University of Technology, Akure, Ondo State, Nigeria ³Department of Radiation Biology and Radiotherapy,

College of Medicine of the University of Lagos, Idi-Araba. Lagos State, Nigeria

Abstract

During the past decade, major breakthroughs in magnetic resonance imaging (MRI) quality were made by means of great improvement in scanner hardware and pulse sequences. Some advanced MRI techniques have truly revolutionized the detection of disease states and MRI can now-within a few minutes-acquire important quantitative information non-invasively from an individual in any plane or volume at comparatively high resolution. However, the very basic physics of this promising technological breakthrough is not well understood. Parameters that are measured from time to time in advanced MRI seem to be logically and functionally related but the theoretical facility to optimally explore them is still missing. In a single experimental investigation, for example, few of huge amount of information available are effectively used. Parts I and II of this study intend to provide a very straightforward theoretical background for measuring diffusion of water protons and specific chemicals encountered in most common advanced MRI methods including diffusion MRI, perfusion MRI, functional MRI.

Keywords: Bloch NMR flow equations, Diffusion, Brownian motion, Perfusion, fMRI, Biological flow

^{*}E-mail address: awojoyogbe@yahoo.com