



# MAPPING OF NATURAL GAMMA RADIATION (NGR) DOSE RATE DISTRIBUTION AROUND GIDAN-KWANO AREA, MINNA, NORTH-CENTRAL NIGERIA



H. O. Shittu<sup>1</sup>, I. O. Olarinoye<sup>2</sup>, M. T. Kolo<sup>2</sup>, A. N. Amadi<sup>3</sup>, S. F. Olukotun<sup>4</sup>, O. F. Oladejo<sup>5</sup> and G. E. K. Samuel<sup>1</sup>

<sup>1</sup>Department of Science Infrastructure, National Agency for Science & Engineering Infrastructure (NASeni), Abuja, Nigeria

<sup>2</sup>Department of Physics, Federal University of Technology, Minna, Nigeria

<sup>3</sup>Department of Geology, Federal University of Technology, Minna, Nigeria

<sup>4</sup>Department of Physics & Engineering Physics, Obafemi Awolowo University, Ile-Ife, Nigeria

<sup>5</sup>Department of Physics, Osun State University, Osogbo, Nigeria

\*Corresponding author: [olukotunsf@oauife.edu.ng](mailto:olukotunsf@oauife.edu.ng)

Received: August 05, 2021 Accepted: October 19, 2021

**Abstract:** An in-situ measurement of the background radiation level was carried out in Gidan-Kwano, Minna, Niger state with an objective to establish a reference data record on the levels and distribution of natural background gamma radiation (NGR). The measurement was carried out using a portable calibrated NaI(Tl) scintillation survey meter. A total of 153 points were surveyed across the study area for background environmental radiation. The results obtained, varied significantly due to geological features of the study area. NGR dose rates values ranged between 87 – 252.3 nGyh<sup>-1</sup> with overall mean value of 136.75 nGyh<sup>-1</sup>. This is more than twice higher than the reported world average value of 59 nGyh<sup>-1</sup>. However, the average annual effective dose obtained from this study is 0.17 mSv/annum, which is less than the recommended limit of 1 mSv/annum by International Commission on Radiation Protection [ICRP] for non-occupational population exposure. An isodose map for the distribution and exposure rate due to natural sources radiation for the study area was also plotted using ArcGIS software.

**Keywords:** Natural gamma radiation, dose rate, mean dose rate, isodose map

## Introduction

Natural background gamma radiation exists all around us. Men are continuously exposed to these radiations due to terrestrial and cosmic sources (Taskin *et al.*, 2009). According to United Nations Scientific Committee on the Effects of Atomic Radiation report (UNSCEAR), the greatest contribution to mankind's exposure comes from natural background radiation, and the global average annual effective dose is 2.4 mSv. An assessment of gamma radiation from natural sources is of particular importance (Kurnaz, 2013; Faanu *et al.*, 2011; Ramli *et al.*, 2009; Gbenu *et al.*, 2020a; 2020b; Oladejo *et al.*, 2020).

Radioactivity levels in environment vary greatly, depending on soil type and its mineral contents (Ramli *et al.*, 2009; Garba *et al.*, 2014; Saleh *et al.*, 2013). Uranium, thorium and potassium are all present in the Earth's crust in concentrations of 2 – 4 and 8 – 12 ppm and around 2.5 wt%, respectively. In rocks, they are present in the form of a number of minerals (Mares, 1984). Radiation measurements and potential risks of exposure to low-dose natural radioactivity have been of interest in recent times (Kapdan *et al.*, 2011; Isik *et al.*, 2012; Khan *et al.*, 2012; Gabdo *et al.*, 2015; Saleh *et al.*, 2015).

This study is aimed at producing a radiological map of the study area – Gidan-Kwano area, Minna, Niger state for the first time. This is essential due to the recent activities of local miners in the study area, and most especially now that the Federal Government of Nigeria has granted the Federal University of Technology, Minna a mining license. More so, the population of the study area has increased tremendously in recent times which is as a result of two major factors; the presence of Federal University of Technology, Minna, with its increasing enrolment over time. Hence, the data obtained will serve as background radiation level baseline to monitor the mining activity of the study area.

## Materials and Methods

### Study area

The study area – Gidan-Kwano area, is located on latitude 9° 30' N and 9° 34' N and longitude 6° 24' E and 6° 28' E. The study area was gridded into a total of 153 points and background environmental radiation survey was carried out at each point shown in Fig. 1. The study area is in Niger State,

which is covered by two major rock formations; the sedimentary and basement complex rocks. The sedimentary rocks to the south are characterized by sand stones and alluvial deposits, particularly along the Niger valley and in most parts of Borgu, Bida, Agaie, Lapai, Mokwa, Lavun, Gbako and Wushishi Local Government Areas. This subarea also contains the extensive flood plains of the River Niger and this has made the state to be one of the largest and most fertile agricultural lands in the country. To the north is the basement complex, characterized by granitic outcrops or inselbergs which can be found in the vast topography of rolling landscape. Such inselbergs dominate the landscape in Rafi, Shiroro, Minna, Mariga and Gurara (Ajibade, 1976). Fig. 2 gives the geological formations of the study area.




Fig. 1: Survey points of the study area




Fig. 2: Geological formations of the study area

**NGR dose rates measurement**

An in situ NGR dose rates measurement was conducted using a portable calibrated NaI(Tl) scintillation survey meter (Inspector Alert™ Nuclear Radiation Monitor, Serial Number – 35440) manufactured by International Medcom, Inc., USA. The meter uses (2.54 X 2.54 cm<sup>2</sup>) sodium iodide (NaI) crystal doped with thallium (Tl) as detector. Measurements were conducted randomly at 1m above soil surface at 153 different locations. Geographical coordinates of the measurement locations were recorded by Global Positioning System (GPS), Garmin eTrex 10 model. The meter displays dose rate reading in  $\mu\text{R}h^{-1}$  which was subsequently converted to  $\text{nGy}h^{-1}$  using a conversion factor  $1 \mu\text{R}h^{-1} \sim 8.7 \text{nGy}h^{-1}$  (Saleh *et al.*, 2013a). The relatively linear energy response of the detector between the gamma ray energies of 0.08 and 1.2 meV makes its excellent for field measurements (Knoll, 2010). Fig. 1 shows the measurement locations. To minimize errors, dose rate readings were recorded when the meter pointer was stable, and at least three set of readings were taken at a given point within the domain of the geological formations and soil types. Thereafter, the mean value for each location was computed from the set of the readings. To measure the actual NGR dose rates, measurements were made in an undisturbed open field and far away from mines and mining installations or facilities.

**Plotting of isodose map**

The data set on NGR dose rate measurements plus the coordinates for all data points were used in plotting an isodose map using ArcGIS version 10.3 – a mapping and spatial analysis software. The isodose map represents the distribution of NGR and exposure rates of the study area. Kriging technique was adopted for the plotting (Aziz Saleh *et al.*, 2014; Gerrard, 2000). This technique uses decreasing weight for the interpolation based on a semivariogram of survey points (Apriantoro, 2008). The datum used for the GPS was set up to the world geodetic system (WGS) 1984 and synchronized with coordinates of survey points.

**Results and Discussion**

The obtained NGR dose rates at each 153 points of the study area with their corresponding coordinates are given in Table 1 and the summary of the obtained dose rates is given in Table 2. The values were in the range of 87.0 – 252.3  $\text{nGy}h^{-1}$ , with a mean value of 136.75  $\text{nGy}h^{-1}$ . The obtained mean value is more than twice the world value of 59  $\text{nGy}h^{-1}$  reported by UNSCEAR (2000).

Table 1: Summary of the dose rate in soil samples

S/N	Longitudes	Latitudes	AVG( $\text{nGy}/h$ )
1	6° 24' 0'' E	9° 30' 0" N	121.80
2	6° 24' 15" E	9° 30' 0" N	113.10
3	6° 24' 30" E	9° 30' 0" N	182.70
4	6° 24' 45" E	9° 30' 0" N	113.10
5	6° 25' 0" E	9° 30' 0" N	139.20
6	6° 25' 15" E	9° 30' 0" N	113.10
7	6° 25' 30" E	9° 30' 0" N	121.80
8	6° 25' 45" E	9° 30' 0" N	113.10
9	6° 26' 0" E	9° 30' 0" N	130.50
10	6° 26' 15" E	9° 30' 0" N	130.50
11	6° 26' 30" E	9° 30' 0" N	104.40
12	6° 26' 45" E	9° 30' 0" N	121.80
13	6° 27' 0" E	9° 30' 0" N	121.80
14	6° 27' 15" E	9° 30' 0" N	121.80
15	6° 27' 30" E	9° 30' 0" N	130.50
16	6° 27' 45" E	9° 30' 0" N	113.10
17	6° 28' 0" E	9° 30' 0" N	156.60
18	6° 24' 0'' E	9° 30' 30" N	130.50
19	6° 24' 15" E	9° 30' 30" N	113.10
20	6° 24' 30" E	9° 30' 30" N	139.20
21	6° 24' 45" E	9° 30' 30" N	156.60
22	6° 25' 0" E	9° 30' 30" N	130.50
23	6° 25' 15" E	9° 30' 30" N	104.40
24	6° 25' 30" E	9° 30' 30" N	113.10
25	6° 25' 45" E	9° 30' 30" N	139.20
26	6° 26' 0" E	9° 30' 30" N	174.00
27	6° 26' 15" E	9° 30' 30" N	121.80
28	6° 26' 30" E	9° 30' 30" N	147.90
29	6° 26' 45" E	9° 30' 30" N	147.90
30	6° 27' 0" E	9° 30' 30" N	121.80
31	6° 27' 15" E	9° 30' 30" N	113.10
32	6° 27' 30" E	9° 30' 30" N	130.50
33	6° 27' 45" E	9° 30' 30" N	104.40
34	6° 28' 0" E	9° 30' 30" N	121.80
35	6° 24' 0'' E	9° 31' 0" N	121.80
36	6° 24' 15" E	9° 31' 0" N	182.70
37	6° 24' 30" E	9° 31' 0" N	191.40
38	6° 24' 45" E	9° 31' 0" N	147.90
39	6° 25' 0" E	9° 31' 0" N	139.20
40	6° 25' 15" E	9° 31' 0" N	147.90
41	6° 25' 30" E	9° 31' 0" N	113.10
42	6° 25' 45" E	9° 31' 0" N	139.20
43	6° 26' 0" E	9° 31' 0" N	113.10
44	6° 26' 15" E	9° 31' 0" N	104.40
45	6° 26' 30" E	9° 31' 0" N	0.00
46	6° 26' 45" E	9° 31' 0" N	121.80
47	6° 27' 0" E	9° 31' 0" N	0.00
48	6° 27' 15" E	9° 31' 0" N	147.90
49	6° 27' 30" E	9° 31' 0" N	121.80
50	6° 27' 45" E	9° 31' 0" N	130.50
51	6° 28' 0" E	9° 31' 0" N	191.40
52	6° 24' 0" E	9° 31' 30" N	182.70
53	6° 24' 15" E	9° 31' 30" N	139.20
54	6° 24' 30" E	9° 31' 30" N	147.90
55	6° 24' 45" E	9° 31' 30" N	121.80
56	6° 25' 0" E	9° 31' 30" N	156.60
57	6° 25' 15" E	9° 31' 30" N	130.50
58	6° 25' 30" E	9° 31' 30" N	147.90

*Determination of Levels & Distribution of Natural Gamma Radiation around Gidan-Kwano, Minna*

59	6 25 45 E	9 31 30 N	104.40	128	6 26 0 E	9 33 30 N	87.00
60	6 26 0 E	9 31 30 N	139.20	129	6 26 15 E	9 33 30 N	113.10
61	6 26 15 E	9 31 30 N	113.10	130	6 26 30 E	9 33 30 N	156.60
62	6 26 30 E	9 31 30 N	130.50	131	6 26 45 E	9 33 30 N	147.90
63	6 26 45 E	9 31 30 N	217.50	132	6 27 0 E	9 33 30 N	130.50
64	6 27 0 E	9 31 30 N	217.50	133	6 27 15 E	9 33 30 N	165.30
65	6 27 15 E	9 31 30 N	104.40	134	6 27 30 E	9 33 30 N	191.40
66	6 27 30 E	9 31 30 N	130.50	135	6 27 45 E	9 33 30 N	0.00
67	6 27 45 E	9 31 30 N	156.60	136	6 28 0 E	9 33 30 N	121.80
68	6 28 0 E	9 31 30 N	200.10	137	6 24' 0''E	9 340 N	165.30
69	6 24' 0''E	9 320 N	156.60	138	6 <sup>0</sup> 24 15 E	9 34 0 N	191.40
70	6 <sup>0</sup> 24 15 E	9 32 0 N	121.80	139	6 <sup>0</sup> 24 30 E	9 34 0 N	165.30
71	6 <sup>0</sup> 24 30 E	9 32 0 N	104.40	140	6 24 45 E	9 34 0 N	156.60
72	6 24 45 E	9 32 0 N	113.10	141	6 25 0 E	9 34 0 N	147.90
73	6 25 0 E	9 32 0 N	121.80	142	6 25 15 E	9 34 0 N	208.80
74	6 25 15 E	9 32 0 N	121.80	143	6 25 30 E	9 34 0 N	165.30
75	6 25 30 E	9 32 0 N	156.60	144	6 25 45 E	9 34 0 N	165.30
76	6 25 45 E	9 32 0 N	147.90	145	6 26 0 E	9 34 0 N	147.90
77	6 26 0 E	9 32 0 N	104.40	146	6 26 15 E	9 34 0 N	182.70
78	6 26 15 E	9 32 0 N	121.80	147	6 26 30 E	9 34 0 N	156.60
79	6 26 30 E	9 32 0 N	156.60	148	6 26 45 E	9 34 0 N	147.90
80	6 26 45 E	9 32 0 N	139.20	149	6 27 0 E	9 34 0 N	139.20
81	6 27 0 E	9 32 0 N	156.60	150	6 27 15 E	9 34 0 N	191.40
82	6 27 15 E	9 32 0 N	121.80	151	6 27 30 E	9 34 0 N	165.30
83	6 27 30 E	9 32 0 N	182.70	152	6 27 45 E	9 34 0 N	113.10
84	6 27 45 E	9 32 0 N	139.20	153	6 28 0 E	9 34 0 N	113.10
85	6 28 0 E	9 32 0 N	147.90				
86	6 24 0E	9 32 30 N	252.30				
87	6 <sup>0</sup> 24 15 E	9 32 30 N	182.70				
88	6 <sup>0</sup> 24 30 E	9 32 30 N	130.50				
89	6 24 45 E	9 32 30 N	121.80				
90	6 25 0 E	9 32 30 N	174.00				
91	6 25 15 E	9 32 30 N	130.50				
92	6 25 30 E	9 32 30 N	139.20				
93	6 25 45 E	9 32 30 N	130.50				
94	6 26 0 E	9 32 30 N	165.30				
95	6 26 15 E	9 32 30 N	139.20				
96	6 26 30 E	9 32 30 N	121.80				
97	6 26 45 E	9 32 30 N	95.70				
98	6 27 0 E	9 32 30 N	147.90				
99	6 27 15 E	9 32 30 N	139.20				
100	6 27 30 E	9 32 30 N	130.50				
101	6 27 45 E	9 32 30 N	147.90				
102	6 28 0 E	9 32 30 N	95.70				
103	6 24' 0''E	9 330 N	147.90				
104	6 <sup>0</sup> 24 15 E	9 33 0 N	147.90				
105	6 <sup>0</sup> 24 30 E	9 33 0 N	113.10				
106	6 24 45 E	9 33 0 N	147.90				
107	6 25 0 E	9 33 0 N	182.70				
108	6 25 15 E	9 33 0 N	139.20				
109	6 25 30 E	9 33 0 N	156.60				
110	6 25 45 E	9 33 0 N	174.00				
111	6 26 0 E	9 33 0 N	147.90				
112	6 26 15 E	9 33 0 N	0.00				
113	6 26 30 E	9 33 0 N	174.00				
114	6 26 45 E	9 33 0 N	165.30				
115	6 27 0 E	9 33 0 N	0.00				
116	6 27 15 E	9 33 0 N	0.00				
117	6 27 30 E	9 33 0 N	121.80				
118	6 27 45 E	9 33 0 N	113.10				
119	6 28 0 E	9 33 0 N	139.20				
120	6 24' 0''E	9 33 30 N	130.50				
121	6 <sup>0</sup> 24 15 E	9 33 30 N	139.20				
122	6 <sup>0</sup> 24 30 E	9 33 30 N	147.90				
123	6 24 45 E	9 33 30 N	200.10				
124	6 25 0 E	9 33 30 N	174.00				
125	6 25 15 E	9 33 30 N	147.90				
126	6 25 30 E	9 33 30 N	191.40				
127	6 25 45 E	9 33 30 N	113.10				

**Table 2: Summary of the basic statistics for external gamma dose rates**

Statistics	Dose rate (nGy/h)
Mean	136.75
Range	87.0 – 252.3
SE	3.18
SD	39.32
Median	139.20
Mode	147.90
World average	59

**Table 3: Mean dose rate for this study compared to other countries of the world**

S/N	Country/Region	Dose rate (nGy/h)	References
1	Southwest, Nigeria	232	Jibiri <i>et al.</i> (2016)
2	Ghana	741	Faanu <i>et al.</i> (2016)
3	Jos Plateau	250	Abba <i>et al.</i> (2017)
4	Minna	154	Olarinoye <i>et al.</i> (2010)
5	Malaysia	209	Nuraddeen <i>et al.</i> (2015)
6	Portugal	84	UNSCEAR (2000)
7	USA	47	UNSCEAR (2000)
8	India	56	UNSCEAR (2000)
9	Iran	105	Baykara and Dogru (2009)
10	Brazil	125	Freitas and Alencar (2004)
11	Spain	76	UNSCEAR (1988)
12	World average	59	UNSCEAR (2000)
13	Gidan-Kwano, Minna	136.75	This study

The highest dose rate was recorded due to the soil type underlain by granitic rock formation while the lowest was recorded because of the sandstone, clay and shale formations which were underlain by sedimentary rocks. Soils derived from granitic parent material are known to contribute to higher dose rates (UNSCEAR, 2000) compared to soils developed as a result of decomposition of organic matter such as peat, muck and shale (Sanusi *et al.* 2014). This is because; the minerals that carry U and Th are generally associated with felsic intrusive rocks, particularly with younger granites compared to ultramafic and volcanic rocks (Amadi *et al.*,

2012). The results of this study are in general agreement with similar studies conducted by Ramli *et al.* (2009), Saleh *et al.* (2013a and b), Lee (2009), and Garba *et al.*, (2015) who reported dose rates of higher values for soils of igneous origin. Dose rate of as low as  $8.7 \text{ nGy}^{-1}$  was reported by Tzortzis *et al.* (2004) for the soil of sedimentary rocks in Cyprus. The total mean dose rate of the surveyed area is found to be more than twice that of the world average, and found to be higher

than that of other places (Olarinoye *et al.*, 2010; Jibiri *et al.*, 2016; Faanu *et al.*, 2016; Abba *et al.*, 2017) as shown Table 3. Figs. 3 and 4 show the isodose map of the study area and the frequency distribution curve for dose rates, respectively. The NGR distribution of the study area is determined majorly by the basement rocks and human activities.




Fig. 3: Isodose map of the NGR dose rates




Fig. 4: Frequency distribution curve for dose rates



### **Estimation of annual effective dose (AED)**

The obtained mean dose rate was used to compute the mean indoor and outdoor annual effective doses due to exposure to natural sources of background gamma radiation. The parameters were estimated by assuming the conversion coefficient for the absorbed dose in air to effective dose of 0.7 Sv Gy<sup>-1</sup>, and the indoor and outdoor occupancy factors of 0.8 and 0.2, respectively, as recommended by UNSCEAR (2000). The indoor and outdoor annual effective dose equivalent was estimated using Equation 1 and 2, respectively.

$$AED_{in}(\text{mSv/y}) = \text{mean dose rate (nGy/h)} * 24(\text{h}) * 365(\text{days}) * 0.8 * 0.7 * 10^{-6} \quad (1)$$

$$AED_{ext}(\text{mSv/y}) = \text{mean dose rate (nGy/h)} * 24(\text{h}) * 365(\text{days}) * 0.2 * 0.7 * 10^{-6} \quad (2)$$

Assuming the population in the study area spend 20% of their day in this area (UNSCEAR, 2000), the obtained annual effective dose is 0.17 mSv/annum. The obtained value is lower than the recommended limit of 1 mSv/annum by International Commission on Radiation Protection (ICRP, 1990).

### **Conclusion**

The high background radiation observed in this study area could be attributed only to natural sources (cosmic and terrestrial). The geology of the town also suggests that the soil in Minna town has a large deposit of granite. It is well known that granites contain high concentrations of uranium, thorium and potassium (Ivanovich and Harmon, 1982). The overall mean value for this study is computed to be 136.75 nGy<sup>-1</sup> with a standard deviation of 39.32 nGy<sup>-1</sup>. This value falls within the highest range of those measured at worldwide scale by other authors and, more specifically, is by a factor of more than twice times higher than the reported world average values of 59 nGy<sup>-1</sup> in the UNSCEAR (2000) report. The mean indoor and outdoor annual effective doses for the public were estimated to be 0.67 and 0.17 mSv/y, respectively, which are less than the dose limit recommended by the ICRP. The isodose map for the distribution NGR and exposure rate for the study area was also plotted using ArcGIS software. Thus, there is a need for a comprehensive radiological study in the areas covered by this work to ascertain the radionuclide responsible for the elevated gamma dose rates.

### **Acknowledgement**

The authors express their gratitude to the entire staff of the Centre for Energy Research and Development (CERD), Obafemi Awolowo University, Ile-Ife, Nigeria especially the members of Gamma Spec Laboratory for their assistance.

### **Conflict of Interest**

The authors declare that there is no conflict of interest related to this study.

### **References**

Abba HT, Saleh MA, Hassan WMSW, Aliyu AS & Ramli AT 2017. Mapping of natural gamma radiation (NGR) dose rate distribution in tin mining areas of Jos Plateau, Nigeria. *Environ Earth Sci.*, 76: 208. DOI 10.1007/s12665-017-6534-8

Ajibade AC 1976. Provisional classification and correlation of the schist belt in Northwestern Nigeria. Elizabethan Publishing Co, Ibadan, pp. 85- 87.

Amadi A, Okoye N, Olasehinde P, Okunlola I, Alkali Y, Ako T & Chukwu J 2012. Radiometric survey as a useful tool in geological mapping of Western Nigeria. *J. Geogr. Geol.* 4: 242

Apriantoro N 2008. Radiological study in Perak state and its radiological health impact. Universiti Teknologi Malaysia Doctor of Philosophy Thesis.

Aziz Saleh M, Termizi Ramli A, Alajerami Y, Damoom M & Sadiq Aliyu A 2014. Assessment of health hazard due to natural radioactivity in Kluang District, Johor, Malaysia. *Isot Environ Health Stud.* 50: 103–113.

Baykara O & Dogru M 2009. Determination of terrestrial gamma, 238 U, 232 Th and 40 K in soil along fracture zones. *Radiat. Meas.*, 44: 116–121.

Faanu A *et al.* 2016. Natural radioactivity levels in soils, rocks and water at a mining concession of Perseus gold mine and surrounding towns in Central Region of Ghana. *Springer Plus.*, 5: 98.

Faanu A, Darko E & Ephraim J 2011. Determination of natural radioactivity and hazard in soil and rock samples in a mining area in Ghana. *West Afr. J. Appl. Ecol.*, 19: 77–92.

Freitas A & Alencar A 2004. Gamma dose rates and distribution of natural radionuclides in sand beaches—Ilha Grande, Southeastern Brazil. *J. Environ Radioact.*, 75: 211–223.

Gabdo HT, Ramli AT, Saleh MA, Sanusi MS & Garba NN 2015. The influence of geology on terrestrial gamma radiation dose rate in Pahang state, Malaysia. *Curr. Sci.*, 109: 177–183.

Garba N, Ramli A, Saleh M, Sanusi M, Gabdo H (2015) Terrestrial gamma radiation dose rates and radiological mapping of Terengganu state, Malaysia. *J. Radioanal Nucl. Chem.*, 303: 1785–1792.

Gbenu ST, Makinde OW, Ajenifuja E, Olukotun SF, Fasasi MK, Balogun FA & Ogunde KT 2020a. Ground radiometric mapping of Ondo State Nigeria using Gamma ray spectrometry and geographic information system. *Ife J. Sci.*, 22: 139–147.

Gbenu ST, Olukotun SF, Oladejo OF, Onumojor CA, Balogun FO, Ibitoye FI & Fasasi MK 2020b. Assessment of the elemental concentration of potassium, uranium and thorium with their associated radiogenic heat production in granite rocks of Southwestern Nigeria. *J. Radiat. Nucl. Appl. An Int. J.*, 5: 205–210. <https://doi.org/10.18576/jma/050307>

Gerrard J 2000. Fundamentals of Soils. Psychology Press, London.

Grant F 1982. Gamma ray spectrometry for geological mapping and for prospecting. In: Mining Geophysics Workshops, pp. 1311–1327.

ICRP 1991. International Commission on Radiological Protection. Publication 60, (1990). Recommendations of the International Commission on Radiological Protection, Annals of ICRP vol. 21, No 13 Pergamon Press, Elmsford, NY.

Isik U, Damla N, Akkoca DB & Cevik U 2012. Mineralogical, geochemical and radiological characterisation of Selmo Formation in Batman area, Turkey. *Isot. Environ Health Stud.*, 48:302–312.

Ivanovich, M & Harmon RS 1982. Uranium series disequilibrium in application to environmental problems. Oxford, Clarendon Press, 51-55.

Jibiri N, Isinkaye M, Bello I & Olaniyi P 2016. Dose assessments from the measured radioactivity in soil, rock, clay, sediment and food crop samples of an elevated radiation area in south-western Nigeria. *Environ Earth Sci.*, 75: 107.

Kapdan E, Altinsoy N, Karahan G & Taskin H 2011. Determination of the health hazards due to background radiation sources in the city of Adapazari, Northwestern Turkey. *Isot. Environ Health Stud.*, 47:93–100.

- Khan HM, Ismail M, Zia MA, Khan K 2012. Measurement of radionuclides and absorbed dose rates in soil samples of Peshawar, Pakistan, using gamma ray spectrometry. *Isot. Evt. Health Stud.*, 48: 295–301.
- Knoll GF 2010. Radiation Detection and Measurement. Wiley, London.
- Kurnaz A 2013. Background radiation measurements and cancer risk estimates for Sebinkarahisar, Turkey. *Radiat. Prot. Dosim.* doi:10.1093/rpd/nct115
- Lee SK, Ramli AT, Wagiran H, Apriantoro NH & Wood AK 2009. Radiological monitoring: Terrestrial natural radionuclides in Kinta District, Perak, Malaysia. *J. Environ. Radioact.* 100: 368e374.
- Mares S 1984. Introduction to Applied Geophysics. Dordrecht, NL: Springer Science & Business Media.
- Nuraddeen Nasiru Garba, Ahmad Termizi Ramli, Muneer Aziz Saleh, Syazwan Mohd Sanusi & Hamman Tukur Gabdo 2015. Radiological mapping of Kelantan, Malaysia using terrestrial radiation dose rate. *Isotopes in Evtal. and Health Stud.*, DOI: 10.1080/10256016.2016.1095189
- Oladejo OF, Olukotun SF, Ogundele LT, Gbenu ST & Fakunle MA 2020. Radiological risk assessment of naturally occurring radioactive materials (NORMS) from selected quarry sites in Edo State, South-south, Nigeria. *Environ. Earth Sci.*, 79: 1–8. <https://doi.org/10.1007/s12665-020-8842-7>
- Olarinoye IO, Sharifat I, Baba-Kutigi AN, Kolo MT & Aladeniyi K 2010. Measurement of Background Gamma Radiation Levels at Two Tertiary Institutions in Minna, Nigeria. *J. Appl. Sci. Environ. Mgt. March*, 14(1): 59 – 62. Available Online at [www.bioline.org.br/ja](http://www.bioline.org.br/ja)
- Ramli AT, Apriantoro NH & Wagiran H 2009. Assessment of radiation dose rates in the high terrestrial gamma radiation area of Selama District, Perak, Malaysia. *Appl. Phys. Res.*, 1: 45.
- Saleh MA, Ramli AT, Alajerami Y & Aliyu AS 2013a. Assessment of environmental 226 Ra, 232 Th and 40 K concentrations in the region of elevated radiation background in Segamat District, Johor, Malaysia. *J. Environ Radioact.*, 124: 130–140.
- Saleh MA, Ramli AT, Alajerami Y, Aliyu AS & Basri NAB 2013b. Radiological study of Mersing District, Johor, Malaysia. *Radiat. Phys. Chem.*, 85: 107–117.
- Saleh MA, Ramli AT, bin Hamzah K, Alajerami Y, Moharib M, Saeed I 2015. Prediction of terrestrial gamma dose rate based on geological formations and soil types in the Johor State, Malaysia. *J. Environ Radioact.*, 148:111–122.
- Sanusi M, Ramli A, Gabdo H, Garba N, Heryanshah A, Wagiran H & Said M 2014. Isodose mapping of terrestrial gamma radiation dose rate of Selangor state, Kuala Lumpur and Putrajaya, Malaysia. *J. Environ Radioact.*, 135: 67–74.
- Taskin H, Karavus M, Ay P, Topuzoglu A, Hidiroglu S & Karahan G 2009. Radionuclide concentrations in soil and lifetime cancer risk due to gamma radioactivity in Kirklareli, Turkey. *J. Environ Radioact.*, 100: 49–53.
- Tzortzis M & Tsertos H 2004. Determination of thorium, uranium and potassium elemental concentrations in surface soils in Cyprus. *J. Environ Radioact.*, 77: 325–338.
- UNSCEAR (2000) Sources and effects of ionizing radiation. United Nations, New York
- UNSCEAR 1988. United Nations Scientific Committee on the Effects of Atomic Radiation. Sources Effects and Risks of Ionizing Radiation. Report to the General Assembly, with Annexes. New York, USA: United Nations Publication, pp. 49-51.
- UNSCEAR 1998. Sources, Effects and Risk of Ionizing Radiation. United Nations Scientific Committee on the Effect of Atomic Radiation. New York, United Nations