

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/303913277>

ASSESSMENT OF NATURAL RADIOACTIVITY LEVELS AND RADIATION HAZARDS IN THE TERTIARY INSTITUTIONS IN MINNA, NIGER STATE, NIGERIA

Article · June 2012

DOI: 10.5707/cjenvsci.2012.6.3.25.31

CITATIONS

10

READS

317

2 authors:



Kolo Matthew

Federal University of Technology Minna

19 PUBLICATIONS 330 CITATIONS

[SEE PROFILE](#)



Oyeleke Ismail Olarinoye

Federal University of Technology Minna

186 PUBLICATIONS 3,084 CITATIONS

[SEE PROFILE](#)

ASSESSMENT OF NATURAL RADIOACTIVITY LEVELS AND RADIATION HAZARDS IN THE TERTIARY INSTITUTIONS IN MINNA, NIGER STATE, NIGERIA.

Kolo, M. T., Baba-Kutigi, A. N., Olarinoye, I. O. and Sharifat, I.

Department of Physics, Federal University of Technology, Minna, Niger State, Nigeria.

ABSTRACT

Activity concentrations of natural radionuclides in 30 surface soil samples collected across the three campuses of the two tertiary institutions in Minna, Niger State, Nigeria, were studied and evaluated. This survey was carried out using gamma spectrometric technique which employs NaI(Tl) gamma detector at the Center for Energy Research and Training (CERT), Ahmadu Bello University, Zaria, Nigeria. The mean values for the activity of ^{226}Ra , ^{232}Th and ^{40}K were found to be $7.8\pm 1.3\text{Bqkg}^{-1}$, $29.4\pm 0.9\text{Bqkg}^{-1}$ and $229.4\pm 1.8\text{Bqkg}^{-1}$ respectively. The activity profile of the primordial radionuclides in the soil samples investigated showed the existence of low level activity across the three campuses. The mean value of the annual effective dose equivalent obtained from this study is 0.04mSvyr^{-1} , with mean external hazard index of 0.2. These average values fall within the internationally provided safety range for outdoor radiation exposure. The values obtained from this investigation for all the radiation parameters for the studied soil samples showed that none of the campuses investigated pose any significant radiological threat to the public.

KEYWORDS: Natural radioactivity, gamma ray spectrometry, annual effective dose equivalent, external hazard index, Federal University of Technology Minna, Niger State College of Education, Minna.

INTRODUCTION

Naturally Occurring Radioactive Materials (NORMs) are and have always been a part of our world. Radiation is present in every environment of the earth's surface, beneath the earth, in the atmosphere (Kannan *et al.*, 2002), and even in our human body. Various geological formations, including rocks, plants, water, air and soils contain NORMs at various trace levels. According to Karahan and Bayulken (2000), natural radionuclides in soils generate a significant component of the background radiation exposure of the population. Soils not only acts as a source of continuous radiation exposure to humans, but also as a medium of migration for transfer of radionuclides to biological systems (Mehra *et al.*, 2010), and hence, the basic indicator of radiological contamination in the environment (Ibrahim and Mohammad, 2009).

Humans are constantly exposed to natural radiation from two prominent sources: internal exposure from radionuclides in food and inhaled radon gas; and the external radiation exposure from cosmic rays and terrestrial gamma rays due to NORMs in soils, rocks and building materials (Alberto, Laura and Valerio, 1996; Alatisse, Babalola and Olowofela, 2008). The earth and the atmosphere contains various levels of radionuclides, but only those with half lives comparable with the age of the earth and their corresponding decay products existing in terrestrial materials, such as ^{40}K , ^{238}U and ^{232}Th are of great interest, since gamma radiation from these represents the main external source of human exposure. The level of natural radioactivity in soils and in the surrounding environments as well as the associated external exposure due to gamma radiation depends primarily on the geological and geophysical conditions of the region (UNSCEAR, 1998; 2000). Since these radionuclides are not uniformly distributed, the knowledge of their distribution in soils and their measurement is very important, not only to determine the amount of change of the natural background activity with time, but also for radiation protection issues.

Minna, the capital of Niger State, is the home of two tertiary institutions of the state: the Federal University of Technology, and Niger State College of Education. Minna and the entire state are covered by two major rock formations-the sedimentary and basement complex rocks. Other secondary geological formations in the state may suggest a radiation level slightly above background. Data on the natural radioactivity concentrations in Minna city and the entire state are still scarce and so a comprehensive study with the objective to systematically

measure the terrestrial gamma radiation and determine its contribution to annual effective dose equivalent to the population in these campuses is highly needed.

In order to assess the radiological hazards due to natural radioactivity of soils of these campuses, calculations of the radium equivalent activity, absorbed gamma dose rate in air out doors and the external hazard index are presented and discussed. Hopefully, the results of this study and the interpretations presented would serve as base line data for radioactivity levels in this environment.

MATERIALS AND METHODS

Surface soil samples were collected from the three campuses of the tertiary institutions. These are Bosso campus (BSC) and Gidan Kwano campus (GKC) of Federal University of Technology, and the Niger State College of Education (COE) campus. Ten surface soil samples were collected per campus. The choice of the sampling points in the campuses, which include, the boys' and girls' hostels, the central mosques, the chapels, the administrative blocks, libraries and the sports complex, was informed by the daily extensive presence of human life and the existence of portable water sources.

Three sub samples per each sample point, at 5cm depth, were collected after removing the stones and vegetation. These sub samples were thoroughly mixed together to give a correct representation of each sample point. The soil samples, neatly packed in well labeled polythene bags were air-dried for 48 hours under normal ambient temperature to remove the moisture content. The dried soil samples were crushed; grind into fine powder to pass through 500 μ m sieve. The representative powder samples were filled into cylindrical plastic containers of dimension 7.2cm in diameter and 6.0cm high to satisfy the selected optimal sample container height. The sample containers were then sealed hermetically with adhesive tape and stored for 24 days to allow them attain radioactive equilibrium. This was necessary to ensure that radon gas is confined within the volume and that the daughters will also remain in the sample (Sroor et al., 2001). Each container accommodated average of 275 \pm 1g.

The gamma ray spectrometry system used for the measurement consist of 7.62cmx7.62cm NaI(Tl) detector at the Center for Energy Research and Training (CERT), Ahmadu Bello University Zaria, Kaduna State, Nigeria. The detector is housed in a 6cm thick lead shield so as to reduce the background radiation levels many times. The inside of the detector is also lined with cadmium and copper sheets, which according to El-Ayadarous (2007), help to absorb the emitted x-rays from lead which may contain radioactive impurities due to antimony impurities. The detector assemblage is coupled to a computer based multichannel analyzer (MCA) with ACCUSPEC computer program used for data acquisition and analysis of gamma spectra. The efficiency and energy calibration of the detector were done over energy range using ^{137}Cs , ^{60}Co and ^{54}Mn standard isotopic sources over energy range of 200keV to 3MeV, being the energy range of radionuclides to be determined. Also, the IAEA gamma spectrometric reference materials, RGK-1, RGU-1 and RGTh-1 were used to recalibrate the system for quantitative determination of ^{40}K , ^{226}Ra and ^{232}Th in the soil samples. The activity concentration of ^{226}Ra was evaluated from 1764keV gamma line of ^{214}Bi , while 2614keV gamma line of ^{208}Tl was used to evaluate the activity concentration of ^{232}Th . The single 1460keV gamma line of ^{40}K was used for its content evaluation. Each sample was counted for 29000 seconds.

It is known that the distribution of ^{226}Ra , ^{232}Th and ^{40}K in soil is heterogeneous in nature with no specific relation between them (Christa et al., 2011). A common index with respect to radiation exposure, called the radium equivalent activity (Ra_{eq}), has therefore been defined to represent the specific activities of ^{226}Ra , ^{232}Th and ^{40}K in different combinations in soil samples. This index is calculated using the relation (Beretka and Matthew, 1985; Yu et al., 1992; UNSCEAR, 2000)

$$Ra_{eq} (\text{Bqkg}^{-1}) = C_{Ra} + 1.43C_{Th} + 0.077C_K \quad (1)$$

Where C_{Ra} , C_{Th} , and C_K are activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K in Bqkg^{-1} respectively. Ra_{eq} is defined according to equation (1) on the assumption that 1 Bqkg^{-1} of ^{226}Ra , 0.7 Bqkg^{-1} of ^{232}Th or 13 Bqkg^{-1} of ^{40}K yields the same gamma ray dose (Malanca et al., 1993; OECD, 1979).

From the measured activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K in the soil samples, the external gamma absorbed dose rate in air 1m above ground level was calculated following the equation (UNSCEAR, 2000)

$$D \text{ (nGyh}^{-1}\text{)} = 0.462C_{\text{Ra}} + 0.621C_{\text{Th}} + 0.417C_{\text{K}} \quad (2)$$

Where D is the dose rate in nGyh⁻¹, C_{Ra}, C_{Th}, C_K (Bqkg⁻¹), and 0.462, 0.621, and 0.417 (ngyh⁻¹ per Bqkg⁻¹) are the activity concentrations and dose conversion factors of ²²⁶Ra, ²³²Th and ⁴⁰K respectively. In equation (2) above, it is assumed that all decay products of ²²⁶Ra and ²³²Th are in radioactive equilibrium with their precursors (Prasong and Susaira, 2008).

The annual effective dose equivalent to the students population in the campuses due to soil radioactivity was estimated using the equation (UNSCEAR, 2000)

$$\begin{aligned} \text{AED (mSvyr}^{-1}\text{)} &= D \text{ (nGyh}^{-1}\text{)} \times 0.7 \text{ SvGy}^{-1} \times 8760 \text{ hr}^{-1} \times \left(\frac{10^3 \text{ mSv}}{10^9} \right) \times 0.2 \\ &= D \text{ (nGyh}^{-1}\text{)} \times 1.2264 \times 10^{-3} \end{aligned} \quad (3)$$

Equation (3) was calculated using the dose conversion factor, 0.7SvGy⁻¹, from absorbed dose rate in air to effective dose received by adults, and the occupancy factor of 0.2 proposed by UNSCEAR (2000), bearing in mind that the students spent an average of 20% of their time outdoors.

The external hazard index (H_{ex}), which is an estimate of radiation risk resulting from exposure to gamma rays of the primordial nuclei is evaluated from the equation (UNSCEAR, 2000)

$$H_{\text{ex}} = \frac{C_{\text{Ra}}}{370} + \frac{C_{\text{Th}}}{259} + \frac{C_{\text{K}}}{4810} \quad (4)$$

Where C_{Ra}, C_{Th}, and C_K are activity concentrations of ²²⁶Ra, ²³²Th and ⁴⁰K in Bqkg⁻¹ respectively. The maximum value for H_{ex} is equal to unity, which corresponds to 370Bqkg⁻¹, the upper limit of Ra_{eq}. The value of this index must therefore be less than unity so as to keep the radiation hazard insignificant.

RESULTS AND DISCUSSION

The results of the gamma spectrometry analysis for the 30 soil samples collected across the three campuses of the tertiary institutions in Minna, Niger state are presented in Table 1.

Table 1 shows that the activity concentrations in Bqkg⁻¹ of ²²⁶Ra, ²³²Th and ⁴⁰K in all the samples analyzed ranged between 1.0±2.3–26.5±1.9, 2.2±0.3–70.3±0.9, and 43.1±3.1–468.5±3.0, with mean values of 7.8±1.3, 29.4±0.9 and 229.4±1.8 respectively. Although the mean value of ⁴⁰K in all the measured samples is higher than those of ²²⁶Ra and ²³²Th as can be seen from the table, the obtained results are much lower compared with the world wide average concentration of these radionuclides in soils as reported by UNSCEAR (2000), which are 40Bqkg⁻¹ for ²²⁶Ra, 40Bqkg⁻¹ for ²³²Th and 370Bqkg⁻¹ for ⁴⁰K. the results therefore shows that the three campuses being investigated have normal levels of natural background and so do not pose any risk to the students.

Table 2 presents a comparison of the mean values of activity concentrations of the natural radionuclides in soil samples of the studied area with those of other literatures and the world wide mean. The comparison shows that the results obtained from this study are in agreement with similar works from other countries.

The values calculated for radium equivalent activity, gamma dose rate, annual effective dose and external hazard index are presented in table 3. The values for Ra_{eq} for all the samples as seen in Table 3 ranged from 11.6 to 131.9Bqkg⁻¹, with a mean value of 67.5Bqkg⁻¹. The values are less than the safe limit of 370Bqkg⁻¹ provided by UNSCEAR (2000). The values for the absorbed gamma dose rate in air fluctuates between 5.6nGyh⁻¹ and 59.7nGyh⁻¹ with an average value of 31.3nGyh⁻¹, while the calculated values for annual effective dose equivalent ranged between 0.01 and 0.07mSvyr⁻¹, with a mean value of 0.04mSvyr⁻¹. These values are found to be comparable to the world average of 57nGyh⁻¹ and 0.48mSvyr⁻¹ respectively provided by UNSCEAR (2000). The mean value for the external hazard index as seen in table 3 is 0.2, which is lower than unity, indicating that the radiation hazard in the three campuses is really insignificant.

Table 1: Activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K in all the soil samples (Bq/kg)

| SAMPLE ID | ^{226}Ra | ^{232}Th | ^{40}K |
|-----------|-------------------|-------------------|--------------------|
| BSC1 | 3.5±0.5 | 8.0±0.1 | 357.3±2.2 |
| BSC2 | 21.0±0.1 | 20.2±0.6 | 292.7±1.2 |
| BSC3 | 1.4±2.2 | 44.5±9.8 | 437.4±0.8 |
| BSC4 | 13.3±0.8 | 14.0±0.1 | 395.0±1.7 |
| BSC5 | 1.0±2.3 | 30.3±1.8 | 286.0±1.6 |
| BSC6 | 2.6±2.3 | 55.8±0.5 | 468.5±3.0 |
| BSC7 | 26.5±1.9 | 3.8±0.5 | 291.6±1.2 |
| BSC8 | 8.6±0.2 | 20.9±0.8 | 378.2±0.9 |
| BSC9 | 7.2±0.5 | 16.3±0.1 | 270.7±3.7 |
| BSC10 | 8.8±2.0 | 70.3±0.9 | 292.5±5.0 |
| GKC1 | 1.5±0.2 | 32.6±0.7 | 416.9±4.4 |
| GKC2 | 4.9±0.5 | 51.8±1.0 | 207.1±0.8 |
| GKC3 | 5.9±1.6 | 44.4±0.9 | 143.1±1.6 |
| GKC4 | 4.3±2.1 | 2.3±0.7 | 187.7±0.8 |
| GKC5 | 3.6±0.6 | 3.1±0.1 | 216.8±0.2 |
| GKC6 | 4.1±1.5 | 18.8±0.1 | 198.9±0.3 |
| GKC7 | 2.8±0.2 | 7.9±0.8 | 259.1±3.7 |
| GKC8 | 9.6±1.3 | 12.7±0.3 | 220.5±0.2 |
| GKC9 | 1.3±1.0 | 21.1±0.3 | 216.8±0.2 |
| GKC10 | 6.3±0.7 | 57.8±0.8 | 177.9±0.5 |
| COE1 | 10.9±2.2 | 30.1±0.5 | 115.2±0.3 |
| COE2 | 13.7±1.5 | 13.6±0.2 | 108.7±0.2 |
| COE3 | 11.9±1.7 | 47.0±1.7 | 137.5±4.8 |
| COE4 | 9.0±1.7 | 66.8±0.7 | 207.6±3.4 |
| COE5 | 2.3±2.9 | 40.7±0.3 | 169.0±0.9 |
| COE6 | 2.4±0.4 | 8.7±0.6 | 43.1±3.1 |
| COE7 | 16.8±1.3 | 52.8±0.8 | 61.6±1.6 |
| COE8 | 22.8±1.9 | 49.4±0.7 | 19.6±2.2 |
| COE9 | 2.8±1.6 | 33.3±1.3 | 241.3±1.1 |
| COE10 | 3.5±0.7 | 2.2±0.3 | 64.4±1.1 |
| RANGE | 1.0±2.3 -26.5±1.9 | 2.2±0.3-70.3±0.9 | 43.1±3.1-468.5±3.0 |
| MEAN | 7.8±1.3 | 29.4±0.9 | 229.4±1.8 |

Table 2: Comparison of activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K in Bq/kg, obtained in present study with that reported for other countries

| Region | ^{226}Ra | ^{232}Th | ^{40}K | Reference |
|------------------------|-------------------|-------------------|-----------------|----------------------------------|
| India (Punjab) | 19.7 | 220.5 | 920.2 | Mehra <i>et al.</i> , (2010) |
| Thailand | 68 | 45 | 213 | Prasong and Susaira (2008) |
| India (Tamilnadu) | 42.9 | 43.9 | 96.0 | Saravanan <i>et al.</i> , (2003) |
| Nigeria (Delta region) | 16.2 | 24.4 | 34.8 | Arogunjo <i>et al.</i> , (2004) |
| Vietnam (south-East) | 19.6 | 31 | 34.6 | Huy and Luyen (2006) |
| Madagascar | 29 | 95 | 294 | Naivo <i>et al.</i> , (2008) |
| Botswana | 34.8 | 41.8 | 432.7 | Murty and Karunakara (2008) |
| Turkey | 115 | 1192 | 1207 | Merdanoglu and Altinsoy (2006) |
| China | 112 | 71.5 | 672 | Yang <i>et al.</i> , (2005) |
| Jordan | 84 | 82 | 560 | Al-jundi <i>et al.</i> , (2003) |
| World average | 33 | 36 | 474 | UNSCEAR (2000) |
| Nigeria (Minna) | 7.8 | 29.4 | 229.4 | Present study |

UNSCEAR (1988) considered bone marrow and the bone surface cells as the organs of interest. Therefore, the annual gonadal dose equivalent (AGDE) in μSvyr^{-1} due to the activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K was calculated using the equation (Mamont-Ciesla *et al.*, 1982; Ibrahim and Mohammad, 2009)

$$\text{AGDE } (\mu\text{Svyr}^{-1}) = 3.09C_{\text{Ra}} + 4.18C_{\text{Th}} + 0.314C_{\text{K}} \quad (5)$$

Where C_{Ra} , C_{Th} , and C_{K} are activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K in Bqkg^{-1} respectively. The results calculated for AGDE for all the samples investigated are presented in Table 3.

As seen in Table 3, AGDE values vary between 40.1 and 410.1 μSvyr^{-1} with an average value of 217.8 μSvyr^{-1} . Generally as seen in Table 3, the mean values of D, ADE, H_{ex} and AGDE are all below the permissible safety limits which show that the hazardous effects of these radiations in the three campuses investigated are negligible.

Table 3: Radium equivalent activity (R_{eq}), gamma dose (D), effective dose equivalent rate (E), external hazard index (H_{ex}), and annual gonadal dose equivalent (AGDE) in the soil samples

| SAMPLE ID | Ra(eq) (Bq/kg) | D (nGy/h) | E (mSv/y) | H(ex) | AGDE (μSvyr^{-1}) |
|-----------|-------------------|--------------|--------------|---------|-----------------------------------|
| BSC1 | 42.5 | 21.2 | 0.03 | 0.1 | 156.1 |
| BSC2 | 72.4 | 34.2 | 0.04 | 0.2 | 240.4 |
| BSC3 | 98.7 | 46.2 | 0.06 | 0.3 | 325.9 |
| BSC4 | 63.7 | 31.0 | 0.04 | 0.2 | 223.1 |
| BSC5 | 66.4 | 31.0 | 0.04 | 0.2 | 218.3 |
| BSC6 | 118.5 | 55.1 | 0.07 | 0.3 | 386.2 |
| BSC7 | 54.4 | 26.6 | 0.03 | 0.1 | 189.2 |
| BSC8 | 67.6 | 32.5 | 0.04 | 0.2 | 231.9 |
| BSC9 | 51.4 | 24.5 | 0.03 | 0.1 | 174.7 |
| BSC10 | 131.9 | 59.7 | 0.07 | 0.4 | 410.1 |
| GKC1 | 80.2 | 38.0 | 0.05 | 0.2 | 270.5 |
| GKC2 | 94.9 | 42.9 | 0.05 | 0.3 | 294.6 |
| GKC3 | 80.4 | 36.2 | 0.04 | 0.2 | 247.0 |
| GKC4 | 22.0 | 11.1 | 0.01 | 0.1 | 81.7 |
| GKC5 | 24.7 | 12.5 | 0.02 | 0.1 | 92.0 |
| GKC6 | 46.3 | 21.7 | 0.03 | 0.1 | 153.0 |
| GKC7 | 34.0 | 16.8 | 0.02 | 0.1 | 122.7 |
| GKC8 | 44.7 | 21.4 | 0.03 | 0.1 | 151.5 |
| GKC9 | 48.2 | 22.6 | 0.03 | 0.1 | 159.4 |
| GKC10 | 102.7 | 46.1 | 0.06 | 0.3 | 314.6 |
| COE1 | 62.8 | 28.5 | 0.03 | 0.2 | 194.5 |
| COE2 | 41.5 | 19.2 | 0.02 | 0.1 | 132.8 |
| COE3 | 89.7 | 40.3 | 0.05 | 0.2 | 274.5 |
| COE4 | 120.5 | 54.2 | 0.07 | 0.3 | 369.5 |
| COE5 | 73.5 | 33.3 | 0.04 | 0.2 | 228.7 |
| COE6 | 18.2 | 8.3 | 0.01 | 0.0 | 57.0 |
| COE7 | 97.0 | 43.1 | 0.05 | 0.3 | 289.8 |
| COE8 | 95.0 | 42.0 | 0.05 | 0.3 | 281.1 |
| COE9 | 69.0 | 31.9 | 0.04 | 0.2 | 222.3 |
| COE10 | 11.6 | 5.6 | 0.01 | 0.0 | 40.1 |
| RANGE | 11.6-131.9 | 5.6-59.7 | 0.01-0.07 | 0.0-0.4 | 40.1-410.1 |
| MEAN | 67.5 | 31.3 | 0.04 | 0.2 | 217.8 |

CONCLUSIONS

This study has analyzed the activity concentrations of terrestrial gamma emitters for soil samples of the three campuses of the two tertiary institutions in Minna, Niger State. The results of this survey provide a pioneering data on the radioactivity levels of the campuses. The obtained results show that the mean values of the radiation hazard parameters calculated for all the soil samples being investigated are lower than the acceptable values for radiation safety provided by UNSCEAR (2000). The calculated dose rates and external hazard indices show that none of the three campuses have any high radiation exposure for either the students or the general public. This means that the radiation hazard is insignificant in the three campuses and so the students and the entire public can freely live and interact on daily basis without any significant radiation threat to the entire population.

REFERENCES

- Alatise, O.O., Babalola, I.A. and Olowofela, J.A. (2008). Distribution of some natural gamma emitting radionuclides in the soils of coastal areas of Nigeria. *Journal of Environmental Radioactivity*, 99: 1746-1749.
- Al-Jundi, J., Al-Bataina, B.A., Abu-Rukah, Y. and Shedadeh, H.M. (2003). Natural radioactivity concentrations in soil samples along the Amman Aqaba highway, Jordan. *Radiation Measurements*, 36: 555-560.
- Arogunjo, A.M., Farai, I.P. and Fowape, A. (2004). Dose rate assessment of terrestrial gamma radiation in the Delta region of Nigeria. *Radiation Protection Dosimetry*, 108: 73-77.
- Beretka, J. and Matthew, P.J. (1985). Natural radioactivity of Australian building materials, industrial wastes and by-products. *Health Physics*, 48 (1):87-95.
- Christa E. Pereira, Vaidyan, V.K., Sunil, A., Ben Byju,S., Reeba Maria Jose, and Jojo, P.J. (2011). Radiological assessment of cement and clay based building materials from southern coastal region of Kerala. *Indian Journal of Pure and Applied Physics*, vol 49: pp 372-376.
- El-Aydarous, A. (2007). Gamma radioactivity levels and their corresponding external exposure of some soil samples from Taif Governorate, Saudi Arabia. *Global Journal of Environmental Research*, 1 (2): 49-53.
- Huy, N.Q. and Luyen, T.V. (2006). Study of external exposure doses from terrestrial radioactivity in southern Vietnam. *Radiation Protection Dosimetry*, 118: 331-336
- Ibrahim, F.A. and Mohammad I. A. (2009). Soil radioactivity levels and radiation hazard assessment in the highlands of northern Jordan. *Radiation Measurements*,44: 102-110.
- Kannan, V., Rajan, M.P., Iyengar, M.A. and Ramesh, R. (2002). Distribution of natural and anthropogenic radionuclides in soil and beach sand samples of Kalpakkam using hyper pure Germanium (HPGe) gamma ray spectrometry. *Applied Radiation and Isotopes*, 57, 109-119.
- Karahan, G and A. Bayulken, (2000). Assessment of gamma dose rates around Istanbul. *Journal of Environmental Radioactivity*, 47: 213-221.
- Malanca, A., Ressina, V. & Dallara, G. (1993). Assessment of the natural radioactivity in soils around the coast of Rio Grande Do Norte. *Health Physics*, 64.
- Mamont-Ciesla, K., Gwaizdowski, B., Biernacka, M. and Zak, A. (1982). Radioactivity of building materials in Poland. In: Vohra, G., Pillai, K. C., Sadarisan, S. (Eds). *Natural radiation environment Halsted Press, New York*, p 551.
- Mehra, R., Badhan, K., Sonkawade, R.G., Kansal, S. and Singh, S. (2010). Analysis of terrestrial natural radionuclides in soil samples and assessment of average effective dose. *Indian Journal of Pure and Applied Physics*, vol 48, pp 805-808.
- Merdonoglu, B. and Altinsoy, N. (2006). Radioactivity concentrations and dose assessments for soil samples from Kestanbal granite area, Turkey. *Radiation Protection Dosimetry*, 121 (4): 399-405.

Murty, V.R.K., and Karunakara, N. (2008). Natural radioactivity in the soil samples of Botswana. *Radiation Measurements*, 43: 1541-1545.

Naivo, R., Martin, R., Frank, T., Asivelo, F.S., Raelina Andriambololona, (2008). Top soil radioactivity assessment in a high natural background area: the case of Vinaninkarena, Antsirabe Madagascar. *Applied Radiation and Isotopes*, 66: 1619-1622.

OECD,(1979). Organization for Economic Cooperation and Development. Exposure to radiation from natural radioactivity in building materials. Report by a group of experts of OECD nuclear energy agency, France.

Prasong, K. and Susaira A. (2008). Natural radioactivity measurements in soil samples collected from municipal area of Hat Yai District in Songkhla province, Thailand. *KMITL Sci. J. vol 8 no 2:52-58*.

Saravanan, S., Jodha, A.S., Deepak, G., Bhati, S.S. and Kumar, S. (2003). Preliminary measurements of natural radioactivity at Madarai District of Tamilnadu, India. *Radiation Measurements*, 36: 397-399.

Sroor, A., El-Bahi, S.M., F. Ahmed, F. and Abdel-Haleem, A.S. (2001). Natural radioactivity and radon exhalation rate of soil in southern Egypt. *Applied Radiation and Isotopes*, 55: 873-879.

UNSCEAR, (2000). Sources and effects of ionizing radiation. Report to the general assembly, with scientific annexes. United Nations, New York.

Yang Ya-Xin, Xin-Min Wu-zhong-Ying, Jiang Wei-Xing, Wang Ji-gen, Lu Jun Lin, Lei-Ming Wang, Yuan-fu Hsia, (2005). Radioactivity concentrations in soils of Xiaz-hung granite area, China. *Journal of Applied Radiation and Isotopes*, 63:255-259.

Yu, K.N., Guan, Z.J., stokes, M.J. and Young, E.C.M. (1992). The assessment of the natural radiation dose committed to the Hong Kong people. *Journal of Environmental Radioactivity*, 17 (1): 31-48.

ACKNOWLEDGEMENT

The authors wish to thank Prof. (Mrs) I. G. E. Ibeanu and the entire staff of Centre for Energy Research and Training (CERT), Ahmadu Bello University Zaria, for their kind assistance in the analysis of the samples.

Received for Publication: 18/09/2012

Accepted for Publication: 05/11/2012

Correspondence Author

Kolo M. T.

Department of Physics, Federal University of Technology, PMB 65, Minna, Niger State, Nigeria.

Email: mat_479@yahoo.com