



## **Malaria Prevalence and Haemoglobin Level Assessment among Patients Attending a Healthcare Facility in North Central Nigeria**

**Chidiebere A. Otuu<sup>1\*</sup>, Rose N. N. Obiezue<sup>1</sup>, Innocent C. J. Omalu<sup>2</sup>,  
Ada Q. A. Otuu<sup>3,4</sup>, Samuel S. Eke<sup>5</sup>, Emmanuel O. Udeh<sup>6</sup>, Innocent C. Ekuma<sup>7</sup>,  
Suleiman C. Hassan<sup>8</sup> and Hadijah U. Yamman<sup>2</sup>**

<sup>1</sup> Parasitology and Public Health Research Unit, Department of Zoology and Environmental Biology, University of Nigeria, Nsukka, Nigeria.

<sup>2</sup> Department of Animal Biology, Federal University of Technology, Minna, Niger State, Nigeria. <sup>3</sup> Department of Public Health, West African Postgraduate College of Pharmacists, Yaba, Lagos State, Nigeria. <sup>4</sup> Department of Pharmacy, Alex Ekwueme Federal University Teaching Hospital, Abakaliki, Ebonyi State, Nigeria. <sup>5</sup>

Department of Biology, Air Force Institute of Technology, Kaduna, Kaduna State, Nigeria. <sup>6</sup> Centre for Integrated Health Programs, Wuse 2, Federal Capital Territory, Abuja, Nigeria.

<sup>7</sup> Department of Biomedical Engineering, Alex Ekwueme Federal Teaching Hospital, Abakaliki, Ebonyi State, Nigeria.

<sup>8</sup> Department of Zoology, Nasarawa State University, Keffi, Nigeria.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors CAO, AQAQ, RNNO and ICJO designed the study and managed the analyses of the study. Authors SSE and EOU performed the statistical analysis, wrote the protocol and the first draft of the manuscript. Authors ICE, SCH and HUY managed the literature searches. All authors read and approved the final manuscript.*

### **Article Information**

#### Editor(s):

(1) Dr. Somdet Srichairatanakool, Chiang Mai University, Thailand.

#### Reviewers:

(1) Muhammad Torequl Islam, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Bangladesh.

(2) Ali Fattahi Bafghi, Shahid Sadoughi University Yazd, Iran. Complete Peer review History: <http://www.sdiarticle4.com/review-history/62670>

**Original Research Article**

**Received 02 September 2020**

**Accepted 07 November 2020**

**Published 30 November 2020**

\*Corresponding author: E-mail: otuuc@yahoo.com;

; Article no.

## ABSTRACT

**Aim:** This study was carried out to assess malaria prevalence and haemoglobin (Hb) level among patients attending a healthcare facility in Abuja, North Central Nigeria.

**Methodology:** Study Design: Malaria parasite infection among the patients was determined by microscopy while the Hb levels was determined using the Mission Haemoglobin test meter. A total of 550 patients who came for consultation at the outpatient clinic of the hospital and whose written informed consents were obtained were randomly recruited for the study. This comprised of 288 adult males and 262 non-pregnant adult females.

**Place and Duration of Study:** The study was carried out at the Outpatient Clinic of Gwarinpa General Hospital, a healthcare facility in Abuja, North Central Nigeria. The study was conducted from March 2019 to May 2019.

**Results:** The overall malaria prevalence was 60.6%. The prevalence was significantly different both by age and gender ( $P < 0.05$ ). In adult males the malaria prevalence was 64.2% while it was 56.5% in non-pregnant adult females. The mean Hb level among malaria positive patients was 10.3g/dL while it was 12.5 g/dL among malaria negative patients. The mean Hb level in malaria positive adult males was 9.7 g/dL while it was 13.8 g/dL in malaria negative adult males. Also the mean Hb level in malaria positive non-pregnant females was 9.2 g/dL while it was 12.3 g/dL in malaria negative nonpregnant females. Pearson's correlation analysis showed that there was a significant difference at the 0.05 level.

**Conclusion:** It was concluded from the results of this study that malaria parasite infection if not confirmed by malaria test and treated promptly may lower haemoglobin levels and lead to anaemia in affected patients.

**Keywords:** Malaria; infection; plasmodium; prevalence; haemoglobin; anaemia.

## 1. INTRODUCTION

Malaria is a fatal insect-borne tropical disease that continues to pose public health challenges globally and is a major cause of mortality, especially in areas where it is endemic

[1,2,3,4,5]. Malaria is caused by five species of *Plasmodium* with *Plasmodium falciparum* being the major cause of malaria cases which still poses a threat to the lives of millions in the tropics [6]. *Plasmodium falciparum* accounted for about 99.7% malaria cases in the World Health Organization African region in 2018 [5]. It remains a major public health problem and in Nigeria it is estimated that about two-thirds of the more than 60 million of its population are constantly exposed to malaria [1]. Haemoglobin is the part of red

blood cells responsible for carrying oxygen in the body. The level of haemoglobin in the blood is very important as levels below normal is an indication that a person has low Red Blood Cell check which may lead to iron deficiency anemia and may cause serious health challenges to the patient. Polycythemia on the other hand is a condition where the Red Blood Cell check is higher than the normal level and also has its health implications [7,8].

It has been reported that malaria is a major cause of anemia in tropical countries where the disease is endemic [8]. The prevalence of anaemia due to malaria parasite infection is dependent on the intensity of transmission which is determined by various factors including the biting habits of the prevalent mosquito vectors [9]. Blood stage malaria parasites are responsible for malarial

disease that might cause anaemia and other health challenges [7]. During the blood stage of the malaria parasite development, malaria trophozoites digest haemoglobin which affects the haemoglobin level in the blood leading to anemia [10]. Recent studies have reported an increasing incidence of resistance of the malaria parasite to currently used antimalarial drugs [11,12,13,4]. According to a recent study, the resistance of malaria to current malaria therapy causes recurrence of infections which has been reported to increase the prevalence and severity of anaemia caused by malaria [8].

## 2. MATERIALS AND METHODS

### 2.1 Study Area

This study was conducted at the General Outpatient Clinic of Gwarimpa General Hospital located a little north-west of Abuja, the Federal Capital of Nigeria. It has a latitude of 9.067318°N and a longitude of 7.39826°E. It lies on 492 m above sea level with an average annual temperature of 25.7°C and rainfall of 1378 mm. Most of the rain is experienced in the month of September with an average of 282 mm. December is the driest month having about 1mm of rain. April is the warmest month with an average temperature of 28.5°C while August with an average temperature of 23.9°C has the lowest temperature of the year.

### 2.2 Study Population and Duration

Adult male and non-pregnant female patients attending the outpatient clinic of Gwarimpa General Hospital Abuja, Nigeria, who gave their written informed consents were enrolled in the study. The criteria for inclusion in the study were fever or a history of fever in the 24 to 48 hours preceding presentation. Patients who had ingested antimalarial drugs in the 24 to 72 hours preceding presentation were excluded from the study. A total of 550 patients whose written consents were obtained participated in this study. This comprised of 288 adult males and 262 nonpregnant adult females. The study was conducted from the months of March to May 2019.

## 2.3 Sample Collection and Laboratory Analysis

### 2.3.1 Collection of blood samples

For the blood samples collection, patients who had taken or ingested antimalarial drugs 24 to 72 hours before the research were excluded from those selected for the study. This was done in order to avoid the effect of such drugs on the result of the malaria tests of the samples. Collection of blood samples to be used for the study was by venipuncture.

### 2.3.2 Preparation of blood films

This was carried out following the methods described by Cheesbrough 2006 [14]. For the thin film, a small drop of blood was placed on the

; Article no.

center of a grease free microscope slide and spread using a slide spreader to produce a thin film which was fixed with methanol for 2 minutes. For the thick film, a larger drop of the blood sample was placed on a grease free microscope slide and then spread with a plastic bulb pipette to produce a thick smear, both of the films were allowed to air dry and then stained with 10% Giemsa stain and allowed to stand for 10 minutes. The stain was washed away and the slides were placed facing up in a draining rack to dry.

## 2.4 Statistical Analysis

Data from the results were statistically analyzed using SPSS (version 20.0). Comparisons between populations, malaria infection and anaemia were made using Chi-Square analysis with an alpha value of  $P < 0.05$  denoting a statistically significant difference. Relationship between age and haemoglobin level was analyzed using Pearson's correlation.

## 3. RESULTS

A total of 550 patients whose consents were obtained participated in this study. This comprised of 288 adult males and 262 nonpregnant adult females. The overall malaria

prevalence was 60.55% (Fig. 1) and was significantly different both by age and gender ( $p < 0.05$ ). The number and percentage of malaria positive and negative patients in the different age groups and the prevalence of malaria among the patients is shown in Table 1. In adult males the malaria prevalence was 64.2% while it was 56.5% in non-pregnant adult females. Malaria positive and malaria negative patients

according to gender are shown in Table 2. Percentage malaria positive and malaria negative patients according to gender are shown in Fig. 2. The percentage Malaria positive and malaria negative patients according to gender are shown in Fig. 2.

**Table 1. Malaria positive and negative patients according to age**

Age	No. Examined (%)	No. Positive (%)	No. Negative (%)
18-27	118 (21.45)	79 (27.72)	39 (17.97)
28-37	117 (21.27)	72 (21.62)	45 (20.74)
38-47	106 (19.27)	63 (18.92)	43 (19.82)
48-57	107 (19.46)	61 (18.32)	46 (21.19)
58-67	102 (18.55)	58 (17.42)	44 (20.28)
Total	550 (100)	333 (60.55)	217 (39.45)


$\chi^2$  Cal = 3.27;  $\chi^2$  tab = 9.49; df = 1

; Article no.


**Table 2. Malaria positive and negative patients according to gender**

Gender	No. Examined (%)	No. Positive (%)	No. Negative (%)
Male	288 (52.36)	185 (55.56)	103 (47.47)
Female	262 (47.64)	148 (44.44)	114 (66.36)
Total	550 (100)	333 (60.55)	217 (39.45)

$\chi^2$  Cal = 3.48;  $\chi^2$  tab = 3.84; df = 1



**Fig. 1. Percentage malaria positive and malaria negative patients and malaria prevalence**



**Fig. 2. Percentage malaria positive and negative patients according to gender**

The mean Hb level among malaria positive patients was 10.3 g/dL while it was 12.5 g/dL among malaria negative patients. The mean Hb levels in malaria positive and negative patients in the different age groups are shown in Table 3. The mean Hb level in malaria positive males was 9.7 g/dL while it was 13.8 g/dL in malaria negative males. Also the mean Hb level in malaria positive females was 9.2 g/dL while it was 12.3 g/dL in malaria negative females. The mean Hb level in malaria positive and negative patients according to gender are shown in Table 4. Percentage mean

Hb level in malaria positive and negative patients according to gender are shown in Fig. 3. The relationship between age and haemoglobin level was determined using Pearson’s correlation. There was a significant difference between age and haemoglobin level at

; Article no.

0.05 level (2-tailed) as shown in Table 5. Percentage mean haemoglobin level in malaria positive and negative patients according to gender are shown in Fig. 3.


Age	No. Positive	No. Negative	Total (Mean Hb level in g/dL)	No. (Mean Hb level in g/dL)
18-27		79 (9.6)	39 (13.8)	118 (23.4)
28-37		72 (11.8)	45 (13.2)	117 (25.0)
38-47		63 (11.6)	43 (12.8)	106 (24.4)
48-57		61 (9.1)	46 (12.5)	107 (21.6)
		58 (9.5)		102 (19.6)
		333 (51.6)		550 (114.0)
58-67			44 (10.1)	
Total			217 (62.4)	

$\chi^2$  Cal = 0.43;  $\chi^2$  tab = 9.49; df = 4

**Table 4. Mean Hb levels in malaria positive and negative patients according to gender**

Gender	No. Positive (Mean Hb Levels g/dL)	No. Negative (Mean Hb Levels g/dL)	Total (Mean Hb Levels g/dL)
Male	185 (9.7)	103 (13.8)	288 (23.5)
Female	148 (9.2)	114 (12.3)	262 (21.5)
Total	333 (18.9)	217 (26.1)	550 (45.0)

$\chi^2_{Cal} = 0.01; \chi^2_{tab} = 3.84; df = 1$



**Fig. 3. Percentage mean Hb level in malaria positive and negative patients according to gender Table 5. Pearson’s correlation for Age vs Hb level of the patients**

Correlations		
	AGE	HB
	1	-.934*
Pearson Correlation		.020
AGE		5
Sig. (2-tailed)	5	
N		1
	-.934*	1
Pearson Correlation	.020	
HB		5
Sig. (2-tailed)	5	
N		5

\*. Correlation is significant at the 0.05 level (2-tailed).

#### 4. DISCUSSION

The prevalence of 60.6% obtained in this study was higher than that earlier reported by Noland et al. 2014 in Abia and Plateau States of Nigeria (36.1% and 36.6% respectively) [15]. It was also higher than the prevalence reported by Otuu et al. 2019 (54.31%) among persons seeking treatment from private drug retailers in North Central Nigeria [4]. Nnamonu et al. 2020 reported a lower prevalence of 43.3% in rice farm settlements, South East Nigeria [16]. Jenkins et al. 2015 reported a lower prevalence of 28.1% in adult patients in malaria endemic area of Kisumu County, Kenya [17]. A lower prevalence of 22% was reported by Njuguna et al. 2016 among febrile patients seeking clinical care at an outpatient health facility in Nairobi, Kenya [18]. Eke et al. 2018 reported a higher prevalence of 69.19% among patients attending General Hospital Minna, North Central Nigeria [3]. A prevalence of 65.5% which was higher than that of this study was reported in a cross-sectional study in a semi-rural Nigerian Medical Centre in Jigawa State by Michael et al. 2019 [19]. Sam Wobo et al. 2014 also reported a higher prevalence of 71.1% in Primary Health Facilities attendees in Ogun State, South Western Nigeria [20]. The prevalence in this is similar to that reported by Dawaki et al. 2016 (60.6%) in Hausa communities in Kano, North Western Nigeria [21]. The prevalence of malaria in this study was higher in male participants (64.2%) than the female participants (56.5%). This agrees with the studies of Jenkins et al. 2015 with 24.1% malaria prevalence in males compared to 32.3% in females [16] and Sakzabre et al. 2020 with prevalence of 30.93% in males as compared to 69.07% in females [22].

The World Health Organization classification of Haemoglobin levels was used to determine whether the haemoglobin level of each patient in the study was within the normal range or not. The World Health Organization classifies normal Hb levels as 13 g/dL in adult males and 12 g/dL in non-pregnant adult females. In this study, haemoglobin levels below 13 g/dl in adult men and below 12 g/dl in non-pregnant adult women were regarded as being below the normal threshold levels. The mean Hb level in malaria positive males was 9.7 g/dL while it was 13.8 g/dL in malaria negative males. Also the mean Hb level

in malaria positive females was 9.2 g/dL while it was 12.3 g/dL in malaria negative females. This result indicates lower Hb levels in malaria positive patients compared to malaria negative patients which may be due to the destruction of red blood cells by the malaria parasite in malaria positive patients. This agrees with the results of Ajibola et al. 2012, Moore, et al. 2008 and Noland et al. 2014 who reported destruction of red blood cells and lowered haemoglobin levels in malaria positive patients [23,10,15] and Sakzabre et al. 2020 who reported anaemia as one of the haematological abnormalities in malaria infected patients [22].

According to Mockenhaupt et al. 1999 [24], acute malaria attacks, chronic and repeated *Plasmodium falciparum* infections are some of the factors that lead to varying degrees of anaemia in malaria endemic regions of the world. They also reported an association between Plasmodium falciparum infection and low haemoglobin levels which they considered as the most significant manifestation of malaria apart from parasitaemia. In a study by Kotepui et al 2014 [25] on the effect of malaria infection on haematological parameters in populations near Thailand-Myanmar, haemoglobin was among the parameters that was significantly low in malaria infected-patients. In a research by Lombardo et al 2017 [26], the results suggested that if an association between haemoglobin and malaria exists, this would be in favour of increased risk of malaria and low haemoglobin levels. Osaro et al 2019 [27] in their study on some haematological parameters of pregnant women in Sokoto, North Western Nigeria, also reported a significant decrease in haemoglobin levels in malaria infected patients compared to non-infected ones.

#### 5. CONCLUSION

In conclusion, the relationship between malaria parasite infection and haemoglobin level in adult male and non-pregnant adult female subjects was investigated. There was a statistically significant difference between malaria infection and haemoglobin level with subjects positive for malaria infection having low haemoglobin levels compared to malaria negative subjects. Pearson's correlation analysis also revealed a significant difference between age and haemoglobin level.

From the results of this study, it was concluded that malaria parasite infection if untreated may lower haemoglobin levels and lead to anaemia in affected patients with the resultant negative health outcomes.

## CONSENT

As per international standard or university standard written patient consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

Ethical approval and permission for this study was obtained from the Research and Ethics Committee (REC) of the Gwarimpa General Hospital, Abuja, Nigeria with REC Approval Number 11/02/2019-30/06/2019.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. World Health Organization (WHO). World malaria Report. World Health Organization Geneva; 2014. Accessed August 20, 2020. Available:[https://www.who.int/malaria/publications/world\\_malaria\\_report\\_2014/en/](https://www.who.int/malaria/publications/world_malaria_report_2014/en/)
2. Omalu ICJ, Olayemi IK, Otuu CA, Eke SS, Uzoaga GO. Entomological and parasitological indices of malaria transmission in Tungan-Goro and Gbaiko communities in Minna, Niger State, Nigeria. *The Zoologist*. 2015;13:1-5.
3. Eke SS, Omalu ICJ, Olayemi IK, Egwim EC, Hassan SC, Otuu CA, et al. Malaria parasitaemia among patients attending general hospital Minna, North Central Nigeria. *Journal of Bioscience and Biotechnology Discovery*. 2018;3(11):7882.
4. Otuu CA, Eke SS, Omalu IC, Hassan SC, Udeh EO, Urama AC. Prevalence of malaria infection among persons seeking treatment from private drug retailers in North Central Nigeria. *Journal of Public Health and Diseases*. 2019;2(1):1-6.
5. World Health Organization (WHO). World malaria Report. World Health Organization Geneva; 2019. Accessed August 20, 2020. Available:<https://www.who.int/publications/item/world-malaria-report-2019>
6. Omalu ICJ, Olayemi IK, Otuu CA, Hassan SC, Eke SS, Paul S, et al. Entomological and parasitological indices of malaria transmission in Minna, Niger State, Nigeria. *Advances in Research*. 2015;3(2): 181-188.
7. Greenwood B, Mutabingwa T. Malaria in 2002. *Nature*. 2002;415(6872):670-672.
8. White NJ. Anaemia and malaria. *Malaria Journal*. 2018;17:371.
9. Omalu ICJ, Eke SS, Olayemi IK, Egwim EC, Yamman HU, Ocha IM, et al. Diversities of *Anopheles gambiae* ss and *Plasmodium falciparum* in Minna, Nigeria. *Nigerian Journal of Parasitology*. 2019; 40(2):135-140.
10. Moore LR, Fujioka H, Williams PS, Chalmers JJ, Grimberg B, Zimmerman P, et al. Haemoglobin degradation in malaria-infected erythrocytes determined from live cell magnetophoresis. *FASEB Journal*. 2008;20(6):747-749.
11. Dondorp AM, Nostern F, Yi P, Das D., Phyto AP, Tarning J, et al. Artemisinin resistance in *Plasmodium falciparum* malaria. *New England Journal of Medicine*. 2009;361: 455-467.
12. Miotto O, Almagro-Garcia J, Manske M, Macinnis B, Champion S, Rockett KA, et al. Multiple population of artemisinin-resistant *Plasmodium falciparum* in Cambodia. *Nature Genetics*. 2013;45(6):648-665.
13. Ajayi NA, Ukwaja KN. Possible artemisinin-based combination therapy resistant malaria in Nigeria: A report of three cases. *Revista da sociedade Brasileira de Medicina Tropical*. 2013;46(4):525-527.
14. Cheesbrough M. Parasitological tests. District laboratory practice in tropical countries, Part 1. University Press, Cambridge; 239-258.
15. Noland GS, Graves PM, Sallau A, Eigege A, Emukah E, Patterson AE. et al. Malaria prevalence, anaemia and baseline intervention coverage prior to mass net distribution in Abia and Plateau States,



- Nigeria. BMC Infectious Diseases. 2014; 14:168.
16. Nnamonu EI, Ani OC, Ugwu FJ, Egba SI, Aguzie IO, Okeke OP, et al. Malaria prevalence in rice farm settlements South East Nigeria. International Journal of Tropical Disease and Health. 2020; 41(9):64-74.
  17. Jenkins R, Omollo R, Ongecha M, Sifuna P, Othieno C, Onger L, et al. Prevalence of malaria parasite in adults and its determinants in malaria endemic area of Kisumu County, Kenya. Malaria Journal. 2015;14:263.
  18. Njuguna HN, Montgomery JM, Cosmos L, Wamola N, Oundo JO, Desai M, et al. Malaria parasitaemia among febrile patients seeking clinical care at an outpatient health facility in an urban informal settlement area in Nairobi, Kenya. American Journal of Tropical Medicine and Hygiene. 2016;94:122-127.
  19. Michael GC, Aliyu I, Idris U, Ibrahim H, Olalere OS, Grema BA, et al. Investigation of malaria by microscopy among febrile outpatients of a semi-rural Nigerian medical center: What happened to malaria control programs? Nigerian Journal of General Practice, 2019;17:23-30.
  20. Sam Wobo SO, Adekunle NO, Adeleke MA, Dedeke GA, Oke OA, Abimbola WA, et al. Epidemiological factors in prevalence of malaria parasites in primary health facilities attendees, Ogun State, Nigeria. Malaria Chemotherapy Control and Elimination. 2014;3:111.
  21. Dawaki S, Al-Mekhlafi HM, Ithoi I, Ibrahim J, Atroosh WM, Abdulsalam AM, et al. Is Nigeria winning the fight against malaria? Prevalence, risk factors and KAP assessment among Hausa communities in Kano State. Malaria Journal. 2016;15:351.
  22. Sakzabre D, Asiamah EA, Akorsu EE, Abaka-Yawson A, Dika, ND, Kwasi DA, et al. Haematological profile of adults with malaria parasitaemia visiting the Volta Regional Hospital, Ghana. Advances in Hematology; 2020. Article ID 9369758.
  23. Ajibola M, Oloruntoba AC, Yetusoko ND. Evaluation of Pcv and haemoglobin variations among malaria positive and malaria negative patients attending the ECWA community health centre, Bukuru, Jos, Nigeria. IOSR Journal of Pharmacy. 2012;2(6):65-69.
  24. Mockenhaupt FP, Bienzle U, May J, Falusi AG, Ademowo OG, Olumese PE, et al. *Plasmodium falciparum* infections: Influence on hemoglobin levels in thalassemia and microcytosis. The Journal of Infectious Diseases. 1999;180:925-928.
  25. Kotepui M, Phunphuech B, Phiwklam N, Chupeerach C, Duangmano S. Effect of malaria infection on haematological parameters in population near ThailandMyanmar border. Malaria Journal. 2014; 13:218.
  26. Lombardo P, Vaucher P, Rarau P, Mueller I, Favrat B, Senn N. Hemoglobin levels and the risk of malaria in papua new guinean infants: A nested cohort study. American Journal of Tropical Medicine and Hygiene. 2017;97(6):1770-1776.
  27. Osaro E, Abdulrahman AE, Erhabor T. Effects of malaria parasitaemia on some haematological parameters of pregnant women of African descent in specialist hospital Sokoto, North Western Nigeria, JOJ Nurse Health Care. 2019;10(4): 555795.

© 2020 Otuu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sdiarticle4.com/review-history/62670>

