

VOLUME 5, NUMBER 1

# JOURNAL OF INFORMATION, EDUCATION, SCIENCE AND TECHNOLOGY (JIEST)



**JUNE, 2018**

**Journal of Information, Education, Science and Technology**

**ISSN: 2360-8846**

## TABLE OF CONTENTS

1. Groundwater Potential Evaluation using Simple Regression Analysis at Gidan Kwano Campus Phase II, Minna, North central Nigeria. **Jonah, S.A., Majekodunmi, S.E., Nmadu, E.N., Suleiman, A.O., Muhammad, J.D., & Adamu, I.B.** 1
2. Sustainable Development of The Nigerian Economy through Social Entrepreneurship Skill Acquisition by Business Education Graduates. **Magnus, P. Udo & Faruk, A. Suleiman** 13
3. Head or Tail: Do Parents' Socio-Economic Status Influence the Performance of Entry Level Senior Secondary Students in Economics. **Oyetoro, Oyebode Stephen Oyetoro, Oyebanke Veronica & Adegboye Abidenu Cornelius** 21
4. Impact of Digital Graphics on Secondary School Biology Students' Retention on the Concept of Pollution in Agaie Metropolis of Niger State. **Koroka, M. U. S. Yakubu, S. E., Osebor, E., Aliyu, N. M., Duru, P. T. & Yusuf J. Amina** 33
5. ICT Access and Integration Into the English Language Classroom. **Chike-Okoli, Chibuogwu Felicia & Ahmed, Muhammed Sadik** 40
6. Entrepreneurship Education as a Mechanism for Employment through Technical Vocational Education & Training (TVET). **Umar, B. Kudu, Musa, Abdulrahaman Ewugi, & Salawu, Chata James** 47
7. Statistical Modeling of Unemployment Duration in Ilorin, Nigeria. **Bushirat T. Bolarinwa** 53
8. Relationship Between Provision of Information Resources and Use of Departmental Libraries In Universities in Nigeria. **Abdulganiy, Okanla Ahmed** 60
9. Availability, Currency and Extent of Information Resources Utilization by Students in Tertiary Institution Libraries in Maiduguri Metropolis, Borno State. **Ibrahim Vandi & Fatima L. Ibrahim** 69
10. Adoption and Use of ICT Tools For Effective Service Delivery in Federal University of Technology, Minna, Library. **Chuks-Ibe, Prisca Oluchi** 80
11. Characteristics of Annual Rainfall Over Guinea Savanna Zone, Nigeria. **Audu, E.B. Abubakar, A.S; Ojoye, S; Muhammed, M. & Mohammed, S.Y.** 87
12. Trend in Heavy Rainfall Over the Guinea Savanna Zone, Nigeria. **Audu, E.B., Abubakar, A. S., Ojoye, S.' Muhammed, M. & Abenu, A.** 95
13. Relationship Between Counsellors' Emotional Intelligence, Study Habits and School Attendance of Students in Federal Unity Colleges in North-Central Nigeria. **Hanior, Ezekiel Aondoascer, Obida Joseph Audu & Igbo, Happiness Ihuoma** 103

## CHARACTERISTICS OF ANNUAL RAINFALL OVER GUINEA SAVANNA ZONE, NIGERIA

Audu, E. B<sup>1</sup>; Abubakar, A. S<sup>2</sup>; Ojoye, S<sup>2</sup>; Muhammed, M<sup>2</sup>. & Mohammed, S.Y.<sup>2</sup>

<sup>1</sup>Government Secondary School, Abuja@30, Pegi, Federal Capital Territory, Nigeria

<sup>2</sup>Department of Geography, Federal University of Technology, Minna, Niger State, Nigeria.

Email: audu\_ebamaiyi@yahoo.com Mobile: +234-803-585-6619

### Abstract

This research was aimed at studying the characteristics of annual rainfall over Guinea Savanna Zone, Nigeria. The data used were monthly rainfall (mm) obtained from the Nigerian Meteorological Agency, Oshodi, Lagos, 1981-2015 (35 years) for Makurdi, Lokoja, Ilorin, Jos, Lafia, Minna and Kaduna; Abuja, 1983-2015 (33 years) and Ibi, 1981-2013 (33 years). Long term rainfall mean, multiple mean comparison, simple linear regression, regression coefficient, maximum, minimum, standard deviation (SD), skewness, kurtosis and cumulative of variation (CV) were used for analysis. Results were presented in table and figures. The conclusion revealed moderate inter-annual rainfall variation, alternate upward and downward trends as well as a general upward trend in recent years. It was also discovered that inter-rainfall characteristics vary across the data collection points even though they are within the same climatic zone. Recommendations focused on the need for similar studies to be carried out to cover other characteristics of rainfall over the study area and in other ecological regions of Nigeria. Short-term weather forecast was also advocated.

**Key words:** Rainfall, variability, global warming, water resources, weather forecast

### 1. Introduction

Weather affects man in almost all his daily activities. In every part of the world, the weather patterns have determined the traditional patterns of food, clothing, housing, agriculture, transportation and social festivals (Asnani, 2005). In Nigeria, rainfall seems to be the most important climate variable because of its role in human activities. The most important human activity in Nigeria remains agriculture which is still largely rain-fed due to inadequate technology for mechanised farming. Rain is also a major source of water for domestic uses and recharging of underground water. According to Eke (2017), rainfall is without doubt the single and most critical physical climatic variable influencing human activities in West Africa.

Rainfall is unevenly distributed across Nigeria as it varies over climatic zones and meteorological stations in onset, cessation, duration, number of rain days, daily, weekly, monthly, annual, pentad and decadal distribution; frequency, magnitude as well as type. Over the Guinea Savanna Zone, Nigeria (GSZN), rainy season generally starts in April and ends in October with changing patterns inform of early/late onset, early/late cessation, normal/abnormal duration (shorter or longer than normal) as well as normal/abnormal accumulated daily, weekly, monthly, and annual. According to Dada (2016), rainfall patterns have changed over the years as a result of global warming and climate change.

Upward and downward trends in both annual and inter-annual rainfall are witnessed in Nigeria and it is an important feature of rainfall. The study of Iornongo (2016) observed an increasing trend of rainfall in most years over Gboko, Benue State. Also, Okoro (2017) observed an increasing trend in rainfall over Bida, Niger State between 2007-2016. The alternate upward and downward trends in rainfall over Nigeria are becoming a major concern to both government and public. Increasing rainfall trend results mostly into flooding as witnessed in September, 2018 in states like Kogi, Niger, Benue and Kwara. According to Iornongo (2016), the inter-annual variation of rainfall mostly in northern Nigeria is much and result into climate and weather hazards especially floods. Previous studies on GSZN regarding rainfall focused on individual data collection points without focusing on the entire GSZN (Mohammed, 2010; Dada, 2016; Iornongo, 2016; Eke, 2017; Okoro, 2017). Hence, this research is aimed at studying the characteristics of annual rainfall over the entire Guinea Savanna Zone, Nigeria (GSZN).

### 2. The Study Area

The Guinea Savanna Zone, Nigeria (GSZN) is centrally located in Nigeria. It is located between longitudes 4°-10°E and latitudes 6°-11°30'N (Figure 1). The study area is characterized by two (2) major

prevailing winds namely; Tropical Maritime Airmass (mT) and Tropical Continental Airmass (cT) (Adakayi, 2000 cited in Ama, 2017). The alternate occurrence of these prevailing winds leads to the occurrence of two (2) distinct seasons which also occur alternately. These seasons are wet (rainy) and dry seasons. Rainfall is moderate with 60% falling in July, August and September. Rainfall occurs in association of squall lines, lightening, thunderstorms and strong winds (Ama, 2017; Omasoro, 2017). The highest mean temperature is recorded in March. Dry season is experienced between October-April (Mohammed, 2010).

The relief of the study area is made up of both high and lowlands (Maxlock, 1980 cited in Omasoro, 2017; Olayinka, 2017). It is predominantly underlain by Precambrian gneisses, granite and schists of crystalline basement complex. Its soils comprise mostly sand, silt, clay and laterite (Iornongo, 2016; Omozuapo, 2016; Oshin, 2008 cited in Ama, 2017; Omasoro, 2017).

The vegetation of the study area is savanna (Physical Setting, Niger State cited in Okesola, 2016). This vegetation has been severely altered by man through numerous activities such as bush burning, farming, firewood harvesting, mining/excavations, constructions and settlement.




Figure 1: The Study Area

Source: National Space Research and Development Agency (NASRDA) (2018)

**Materials and methods**

Secondary data on monthly rainfall (mm) which were in numerical form for Makurdi, Lokoja, Ilorin, Lafia, Minna Jos and Kaduna, 1981-2015; Abuja, 1983-2015 and Ibi, 1981-2013 were used for this study. These data were sourced from the Nigerian Meteorological Agency, Oshodi, Lagos. The data points are evenly distributed across the GSZN.

The annual rainfall (mm) was calculated thus:

Let  $y$  = years,  $x$  = monthly rainfall (mm)

$$y_1 = y_{1,1} + y_{1,2} + \dots + y_{1,3} + \dots + y_{1,12}$$

$$y_2 = y_{2,1} + y_{2,2} + y_{2,3} + \dots + y_{2,12}$$

$$y_3 = y_{3,1} + y_{3,2} + y_{3,3} + \dots + y_{3,12}$$

$$y_{35} = y_{35,1} + y_{35,2} + y_{35,3} + \dots + y_{35,12}$$

$$y_i = \sum_{j=1}^n x_{ji}$$

where  $j=1,2,\dots,35; i=1,2,\dots$  number of months in a year

Simple linear regression analysis was used to ascertain the relationship between the independent and dependent variables per year. It was calculated thus:

$$y = mx + c$$

Where:  $y$  = dependent variable,  $x$  = independent variable,  $c$  = intercept of the trend on  $y$  axis,  $m$  = slope.  
 To specify how much of the variation in the dependent variable is characterized by a variation in the independent variable  $x$ , R-square ( $R^2$ ) was used and calculated as thus:

$$R^2 = 1 - \frac{SSE_{reg\ line}}{SSE_{mean\ y}} \quad 3$$

$$SSE_{reg\ line} = \sum_{i=1}^n (y_i - (mx_i + b))^2$$

$$SSE_{mean\ y} = \sum_{i=1}^n (y_i - \bar{y})^2$$

Where: SSE = sum of square error, regline = regression line  
 The annual rainfall mean was calculated after Ekeruo *et al* (1989) as thus:

$$\bar{x} = \frac{\sum x}{N} \quad 4$$

Where:  $\bar{x}$  = mean,  $\sum x$  = sum of rainfall,  $N$  = number of years

Multiple mean comparison was used to determine the difference in terms of mean annual rainfall across the data collection points.

The skewness coefficient was used to determine the measure of deviation of the data from symmetry of the distribution. It was calculated after Brown (2016) as thus:

$$g_1 = m_3 / m_2^{3/2} \quad 5$$

Where:  $m_3 = \frac{1}{n} \sum (x - \bar{x})^3$  and  $m_2 = \frac{1}{n} \sum (x - \bar{x})^2$

To determine the tailedness of the probability distribution of a real-valued random rainfall, kurtosis was used and calculated after Brown (2016) as thus:

$$a_1 = \frac{m_4}{m_2^2} \text{ and excess kurtosis} = g_2 = a_1 - 3 \quad 6$$

Where:  $m_4 = \frac{1}{n} \sum (x - \bar{x})^4$  and  $m_2 = \frac{1}{n} \sum (x - \bar{x})^2$

In equations 7 and 8,  $n$  = sample size,  $m_3$  and  $m_4$  = 3rd and 4th moments of data set,  $m_2$  = variance

To show how spread out the data values are, standard deviation ( $\sigma$ ) was used and calculated as thus:

$$\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2} \quad 7$$

Where:  $\mu$  = mean,  $x$  = total annual rainfall,  $\sum$  = sum,  $n$  = number of data set (years)

Cumulative of variation was calculated after Umar (2010) to determine the degree of variation in rainfall over the years as thus:

$$CV = \frac{SD}{\bar{x}} \times 100\% \quad 8$$



Where: CV = cumulative of variation, SD = standard deviation,  $\bar{x}$  = mean

## Results and Discussion



Figs. 2-11. Show the annual rainfall (mm) over data collection points in the study area. The study area enjoys moderate annual rainfall as well as rainfall variability. Rainfall variability over the study area shows diverse spacio and temporal (space and time) patterns of occurrence due to differences in latitudinal locations and other localized factors among which are relief and vegetation. The highest rainfall of about 2456.90 mm in the study area was recorded at Ilorin in 2014, while the lowest rainfall of about 697.1 mm was also recorded at Ilorin in 2002. The data collection points are experiencing alternate upward and downward trends. Results also showed that in recent time, Makurdi, Lokoja, Ilorin, Jos, Lafia, Minna and Kaduna are experiencing upward trend in annual rainfall hence, positive linear trend while Abuja and Ibi are experiencing downward trend. Nigerian Meteorological Agency (NiMet) (2017) confirmed increase in annual rainfall over Nigeria from 2006 to 2017. The alternate upward and downward trends are due to the effects of global warming, climate variability and climate change (Audu *et al*, 2012).

The annual rainfall of below normal (less than mean) recorded by some meteorological stations over the study area in 1980s are attributable to drought (Ishiaku *et al*, 2018) which occurred in Nigeria mostly in the Sudano-Sahelian and Guinea Ecological Zones in the late 1970s and extended to 1980s (Adejuwon and Jegede, 2011; Ojoye, 2013; Adeogun *et al*, 2016). Rainfall of below normal is also evidence of dry spells and droughts over the study area.



In addition, rainfall of above normal (above the mean) (table 1) in all the data collection points is the direct effect of global warming occasioned mostly by the socio-economic activities of man among which are gas flaring (Audu, 2013), fossil fuel consumption, farming, over grazing, lumbering, urbanization, construction and industrialization. Adakayi (2015) observed an increase in maximum temperature over Northern Nigeria. NiMet (2017) confirmed increase in temperature across Nigeria in recent years.



**Figure 2:** Annual rainfall over Makurdi, 1981–2015 **Figure 3:** Annual rainfall over Lokoja, 1981–2015  
**Source:** Authors' computation, 2018 **Source:** Authors' computation, 2018



**Figure 4:** Annual rainfall over Ilorin, 1981–2015 **Figure 5:** Annual rainfall over Jos, 1981–2015  
**Source:** Authors' computation, 2018 **Source:** Authors' computation, 2018



**Figure 6:** Annual rainfall over Lafia, 1981–2015 **Figure 7:** Annual rainfall over Abuja, 1983–2015  
**Source:** Authors' computation, 2018 **Source:** Authors' computation, 2018





Figure 8: Annual rainfall over Minna, 1981-2015  
Source: Authors' computation, 2018

Figure 9: Annual rainfall over Kaduna, 1981-2015  
Source: Authors' computation, 2018





Figure 10: Annual rainfall over Ibi, 1981-2013  
Source: Authors' computation, 2018

Figure 11: Mean annual rainfall over GSZN, 1981-2015  
Source: Authors' computation, 2018

The coefficient of multiple determinations ( $R^2$ ) for the study area is positive (Figure 11). However, the values vary among data collection points (Figures 2-9). The implication of the  $R^2$  is that positive changes are occurring in the inter-annual rainfall across the study area. On the contrary, Ibi is experiencing no change (Figure 10).

According to Figure 11, lowest mean rainfall in the GSZN (study area) was recorded in 1983 while the highest mean was recorded in 2014. The zone is experiencing moderate rainfall variability, while the range is about 446.96 mm. The linear equation is positive supporting the upward trend in rainfall over the region. Ibrahim *et al* (2018), observed an insignificant upward trend in rainfall over some meteorological stations in Savanna region, Nigeria.

Table 1 shows the simple statistical analysis of annual rainfall over the study area. The mean in the study area is above 1000 mm and it is within the range of 1000 mm–1200 mm except for Abuja which is above 1400 mm. Rainfall around the mean is regarded as normal, while rainfall below or above the mean is regarded as abnormal. Mean rainfall for the entire study area (GSZN) based on this study is moderate about 1220.43 mm.

All the data collection points have high ranges signifying great inter-annual variations of rainfall. The high SD also confirms the great size in deviations in inter-annual rainfall over the area. The coefficient of variation (CV) for the data points is high. This by implication means that variation in inter-annual rainfall is great. It is greater over Ilorin with the highest value of 24% and lowest over Jos with a value of 10%. The CV for Ilorin is the highest partly due to the astronomical increase in annual rainfall in recent years especially 2014. The kurtosis and skewness are not close to zero (0) hence depicting that inter-annual rainfall variation is great over the study area.

**Table 1: Simple statistical analysis of annual rainfall (mm) over the data collection points**

Data point / statistics	Makur di	Lokoja	Ibi	Ilorin	Lafia	Abuja	Minna	Jos	Kaduna
Mean(mm)	1173.50	1213.22	1061.92	1233.0	1249.92	1421.20	1191.69	1237.09	1202.35
Range(mm)	855.60	962.60	850.60	1759.8	777.20	923.50	724.80	768	865.50
Min(mm)	761.50	804.50	718.50	697.10	839.90	1088.20	818.40	814.70	793.4
year	(2003)	(1982)	(2003)	(2001)	(1988)	(2015)	(1987)	(1995)	(2008)
Max(mm)	1617.10	1767.10	1569.1	2466.6	1617.10	2011.70	1543.20	1582.70	1658.90
year	(1999)	(1999)	(2009)	(2014)	(1999)	(2002)	(2012)	(2002)	(2014)
Skewness	.157	.557	.530	1.874	-.039	.638	-.236	-.261	.438
Kurtosis	-.526	-.134	-.292	7.358	-.248	.925	.022	3.809	.384
SD	212.92	234.68	206.03	302.98	193.59	200.78	163.35	126.77	195.37
CV	18%	19%	19%	24%	15%	14%	13%	10%	16%

**Source:** Authors' computation, 2018

The implications of the results on water resources and agriculture are enormous. When rainfall is above normal, there will be abundant surface and underground water. Excessive surface water leads to soil erosion and flooding as well as high water level in water bodies which also leads to riverine flooding and landslides (as experienced in Kogi State in 1999) as well as high rate of recharging of underground water especially where rainfall is concentrated within few months, weeks and/or days. Above normal rainfall is due to global warming which is causing high rates of evaporation and relative humidity. Below normal rainfall, leads to acute shortage of surface as well as underground water (due to low infiltration) which also affects domestic water supply especially in rural areas with high dependence on rain water harvesting and collection of water from streams, ponds, rivulets, lakes, dams and rivers.

In the area of agriculture, rainfall below the mean has negative effects in form of crop failure. Crop growth and yields are usually affected in form of delayed planting, poor germination, stunted growth, low flowering and low yield. According to Anuforom (2016), soil moisture conditions respond to precipitation anomalies on a relatively short scale; while the ground water, steam flow and reservoir storage reflects the longer-term precipitation anomalies.

### Conclusion and Recommendations

This study is to ascertain the characteristics of annual rainfall over Guinea Savanna Zone, Nigeria (GSZN). The findings revealed moderate inter-annual rainfall, diverse spacio and temporal rainfall variation, alternate upward and downward trends as well as a general upward trend in recent years. The lowest accumulated mean rainfall amount was recorded in 1983 while the highest was in 2014. The range is 446.96 mm. It was also discovered that rainfall characteristics vary across the data collection points even though they are within the same climatic zone. It is therefore recommended that similar studies should be carried out to cover other characteristics of rainfall over the study area and in other ecological regions of Nigeria. Rainfall forecast (very short, short, medium and long-term) should be given more attention with its results disseminated to the grass root so as to serve as early warning against both annual and inter-annual rainfall variations.

### References

- Adakayi, P. E. (2015). Annual Trend of maximum temperature in Northern Nigeria. *Journal of Development and Society*. Faculty of Social Science, University of Abuja, FCT, Nigeria, 3 (1):56-71
- Adeogun, B. K; Ismail, A. & Nwude, M. O. (2016). Comparism of standardized precipitation index and Normalised Diffeence Vegetation Index Anomalies for Drought Analysis. In O. D. Jimoh, M. Y. Otache, .A. Adesiji & M. Saidu (eds). Nigerian Association of Hydrological Sciences. Proceedings of the 7<sup>th</sup> International Conference on Water Resources and Sustainable Development. Pp.71-81.



- Adejuwon, J. O. & Jegede, M.O. (2011). The Impact of Drought on School Environment and Attendance in Rural-Urban setting of Sokoto State. *Journal of Meteorology and Climate Science*. Published by Nigerian Meteorological Society (Nmets), 9 (1),1-7.
- Anuforom, A. C. (2016). Preface. NiMet Drought and Flood Monitoring Bulletin. Monthly hydromet analysis of Standardised Precipitation Index. Published by Nigerian Meteorological Agency (NiMet). 6,1-39.
- Ama, C. K. (2017). Impact of weather elements on the occurrence of respiratory infections in Kuje Area Council, Federal Capital Territory, Abuja, Nigeria. Department of Geography, School of Physical Sciences, Federal University of Technology, Minna, Niger State, Nigeria. Pp. 7-11.
- Asnani, G. C. (2005). Tropical Meteorology. Revised Edition. 1:1-2.
- Audu, H.O., Binbol, N.L., Ekanem, E. M., Felix, I., & Bamaiyi, E.A. (2012). Effects of climate change on length of growing season in Uyo, Akwa Ibom State, Nigeria. *International Journal of Applied Research and Technology*. 1(6):289-294.
- Audu, E. B. (2013). Gas Flaring, A Catalyst to Global Warming in Nigeria: A Catalyst to Global Warming in Nigeria. *International Journal of Science and Technology*. 3 (1): 6–10.
- Brown, S. (2016). Measures of shape: Skewness and kurtosis, n.p
- Dada, S. (2016). Assessment of rainfall variability and its impact on Typhoid Fever occurrence in parts of Minna, Niger State, Nigeria. Department of Geography, School of Physical Sciences, Federal University of Technology, Minna, Niger State, Nigeria. Pg. 4.
- Eke, L. (2017). Weather and its on rice production in Wushishi Local Government Area, Niger State, Nigeria. department of Geography, School of Physical Sciences, Federal University of Technology, Minna, Nigeria. pg. 8
- Ekerou, A. I. C; Ikediashi, A. E; Ekwe, A. O. & Nwamuo, P. A. (1989). Essentials of Educational Psychology. Central Books Ltd, Agbor, Bendel State, Nigeria. Pg. 230.
- Iomongo, T. (2016). Impact of some Agro-climatic variables on cassava yield in Gboko Local Government Area, Benue State, Nigeria. Department of Geography, Federal University of Technology, Minna, Niger State, Nigeria. Pp. 2, 10 & 38.
- Ibrahim, I., Emigilati, M.A. Suleiman, Y. M., Ojoye, S., & Yahaya, T. I. (2018). Effectiveness of early warning methodology and standardized precipitation index for drought monitoring over Guinea Savanna Zone, Nigeria. *Journal of Science, Technology, Mathematics and Education (JOSTMED)*. Department of Science Education, Federal University of Technology, Minna, Nigeria, Africa. 14(2):1-8.
- Mohammed, H. (2010). Mean monthly rainfall pattern of the Shiroro (HEP) during the Pre and Post Dam periods. Department of Geography, Federal University of Technology, Minna, Niger State, Nigeria. pg.8
- National Space Research and Development Agency, Abuja (2018). The Study Area.
- Ojoye, S. (2013). Drought trend analysis and implications for water resource management in the Sudano-Sahelian Zone of Nigeria. *Journal of Science, Technology, Mathematics and Education (JOSTMED)*, 9(3):6-16.

- Okesola, M. S. (2016). Drought dynamics and farmers perceptions in Bida Environs, Niger State, Nigeria. Department of Geography, Federal University of Technology, Minna, Niger State, Nigeria. Pp. 5-9.
- Okoro, J. O. (2017). Impact of climate variability on rice production in Bida Environs, Niger State, Nigeria. Department of Geography, School of Physical Sciences, Federal University of Technology, Minna, Niger State, Nigeria. Pg. 33
- Olayinka, J. I. (2017). Changes in weather variables and the impact on population growth in Bosso Local Government Area, Niger State, Nigeria. Department of Geography, School of Physical Sciences, Federal University of Technology, Minna, Niger State, Nigeria. Pp. 5-8.
- Omasoro, A. Y. (2017). Assessment of micro climate effect on human thermal comfort in part of Chanchaga Local Government Area, Niger State. Department of Geography, School of Physical Sciences, Federal University of Technology, Minna, Niger State, Nigeria. Pp. 6-10
- Omozuapo, A. P. (2016). The effectiveness of Urban Development Control Board in managing environmental challenges in Minna, Niger State, Nigeria. Department of Geography, Federal University of Technology, Minna. Pg. 10
- The Nigeria Meteorological Agency (NiMet) (2017). Climate Review Bulletin. Pp. 3 and 10