

## DEVELOPMENT OF A MATHEMATICAL MODEL FOR UPLAND RICE PRODUCTION (A CASE STUDY OF NATIONAL CEREAL RESEARCH INSTITUTE BADDEGI, NIGERIA)

HAKIMI, D<sup>1</sup>. BATAGI, S. A<sup>2</sup>., & SHEHU, M. D.<sup>3</sup>

Department of Mathematics,  
Federal University of Technology, Minna, Nigeria  
Email: hakimi\_shenfu@yahoo.com, alhajibatagishehu@yahoo.com,  
m.shehu@futminna.edu.ng  
Phone No: +234-803-451-3313

### Abstract

*In this paper, mathematical development of two quadratic models was used to obtain minimum, maximum and saddle points of yield response of upland rice production. The results showed that the models were adequate and significant at 5% by using canonical analysis through the methods of least square,  $R^2$  (coefficient of determination) strength,  $R^2$ -adjust, coefficient of variation (CV) and root mean square error (RMSE). The results also indicated that the quadratic effect of irrigation is important during the dry season than nitrogen for optimum yields. It was also established that there was an increase in yield when variance (ANOVA) was used for the data collected from National Cereal Research Institute Baddegi, Niger State. And computation of the data was adequate with  $R^2$  above 60%,  $R^2$ -adjusted above 55% and RMSE was very small.*

**Keywords:** Upland, Nerica Rice, Quadratic Model, Minimum, Maximum and Saddle Points, Yield Response and Canonical Analysis.

### Introduction

Finley (1972), said response surface methodology examines the relationship between several explained variables. And one or other necessary variables polynomial models are examined through the use of factorial experiment or a fractional factorial design. Response surface methodology is an important way of keeping records to help researchers improve products and services.

Cox (1958), said the application of Mathematics to the production of crops has gained dominance since rice production has become a global issue today through irrigational system and fertilizer application; the application of mathematical modeling becomes imperative. It is known that the mathematical examination of the impact on the objective of production at both rural and urban areas requires appropriate tool such as Mathematical modeling.

Friedman, et al (1948) said that mathematical modeling shows the understanding of Mathematics and help in upland rice production to give yield results. Mathematical models are useful examining tools for manipulating and testing theories, accessing quantitative variables, answering constructive questions, accessing sensitivities to changes in parameter values, and examining key parameters from data observed or collected.

Box, et.al (1990) said the closer Mathematics believes are to reality of behaviors, the more difficult the Mathematical analysis, hence the need to simplify our fillings without losing track of the situation or fillings at hand. Thus, the choice of using Mathematical modeling approach in this research work cannot be over stated.

Hakimi (2005), said the most appropriate model depends on the precision or generality

required, the available data, and the time frame in which results are needed. It is therefore, difficult to express definitively which model is "right", though naturally we are interested in developing models that capture the essential features of a system. Ultimately, we are faced with the usefulness of any model.

This paper is aimed at developing a Mathematical model and analysis of a response surface outlook for upland rice production to meet up with the food consumption of people in Nigeria.

### Materials and Methods

In this research work, we intend to know whether there was loss or increase in the yield result of rice production in the direction of application of irrigation system 'I'. Also, we intend to examine the coefficient of determination  $R^2$ ,  $R^2$  - adjusted, root mean square error (rmse) and coefficient of variation (CV) to check the model adequacy. So also the model equation formulated will be used to determine the point of maximum, minimum or a saddle point of the rice production.

### Quadratic Fit Model

To develop the model equation for the farming system of rice production during the dry season of 2013 and 2014, all possible parameters were taken. We considered the irrigation system 'I', nitrogen fertilizer 'N', and seed varieties of rice 'V' to see the improvement of rice production during the dry season of the years mentioned above.

We also considered the interaction of irrigation system and nitrogen as 'IN', irrigation system and variety of rice as 'IV', nitrogen fertilizer and variety of rice as 'NV'.

The data we used for this research work were collected from NCRI Baddegi, Niger State. The data were based on the field trials during the third quarter of the years mentioned above. The treatments that were applied in the course of experimentation comprised of three (3) irrigation intervals (i.e 7, 14 and 21 days), four (4) fertilizer rates (i.e 30, 60 and 120kg  $Nha^{-1}$ ) which were randomly allocated to the main plots (i.e the region of interest), while four (4) Nerica rice varieties (i.e 2, 3, 4 and 14) constituted the sub-plots.

The fitting modeled equation for the three factors is;

$$y = b_0 + b_1x_1 + b_2x_2 + b_3x_3 + b_{11}x_1^2 + b_{22}x_2^2 + b_{33}x_3^2 + b_{12}x_1x_2 + b_{13}x_1x_3 + b_{23}x_2x_3 + \varepsilon \quad (1)$$

Where

$y$  = yield response

$b_i$  = estimated Parameters ( $i = 0, 1, 2$  and  $3$ )

$x_1=I$  = Irrigation system

$x_2=N$  = Nitrogen fertilizer applied

$x_3=V$  = Variety of rice to be produced

$\varepsilon$  = random error (i.e constant)

### Complete Factorial versus Composite Design

For this research work, we used a complete factorial to produce a model of quadratic surface and it was used instead of a composite design.

A composite design has a  $2^k+2k+1$  treatment combination, and when we applied it to our research design, we had one factor at three levels i.e  $2^1+2(1) + 1 = 5$  and  $2^2+2(2) + 1 = 9$  for two factors at four levels. Therefore, the total treatment combinations will be  $5 \times 9 = 45$ . On the

other hand, for a complete factorial design, we had  $3 \times 4^2$  factorial which is  $3 \times 4 \times 4 = 48$  treatment combinations. Combinational treatment for the composite design is reduced compared to the complete factorial design. Hence, we chose a complete factorial design for the following reasons:

- (i) A full factorial approach will enable equal variance in the estimated effects.
- (ii) It will also allow adequate degrees of freedom for error

### Quadratic Surface for More than one Factor

Equation (1) gives a response function of a quadratic fit model for the three factors used in this research work.

$$y = b_0 + b_1x_1 + b_2x_2 + b_3x_3 + b_{11}x_1^2 + b_{22}x_2^2 + b_{33}x_3^2 + b_{12}x_1x_2 + b_{13}x_1x_3 + b_{23}x_2x_3 + \varepsilon$$

#### Step 1:

The surface is obtained by taking the partial derivatives with respect to  $x_1, x_2, x_3$  and setting them equal to zero, we had

$$\frac{\partial y}{\partial x_1} = b_1 + 2b_{11}x_1 + b_{12}x_2 + b_{13}x_3 = 0 \tag{2}$$

$$\frac{\partial y}{\partial x_2} = b_2 + b_{12}x_1 + 2b_{22}x_2 + b_{23}x_3 = 0 \tag{3}$$

$$\frac{\partial y}{\partial x_3} = b_3 + b_{13}x_1 + b_{23}x_2 + 2b_{33}x_3 = 0 \tag{4}$$

The solutions of these equations give the factorial combinations at which  $y$  is a local maximum, minimum or a local stationary value.

### Canonical Analysis

We applied canonical analysis to determine whether the stationary point is a maximum, minimum or saddle point. Contour plots may also be used.

#### Step 2:

Solving equation (2), (3) and (4) to obtain the points of optimum for  $x_1, x_2$  and  $x_3$

$$\left. \begin{aligned} b_1 + 2b_{11}x_1 + b_{12}x_2 + b_{13}x_3 &= 0 \\ b_2 + b_{12}x_1 + 2b_{22}x_2 + b_{23}x_3 &= 0 \\ b_3 + b_{13}x_1 + b_{23}x_2 + 2b_{33}x_3 &= 0 \end{aligned} \right\} \tag{5}$$

$$2b_{11}x_1 + b_{12}x_2 + b_{13}x_3 = -b_1$$

$$b_{12}x_1 + 2b_{22}x_2 + b_{23}x_3 = -b_2$$

$$b_{13}x_1 + b_{23}x_2 + 2b_{33}x_3 = -b_3 \tag{6}$$

We write the above system of equation in Matrix form

$$\begin{pmatrix} 2b_{11} & b_{12} & b_{13} \\ b_{12} & 2b_{22} & b_{23} \\ b_{13} & b_{23} & 2b_{33} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -b_1 \\ -b_2 \\ -b_3 \end{pmatrix} \tag{7}$$

By using Cramer's Rule we obtained the value of  $x_1$ ,  $x_2$  and  $x_3$  as follows:

Let

$$\Delta_1 = \begin{pmatrix} 2b_{11} & b_{12} & b_{13} \\ b_{12} & 2b_{22} & b_{23} \\ b_{13} & b_{23} & 2b_{33} \end{pmatrix} \quad (8)$$

$$\det \Delta_1 = \begin{vmatrix} 2b_{11} & b_{12} & b_{13} \\ b_{12} & 2b_{22} & b_{23} \\ b_{13} & b_{23} & 2b_{33} \end{vmatrix}$$

We solved and arrived

$$x_1 = \frac{(\Delta_2)}{(\Delta_1)} = \frac{\Delta_2}{\det \Delta_1} \quad (9)$$

$$x_1 = \frac{-4b_1b_{22} + b_1b_{23}b_{23} + 2b_2b_{12}b_{33} - b_3b_{12}b_{23} - b_2b_{13}b_{23} + b_3b_{13}b_{22}}{8b_{11}b_{12}b_{33} - 2b_{11}b_{23}b_{23} - 2b_{12}b_{12}b_{33} + b_{12}b_{12} + b_{12}b_{13}b_{23} + 2b_{13}b_{13}b_{22}} \quad (10)$$

$$x_2 = \frac{(\Delta_3)}{(\Delta_1)} = \frac{\Delta_3}{\det \Delta_1} \quad (11)$$

$$x_2 = \frac{-4b_1b_{11}b_{33} + 2b_3b_{11}b_{23} + 2b_1b_{12}b_{33} - b_1b_{13}b_{23} - b_3b_{13}b_{12} + b_2b_{13}b_{13}}{8b_{11}b_{12}b_{33} - 2b_{11}b_{23}b_{23} - 2b_{12}b_{12}b_{33} + b_{12}b_{12} + b_{12}b_{13}b_{23} + 2b_{13}b_{13}b_{22}} \quad (12)$$

$$x_3 = \frac{(\Delta_4)}{(\Delta_1)} = \frac{\Delta_4}{\det \Delta_1} \quad (13)$$

$$x_3 = \frac{-4b_3b_{11}b_{22} + 2b_2b_1b_{23} + b_3b_{12}b_{12} - b_2b_{12}b_{13} - b_1b_{12}b_{23} + 2b_1b_{13}b_{22}}{8b_{11}b_{12}b_{33} - 2b_{11}b_{23}b_{23} - 2b_{12}b_{12}b_{33} + b_{12}b_{12} + b_{12}b_{13}b_{23} + 2b_{13}b_{13}b_{22}} \quad (14)$$

**Step 3:**

To obtain the optimum response  $y_m$  ( $m$  is the optimum) we substitute the values of  $x_1, x_2$  and  $x_3$  in to equation (3) below;

$$y = b_0 + b_1x_1 + b_2x_2 + b_3x_3 + b_{11}x_1^2 + b_{22}x_2^2 + b_{33}x_3^2 + b_{12}x_1b_{13}x_1x_3 + b_{23}x_2x_2x_3 + e \quad (15)$$

**Step 4:**

A determinant matrix is formed to determine the coefficients in the canonical form i.e

$$\begin{pmatrix} 2b_{11} & b_{12} & b_{13} & x_1 \\ b_{12} & 2b_{22} & b_{23} & x_2 \\ b_{13} & b_{23} & 2b_{33} & x_3 \end{pmatrix} \begin{pmatrix} -b_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -b_2 \\ -b_3 \end{pmatrix}$$

i.e

$$\begin{pmatrix} 2b_{11} & b_{12} & b_{13} \\ b_{12} & 2b_{22} & b_{23} \\ b_{13} & b_{23} & 2b_{33} \end{pmatrix} \quad (16)$$

We divided  $R_1, R_2$  and  $R_3$  by 2 and we had

$$\begin{pmatrix} \frac{b_{11}}{2} & \frac{b_{12}}{2} & \frac{b_{13}}{2} \\ \frac{b_{12}}{2} & b_{22} & \frac{b_{23}}{2} \\ \frac{b_{13}}{2} & \frac{b_{23}}{2} & b_{33} \end{pmatrix}$$

We had our characteristics equation as:

$$\begin{pmatrix} (b_{11} - \lambda) & \frac{b_{12}}{2} & \frac{b_{13}}{2} \\ \frac{b_{12}}{2} & (b_{22} - \lambda) & \frac{b_{23}}{2} \\ \frac{b_{13}}{2} & \frac{b_{23}}{2} & (b_{33} - \lambda) \end{pmatrix} = 0 \quad (17)$$

Therefore, we solved and arrived at characteristics equation as:

$$-8\lambda^3 + (8b_{11} + 8b_{22} + 8b_{33})\lambda^2 - (2b_{12}b_{12} + 2b_{13}b_{13} + 2b_{23}b_{23} - 8b_{11}b_{22} - 8b_{11}b_{33} + 8b_{22}b_{23})\lambda - 6b_{11}b_{22}b_{33} + 2b_{11}b_{23}b_{23} - 2b_{12}b_{13}b_{23} + 2b_{13}b_{13}b_{22} = 0 \quad (18)$$

### Step 5

We substituted the values of  $x_1, x_2$  and  $x_3$  by using computer Algebraic symbolic package "Marple software", and subtracted the result from the original equation (i.e equation 3) to have.

$$y - y_m = b_{11}x_1^2 + b_{11}b_{11}^2 + b_{22}x_2^2 + b_{33}x_3^2 \quad (19)$$

### Results

The result obtained in step 5;  $y - y_m = b_{11}x_1^2 + b_{22}x_2^2 + b_{33}x_3^2$  can be interpreted as a change in yield from the point of optimum, m to some points  $((x_1, x_2$  and  $x_3)$ .

The following reasons are observed:

If the coefficients  $b_{11}, b_{12}$  are all negative, then there is a loss in yield whichever way one goes from m.

- i. If  $b_{11}, b_{12}$  are negative and  $b_{33}$  is zero, there is still a loss in yield.
- ii. If  $b_{11}$  is negative and  $b_{22}$  and  $b_{33}$  are both zero, there is no unique centre and we have a stationary ridge.
- iii. If  $b_{11}$  is negative,  $b_{22}$  and  $b_{33}$  are both positive and measure an increase in yield along the  $x_2$  or  $x_3$  axis at infinity, then we have a rising ridge.

### Fitting Model for 2013 Rice Yield

The model that was fitted for rice yield in 2013 is:

$$y = 26.5 + 0.1x_1 - 0.02x_1^2 + 0.54x_2 - 0.01x_2^2 + 0.000013x_2^3 - 0.098x_1x_2 + 0.0014x_1x_2^2 - 0.000004x_1x_2^3 + 0.004x_1^2x_2 - 0.0001x_1^2x_2^2 + 0.00000021x_1^2x_2^3 \quad (20)$$

where,

$y$  = response estimate for yield

$x_1$  = irrigation linear effect

$x_1^2$  = irrigation quadratic effect

$x_2$  = nitrogen linear effect

$x_2^2$  = nitrogen quadratic effect

$x_2^3$  = nitrogen cubic effect

Equation (20) is the response surface polynomial for 2013 rice yield. The ANOVA table below helps us to select the parameters that are significant and needed for the response surface model.

**Table 4.1: ANOVA table for 2013 rice yield**

SOURCE	Degree of Freedom (df)	Sum of Square (SS)	Mean Squares (MS)	F	P-VALUE
REP	1	17.17	17.17	2.10	0.1508
$x_1$	2	742.95	317.47	45.52	0.0001
$x_2$	1	741.84	741.84	90.91	0.0001
$x_1^2$	1	1.11	1.11	0.136	0.7134
$x_2^2$	1	198.15	66.05	8.09	0.0001
$x_1x_2$	3	193.40	193.40	23.70	0.0001
$x_1^2x_2$	1	193.40	4.75	0.58	0.4479
$x_1^2x_2^2$	1	4.75	0.0002	0.000025	0.9957
$x_1^2x_2^3$	1	0.0002	26.10	3.198	
$x_1x_2^2$	6	156.62	38.15	4.67	0.0335
$x_1^2x_2^2$	1	38.15	24.15	2.96	0.0892
$x_1^2x_2^3$	1	24.15	2.51	0.31	0.5805
$x_1^3x_2^2$	1	2.51	85.50	10.47	0.0017
$x_1^3x_2^3$	1	85.50	5.55	0.68	0.4119
ERROR	1	5.55	0.76	0.09	0.7613
TOTAL	83	0.76	677.66	8.16	
	95	1792.54			

Source: (Author, 2016)

From the table above it can be seen that at 5% level of significance the parameters  $x_1x_2$ ,  $x_1^2x_2$  and  $x_1^2x_2^2$  are significant since their p-values are less than  $\alpha = 5\%$  significance level and our final response surface function is;

$$y = 26.5 + 0.25x_1 + 0.54x_2 - 0.098x_1x_2 + 0.004x_1^2x_2 \quad (21)$$

From the ANOVA table above we observed that  $x_1^2$ ,  $x_2^2$  and  $x_1^2x_2^2$  are not significant at 5% level of significant, therefore, we concluded that effects on rice yield are not significant.

### Discussion

Here we will determine whether the stationary point is a point of maximum, minimum or a saddle point.

The fitted model to be used is equation

$$y = 26.5 + 0.25x_1 + 0.54x_2 - 0.098x_1x_2 + 0.004x_1^2x_2 \quad (22)$$

Differentiate equation (4.2) partially w.r.t.  $x_1$  and  $x_2$  and equate to zero to find the optimum point we have;

$$\frac{\partial y}{\partial x_1} = 0.25 - 0.098x_2 + 0.008x_1x_2$$

$$0 = 0.25 - 0.098x_2 + 0.008x_1x_2$$

$$0.098x_2 - 0.008x_1x_2 = 0.25 \quad (23)$$

$$0 = 0.54 - 0.098x_1 + 0.004x_1^2$$

$$0 = 0.54 - 0.098x_1 + 0.004x_1^2$$

$$0.098x_1 - 0.004x_1^2 = 0.54 \quad (24)$$

Solving simultaneously equation (1.23) and (1.24), we have;

$$x_1 = 16.13 \text{ or } 8.37$$

Substituting  $x_1 = 16.13$  and  $8.37$  into equation (1.24) to obtain values for  $x_2$ , we have;  $x_2 = -8.051$  and  $8.054$  when  $x_1 = 16.13$  and  $8.369$  respectively; these are the optimum points, that is,  $x_1 = 16, x_2 = 0$  or  $x_1 = 8, x_2 = 8$ .

We then substitute  $x_1$  and  $x_2$  using  $x_1 = 16, x_2 = 0$  and  $x_1 = 8, x_2 = 8.05$  in equation (1.20) to obtain the optimum response  $Y_m$  as;

$$Y_m = 26.5 + 0.25(16) + 0.54(0) - 0.098(16)(0) - (0.004 \cdot 0)(16)^2$$

$$Y_m = 30.5 \text{ is the optimum response for } x_1 = 16, x_2 = 0$$

$$Y_m = 26.5 \text{ is the optimum response for } x_1 = 8 \text{ and } x_2 = 8$$

We therefore constructed a determinant matrix as follow to determine the coefficient in the canonical form.

$$\begin{vmatrix} b_{11} - \lambda & (b_{12})/2 \\ \frac{b_{12}}{2} & b_{22} - \lambda \end{vmatrix}$$

Determine the Eigen-values or characteristic roots and equate the determinant to zero, we have;

$$\begin{vmatrix} 0 - \lambda & -0.049 \\ -0.049 & 0 - \lambda \end{vmatrix} = 0$$

Solving the determinant matrix we have;

$$-\lambda(-\lambda) - (-0.049 * -0.049) = 0$$

$$\lambda^2 = 0.002401$$

$$\lambda = + 0.049$$

taking the values of  $\lambda$  we have two set of canonical equations as

$$Y - 30.5 = 0.049x_1^2 - 0.049x_2^2 \tag{25}$$

$$Y - 33.3 = 0.049x_1^2 - 0.049x_2^2 \tag{26}$$

Equation (1.25) is obtained when  $x_1 = 16$  and  $x_2 = 0$  while, equation (1.26) is obtained when  $x_1 = 8$  and  $x_2 = 8$ . We observed that the contours of both equation (1.25) and equation (1.26) are saddle surfaces because their coefficients  $b_{11}$  and  $b_{22}$  are positive and negative respectively, which means that there will be a rapid increase in yield if  $x_1$  is increased. This shows that there will be an increase in yield in the direction of  $x_1$  axis from the optimum response M.





Figure1: Normal probability plot of the residuals for 2013 rice yield

Figure 1 above is the normal probability plot for 2013 rice yield data to check for normality of the data and it appears that the shape confirms the normality of the data.



**Fig 2: Residuals versus the order of the data for 2013 rice yield**

Figure 2 above is a graph showing the residuals versus the order of the data for 2013 rice data it is a confirmatory graph of figure 4.2 (residual versus fitted values).

### Conclusion

Basically, a complete factorial experiment where each factor of all the levels are absolutely combined and the model was adequate. We found that the linear and quadratic response surfaces were significant. The model showed that in the optimum response region, there was an increase in yield when irrigation and nitrogen are combined at quadratic and linear effect levels respectively.

### Acknowledgments

We are thankful to Almighty Allah by granting us this very cost opportunity to write this paper and for all the blessings.

Special credits go to Prof. K. R. Adeboye, Prof. N. I. Akinwande and Prof. Y. A. Yahaya. All from Department of Mathematics, Federal University of Technology, Minna, Niger State. Our special gratitude also goes to Dr. M. D. Shehu, Dr. Sirajo Abdulrahman, Dr. A. Ndanusa, Dr. Jiya Mohammed and Dr. Adamu Mohammed.

### References

- Box, e tal (1990). *Experimental designs*. New York: John Wiley and Sons. Inc. pp. 335 - 370.
- Cox, D. R (1958). *Planning of experiments*. New York: John Wiley and sons, Inc. Wiley Classics Library Edition (1992). 120 - 128.
- Finley, D. J. (1972). *Statistical science in agriculture*. Blackwell scientific publications. Oxford, London: Edinburgh and Melbourne 110 - 166.



Friedman, M., & Savage, L. J. (1947). *Planning Experiments seeking maxima in techniques of statistical analysis*. New York: McGraw Hill.

Hakimi, D. (2005). The numerical simulates approaches to the two dimensional equasilinear equipotential fluidics. *Proceedings of the 1<sup>st</sup> Annual Conference of School of Science and Science Education, Federal University of Technology, Minna*, 88 – 94.