



**PROCEEDINGS**



**OF THE**  
**GENETICS SOCIETY OF NIGERIA**  
IN COLLABORATION WITH  
**FEDERAL UNIVERSITY OF AGRICULTURE  
MAKURDI**

**41<sup>ST</sup>**  
**GSN**

**ANNUAL CONFERENCE  
(MAKURDI 2017)**

THEME:  
**"GENETICS & NATIONAL  
ECONOMIC REVIVAL"**

SUNDAY 15TH - THURSDAY 19TH OCTOBER, 2017

EDITED BY  
VANGE T. AND UGBAA M.S.

# **PROCEEDINGS**

of the

# **41<sup>st</sup>**

**National Annual Conference  
of the**

**Genetics Society of Nigeria (GSN)**

**October 15-19, 2017**

Publication of Genetics Society of Nigeria (GSN)

## LOCAL ORGANIZING COMMITTEE AND REVIEWERS

1. Prof. Terkimbi Vange - LOC Chairman
2. Dr. G.O.S. Ojo - LOC Secretary
3. Mr. Macsamuel S. Ugbaa - LOC Assist. Sec.
4. Chairman Technical Sub-Committee-Prof (Mrs) N. I. Odiaka  
Member- Prof. D.K Adedzwa  
Member- Prof. L.L Bello  
Member-Prof. M.O. Adeyemo  
Member-Prof. Momoh O.M
5. Chairman Financial Sub-Committee-Prof. M.I. Ogbaji  
Member- Prof. D.K Adezwa  
Member-Dr. Kortse P.A  
Member-Mrs E.A. Ochigbo  
Member-Prof. E.H. Kwon-Ndung
- 6.Chairman Publicity sub-committee-Dr. Egahi J.O.  
Member-Dr. Ochefu Joshua  
Member-Mr. MacsamuelUgbaa  
Member-Mr. Akombo Richard
- 7.Chairman Program Sub-committee-Prof. L.L Bello  
Member - Prof. Momoh O.M.  
Member - Dr. Moses Ogah  
Member - Mr. Olalekan Olassan  
Member - Mr. Ojobo Omoche
- 8.Chairman Transport Sub-committee-Dr. J. Okoh  
Member-Dr. Ochefu Joshua
- 9.Chairperson Welfare/Entertainment-A.E Ochigbo  
Member-E.N Etukudo  
Member-Prof. Odiaka N.  
Member-Mrs. O. Vange
- 10.Chairman Accommodation Sub-committee-Dr. Gwaza  
Member-Dr. A. A Jimin
- 11.Chairman Venue Sub-committee Dr. Ugese F.D  
PG students (Agronomy)
- 12Chairman Protocol Sub-committee- Dr T.S Msaakpa  
Mr. Christopher Nwadike  
Mr. Titus Yongu

# National Officials Genetics Society of Nigeria (GSN)

**President** - Prof E. H. Kwon-Ndung  
Department of Botany, Faculty of Science, Federal University Lafia  
08036488345 [kwon\\_ndung@yahoo.com](mailto:kwon_ndung@yahoo.com)

**Vice President** - Prof S. A. Olakojo  
Institute of Agric Research and Training I A R&T Obafemi Awolowo University  
Ile Ife Pmb 5029 Moor Plantation Ibadan 08034671714 [saolakojo@yahoo.co.uk](mailto:saolakojo@yahoo.co.uk)

**Secretary General**-Dr D. M. Ogah  
Animal Science Dept, faculty of Agriculture, Nasarawa State University, Keffi.  
Lafia Campus [mosesdogah@yahoo.com](mailto:mosesdogah@yahoo.com) 08065303418

**Assistant Secretary** - Dr J. Kaye  
Animal Science Dept, faculty of Agriculture, University of Jos  
08039354507

**Treasurer** -Dr Mrs Ene Ochigbo  
Department of Plant breeding, College of Agronomy, University of  
Agriculture Makurdi 08160686050 [ochigboene2016@gmail.com](mailto:ochigboene2016@gmail.com)

**Financial Secretary**- Dr Macaulay Ittah  
University of Calabar 08036767013 [macittah@yahoo.com](mailto:macittah@yahoo.com)

**Public Relation Officer**- Dr J. C. Onovo  
Department of Biological Science, Faculty of Natural Science, Nasarawa  
State University keffi 08065709649 [onovojos@yahoo.com](mailto:onovojos@yahoo.com)

**Editor in Chief** - Dr F T Rotimi  
Dept of Animal Production, University of Ilorin  
08060395678 [fayetiro@yahoo.com](mailto:fayetiro@yahoo.com)

**Associate Editor**- Dr Kenneth Omokhafe  
Rubber Research Institute of Nigeria, Benin City  
08078047096 [kenomokhafe2001@yahoo.com](mailto:kenomokhafe2001@yahoo.com)

**Internal Auditor**- Dr O. F. Owolade  
I A R & T Obafemi Awolowo University  
Moor plantation Ibadan 08035478678 [ofowolade@yahoo.com](mailto:ofowolade@yahoo.com)

**Ex-officio** Prof W. Akin-Hassan  
Animal Science Dept, Faculty of Agriculture, Usmanu Danfodio University,  
Sokoto.  
08035875473 [akinola19@yahoo.com](mailto:akinola19@yahoo.com)

# Table of Contents

## SECTION ONE

<b>HERITABILITY AND GENETIC ADVANCE IN WHITE MAIZE (<i>Zea mays</i> L.) INBRED LINES</b>		
K.D. Dawaki, M. Yusuf, A.K Ibrahim, H.S.Hamisu, and M.S. Jibrin	- -	1
<b>TREES GENETIC TRANSFORMATION FOR DROUGHT TOLERANCE: A PROSPECT FOR <i>ACACIA SENEGAL</i> [L.] WILLD</b>		
Fakuta, N. M., Ojiekpon, I.F., Sagir, M., Usman, A., Gashua, I. B., Ogunremi O. C <sup>4</sup> , Jibrin, M.S, Jonah, P.M, Simon, S.Y.	- - - -	6
<b>INHERITANCE OF QUALITATIVE CHARACTERS AND HETEROSIS AMONG THE F<sub>1</sub> AND F<sub>2</sub> HYBRIDS IN RICE (<i>Oryza sativa</i> L. and <i>Oryza glaberrima</i> Steud)</b>		
Chimdi, Grace O, Abu, Ngozi E, Russom, Z and Fagam, S. A.	- -	14
<b>CORRELATION AND PATH ANALYSIS OF QUANTITATIVE TRAITS IN WEST AFRICAN OKRA (<i>Abelmoschuscaillei</i>) GENOTYPES</b>		
Jonah, P.M., Fakuta, N.M., Vange, T. and Iyongo, T.T	- - - -	28
<b>COMPARATIVE YIELD PERFORMANCE OF 47 DIFFERENT ORANGE FLESHED SWEET POTATO (OFSP) GENOTYPES FOR ROOT YIELD, DRY MATTER AND <math>\beta</math>-CAROTENE.</b>		
Harriman, J. C., Okocha, P. I., Nwofia, G. E. and Solomon, A.	- - - -	33
<b>EXPLOITING GRAIN YIELD PERFORMANCE AND STABILITY OF 40 PROMISING RICE ACCESSIONS USING AMMI MODEL</b>		
M. Bashir*, B.O. Ehirim, A. Odoba, A.T. Maji, S. O. Bakare <sup>1</sup> and S.T. Gbadeyan	- -	45
<b>SAFEGUARDING THE NIGERIAN COCOA ECONOMY- MOLECULAR- BASED EVIDENCE OF <i>CACAO SWOLLEN SHOOT VIRUS</i> (CSSV) AND ITS NATURAL VECTORS IN THE NIGERIAN COCOA (<i>Theobroma cacao</i> L.)</b>		
Ekemini Obok, Peter Aikpokpodion, Paul Hadley and Andy Wetten	- -	54
<b>MUTAGENIC EFFECTS OF SODIUM AZIDE ON MORPHOLOGICAL PARAMETERS OF LAGOS SPINACH (<i>Celosia argentea</i> L.)</b>		
Abubakar, A., Falusi, A. O., Daudu, O. A. Y., Dangana, M.C., Yahaya, S.A	- -	61
<b>ASSESSMENT OF GENETIC DIVERSITY IN BAMBARA GROUNDNUT (<i>Vigna subterranea</i> (L.) VERDC) USING MORPHOLOGICAL, PROXIMATE AND PROTEIN PROFILE MARKERS</b>		
Alege, Gbenga Olorunshola, Daisi, Roseline Omolade, Omoleye, Wemimo Atinuke and Isiaka, Franca	- - - - - - - -	70
<b>INFLUENCE OF ELECTRICITY SUPPLY ON STORAGE CONDITIONS AND SUBSEQUENT GERMINATION OF AMARANTH SEEDS (<i>Amaranthus cruentus</i> L.)</b>		
Olosunde A. A and O. A. Ojo	- - - - - - - -	80

**PHYSIOLOGICAL CHARACTERIZATION OF GROUNDNUT SEEDLINGS  
USING VIGOUR AND PERFORMANCE INDICES**

Olalekan J. Olasan, Celestine U. Aguoru, Macsamuel S. Ugbaa and  
Ebenezer J. Ekefan

84

**RELATIONSHIPS BETWEEN NORMALIZED DIFFERENCE VEGETATION  
INDEX (NDVI) AND OTHER TRAITS OF COWPEA HYBRIDS UNDER  
DROUGHT AND WELL-WATERED CONDITIONS**

Ibitoye, D.O., Tongoona, P., Blay, E.T., Offei, S.K., Boukar, O., Olomide, O.A.K.

92

**GENETIC DIVERSITY OF *STRIGA*-RESISTANT MAIZE (*Zea mays* L.)  
INBRED LINES USING DArTseq MARKERS**

S. O. Afolabi, A. Menkir, M. Oyekunle<sup>1</sup>, U. S. Abdullahi

100

**GENE ACTION AND HERITABILITY STUDIES OF YIELD IN COWPEA  
(*VIGNA UNGUICULATA* L. WALP)**

Aliyu D. J., Ogbaji, M.I., and Akombo .R.

108

**BIOFORTIFICATION QUANDARY ASSOCIATED WITH DISTRIBUTION  
PATTERN OF IRON ACCUMULATIONS IN THE THREE MAIN PORTIONS  
OF THE COWPEA (*VIGNA UNGUICULATA* L. WALP) SEEDS**

\*Umar M. <sup>1</sup>, Y. Caro<sup>1</sup>, M.F. Ishiyak<sup>1</sup> and B. S. Mohammed

113

**GENETIC VARIABILITY OF SOME PHENOTYPIC TRAITS OF SOME CASSAVA  
GENOTYPES IN A GENOMIC SELECTION  
TRAINING POPULATION**

Amaefula C<sup>1</sup>, Nnaji C.I. <sup>1</sup>, Ahamefule S. N

119

**PERFORMANCE OF INTRODUCED MAIZE VARIETIES IN MAKURDI,  
SOUTHERN GUINEA SAVANNA OF NIGERIA**

Akaazua, Barnabas Wanger, Yadoo Adzer and Owoyemi Stephen Segun

126

**GRAIN ZINC AND IRON CONCENTRATIONS IN MAIZE TOPCROSS AND  
VARIETY HYBRIDS AND THEIR RELATIONSHIP WITH SEED QUALITY**

Akinwale, R. O., Sulola, B. H., Awosanmi, F. E. and Fadoju, A. O.

132

**GENETIC VARIANCES AND HERITABILITY FOR GRAIN YIELD AND YIELD  
COMPONENTS IN A DIALLEL CROSS OF EARLY MAIZE**

Akinwale, Richard Olutayo, Adewale, Samuel Adeyemi, Fadoju,

Abigail Opeyemi, and John, Stephen, Olurankinse, Olumuyiwa Paul

139

**COMPARATIVE ANALYSIS OF YIELD AND AGRONOMIC PERFORMANCE  
OF MAIZE TOP-CROSS AND VARIETY HYBRIDS**

Fadoju, A.O. and Akinwale R.O.

145

**YIELD POTENTIAL AND TRAITS ASSOCIATION AMONG  
SELECTED NSUKKA-YELLOW PEPPER (*Capsicum annum* L.)  
BREEDING LINES IN SOUTH WESTERN NIGERIA**

Anyaocha Christian Okechukwu\*, Yusuf Rashidat Oyindamola,

Fariyike Timothy Alaba

152

<b>VARIATIONS IN SOME AGRONOMIC TRAITS AMONGST M<sub>2</sub> MUTANTS OF SUNFLOWER (HELIANTHUS ANNUS L.) INDUCED BY ETHYLMETHANE-SULFONATE (EMS)</b>	
Yekini A. E., Usman A., Mansur Y., Katung M. D., Mohammed, S. M.	158
<b>CORRELATION AND FACTOR ANALYSES OF MARKETABLE LEAF YIELD IN FLUTED PUMPKIN (<i>Telfairia occidentalis</i>)</b>	
Fayeun, L. S.	166
<b>FLOWERING, SEXUALITY AND YIELD CHARACTERISTICS ANALYSES OF INDUSTRIAL SUGARCANE IN HUMID TROPICAL AGRO-ECOLOGY</b>	
M. A. Ittah and A. P. Etefia	172
<b>EFFECTS OF GAMMA RADIATION ON GERMINATION, SURVIVAL, GROWTH AND YIELD OF AFRICAN YAM BEAN (<i>SPENOSTYLIS STENOCARPA</i> (HOCHST. EX A. RICH) HARMS)</b>	
Ihuoma, Ogechi and Adesoye, Adenubi, I.	178
<b>GENETIC PROGRESS IN THE EARLY HYBRID (F<sub>1</sub> - F<sub>4</sub>) POPULATIONS ARISING FROM INTERSPECIFIC CROSSES AMONG FOUR SOLANUM VARIETIES</b>	
Atugwu A.I*, Onyia, V.C and Soluobi B.I	188
<b>TRAITS ASSOCIATED WITH SOYBEAN YIELD AND THEIR IMPLICATIONS</b>	
T.Vange; L. A. Ihyembe, A.C. Iorkyaa and P. Swende	208
<b>EVALUATION OF BODY BIOMETRY AND LOSS IN HYBRID VIGOUR AND THEIR MIXED GENETIC GRADES OF THE DUAL PURPOSE FRENCH GUINEA FOWL</b>	
Elkana, H.and Gwaza, D. S.	217
<b>GENOMIC DNA EXTRACTION FROM NIGERIAN DATE PALM (<i>Phoenix dactylifera</i> L.) USING DIFFERENT EXTRACTION METHODS WITHOUT LIQUID NITROGEN</b>	
B. O. Emoghene, O. Asemota, C. R. Eke and M. Idu	227
<b>EFFECTS OF COMBINED COLCHICINE AND GAMMA RAYS TREATMENTS ON THE VARIABILITY OF AGRONOMIC TRAITS IN FONIO (<i>Digitaria exilis</i> [Kippist] Stapf.)</b>	
Nura, S., Adamu, A.K., Adelanwa, M.A., Usman, I.S., and Shehu, K.	234
<b>EVALUATION OF SOME SOYBEAN (<i>Glycine max</i> L. MERRILLS) GENOTYPES USING MORPHO-AGRONOMIC CHARACTERISTICS IN IBADAN, NIGERIA</b>	
Yekeen, O. M., Amoo, M. A., Olomola, A., and Adeyemo, Y. A.	243
<b>IDENTIFICATION OF GENETIC POLYMORPHISM AMONG GROUNDNUT GENOTYPES USING SIMPLE SEQUENCE REPEAT (SSR) MARKERS.</b>	
Dughdugh Paul, Adedzwa K. Daniel, Omoigui O. Lucky, Olasan J. Olalekan, Ugbaa S. Macsamuel	252

<b>IDENTIFYING NEW SOURCES OF APHIDS [APHIS CRACCIVORA (KOCH)] RESISTANCE IN COWPEA GERMPLASM USING PHENOTYPIC AND MOLECULAR MARKER APPROACHES</b>	
G. C. Ekeuro , L. O. Omoigui, L. L. Bello, M. P. Timko, G. O. Ogunwolu, C.C. Danmaigona, M.S Ugbaa, N. Iyorkaa. - - - - -	257
<b>ASSESSMENT OF GENETIC DIVERSITY AMONG SELECTED MAIZE LANDRACES AND HYBRIDS USING SSR MARKERS LINKED TO QTLS FOR DROUGHT AND NITROGEN TOLERANCE</b>	
Oyenike A. Adeyemo, Olatunde. A. Bhadmus, Olumide A. Adebessin, and Uzezi E. Okinedo - - - - -	268
<b>OPTIMIZED PROTOCOL FOR ELUTION OF COWPEA (<i>Vigna unguiculata</i> (L.) Walp.) PLANT DNA ON FTA CARD FOR USE IN SCREENING OF MARKERS IN MAPPING POPULATIONS</b>	
Catherine C. Danmaigona, Lucky O. Omoigui, Michael P. Timko - - - - -	278
<b>BIOFORTIFICATION OF STAPLE FOOD: A PARADIGN SHIFT IN BREEDING CROP FOR BETTER NUTRITION AND IMPROVED FOOD SECURITY IN NIGERIA</b>	
Nwadike, C., Terkimbi Vange., Ojo.G.O.S., Nwosu, D.J and A.U. Okere. - - - - -	285
<b>FAST NEUTRONS INDUCED VARIATION IN FOXTAIL MILLET (<i>SETARIA ITALICA</i> (L.) P. BEAUV.)</b>	
Esson, A.E., Adelanwa, M.A., Adamu, A.K., and Adebola, M.I. - - - - -	296
<b>IDENTIFYING OIL PALM FRUIT FORMS USING RAPD AND AS-PCR MARKERS</b>	
Odenore, V.D., Anoliefo, G.O., Asemota, O., Ikhajiagbe, B., Eke, C.R., Ihase, L.O. and Obahiagbon, O. - - - - -	303
<b>RELATIONSHIPS OF BEAN YIELD TRAITS IN SOME Cacao (<i>Theobroma cacao</i> L.) GENOTYPES</b>	
Adenuga, O.O, Adepoju, A.F., Olaniyi, O.O., Mapayi, E.F. and Idrisu M. - - - - -	307
<b>CORRELATION AND PRINCIPAL COMPONENT ANALYSIS (PCA) OF IMPORTANT TRAITS OF COWPEA SEEDLINGS UNDER DROUGHT STRESS</b>	
Ajayi, A.T., Gbadamosi, A.E., Olumekun, V.O. - - - - -	314
<b>PRELIMINARY EVALUATION OF SOME COWPEA CULTIVARS FOR YIELD AND YIELD PARAMETERS AT LAFIA, NASARAWA STATE, NIGERIA</b>	
H.A Kana, O.P. Ifenkwe and E.E. Goler - - - - -	322
<b>THE PROSPECTS OF EARLY-MATURING CROPS IN ENSURING FOOD SECURITY IN NIGERIA</b>	
Garba Linda Larai and Jury Esther Jonah - - - - -	327
<b>GENETIC RELATIONSHIPS BETWEEN FRESH RHIZOME YIELD AND OTHER QUANTITATIVE CHARACTERS IN GINGER (<i>Zingiber officinale</i> Roscoe)</b>	
Umar, U. U., Musa, N. Mijinyawa, A. <sup>3</sup> , Bashir, Y.R. and Abdulrahman, M.D. - - - - -	331



# ASSESSMENT OF SOME GENETIC DIVERSITY OF SOME COTTON CULTIVARS IN MINNA NIGER STATE, NIGERIA

Dangana, M C , Falusi O A., Daudu, O A Y , Abubakar, A., Gado, A A and Oluwajobi, A.O

<sup>1</sup> Department of Biological Sciences, Federal University of Technology, Minna  
Corresponding Author: danganachata@gmail.com:08035890169

## Abstract

Cotton (*Gossypium* sp.) is an important cash crop of the family Malvaceae which produces lint and seed that serve as food and raw materials for textile and seed crushing industries. Despite, the economic importance of this crop to human livelihood, its total production remains far below the national requirements for textile and oil mills due to obstacle posed by poor seed quality and germination. Pollen quality and pollination play a fundamental role in fertilization and seed sets. Therefore, study on pollen parameter is essential for understanding fertility of this crop and its improvement. A total of six accessions of cotton (SAMCOT 8, 9, 10, 11, 12, and 13) were evaluated for morphological parameters on the field and their pollen studies were carried out in the laboratory at maturity. The accessions were evaluated for plant height, number of branches, days to bract square and boll formations, pollen viability and size (both intine and exine). The results of the morphological parameters showed significant variations in plant height with SAMCOT 008 having the highest plant (49.75cm) and SAMCOT 010 had the least (26.75cm) at the end of the study period. In terms of earliness to maturity SAMCOT 008 matured earliest, with 37 days and 55 days for bract square and boll formation, while SAMCOT 013 had the least maturity status of 56 days and 77 days for both bract square and Boll formation. Significant variations were recorded in the pollen parameters with highest percentage viability recorded in SAMCOT 009 (94.25%) and the lowest in SAMCOT 008 77.78%. Variation in the pollen size (both intine and exine) among taccessions was recorded, with the lowest intine (20.50 $\mu$ m) and exine (27.00 $\mu$ m) in SAMCOT 011. The highest pollen intine was recorded in both SAMCOT 010 & SAMCOT 012 (25.50 $\mu$ m) and exine (33.00) in SAMCOT 008. The highest pollen size, plant height coupled with least days for bract square and boll formation might indicate the presence of genes for earliness in the accessions which is a good trait for selection and improvement in breeding programs.

**Keywords:** cotton, accessions, variations, pollination

## INTRODUCTION

### Background to the study

Cotton (*Gossypium* sp.) is a soft staple fiber belonging to the family Malvaceae. It is one of the most important and earliest domesticated plants in the world. Among the different domesticated species, *G. arboreum*, *G. herbaceum*, *G. hirsutum* and *G. barbadense* are the most common. Cotton plant is a native to tropical and subtropical regions around the world, including, Africa, America and India (Wendel and Cronn, 2003; Rahman, et al., 2012). The greatest diversity of wild cotton species is found in Mexico, followed by Australia and Africa (Cotton Australia, 2016).

It is an important cash crop in Nigeria which produces lint and seed that serve as food and raw materials for the local textiles and seed crushing industries (Adeniji, 2011). The seeds are about 15% of the value of the crop and are pressed to produce edible oil for human

consumption and cotton seed cake for feeding of ruminant animals due to its high protein content. The fiber is pure cellulosic material with unique structure from which paper is made (Adeniji, 2007; Hornyak et al., 2008). Cotton fibers are grown commercially for the seed coat to form a boll of cotton lint. The seed in the "boll", is surrounded by fibers of two types, commercially produced by a process called ginning. At the first ginning, the longer fibers, called staples, are removed and these are twisted together to form yarn for making thread and weaving into high quality textiles. At the second ginning, the shorter fibers, called "Inters", are removed, and these are woven into lower quality textiles (which include the eponymous lint). Commercial species of cotton plant are *G. hirsutum* (>90% of world production), *G. barbadense* (30-45%), *G. arboretum* and *G. herbaceum* (together 2%). Many varieties of cotton have been developed

by selective breeding and hybridization of these species.

Cotton was the fifth most important export source of foreign exchange in Nigeria before now, unfortunately, its total production remains far below the national requirements for consumption and industries. This is as a result of low average yield of the crop on the field below its genetic yield potentials (Ogunlela, 2004). Pollen and pollination have been reported to play a fundamental role in fertilization and consequently seed and fruit sets in spermatophytes (Asma, 2008). Deficiency in pollen production and performance could have direct effects on seed formation, seed viability, seed germinability and fruit production (Raji *et al.*, 2012). Therefore, High pollen quantity and quality profile are central to viable seed formation and propagation in seed plants. However, there are limited studies on the pollen parameter of indigenous cotton species in Nigeria. Therefore, this study was carried out to investigate morphological and some pollen parameters in selected accessions of *Gossypium* sp.

## MATERIALS AND METHODS

### Study area

The study was carried both on the field (Garden) and laboratory of the Department of Biological Sciences, Federal University of Technology, Minna, Niger State.

### Sample Collection

Cotton seed accessions used for the study were collected from institute of Agricultural Research, Ahmadu Bello University Zaria. The accessions were SAMCOT 8, 9, 10, 11, 12, and 13. The samples were packed in a brown envelope and labeled properly as reported by Daudu *et al.* (2015).

### Experimental Design and Planting of Seeds

The seeds were sown in twenty-eight (28) plastic buckets of 20 litres each, filled with rich sandy-loamy soil and arranged in Randomized complete Block Design (RCBD). Four plastic buckets were allocated to each accession.

A total of six seeds of each accession were planted per pot at a depth of 2-3 inches below the soil surface. After two weeks of seed emergence, the less vigorous ones were thinned out and two plants were left per bucket. The plants were monitored for morphological and phenotypic variables at budding, flowering and fruiting stages and using standard procedures as

described by Akinyele and Osekita (2006); and Falus *al.* (2014).

Parameters investigated and recorded include plant height: the distance from ground level up to the terminal bud on main axis of the plant in (cm), Number of branches per plant (both monopodial and sympodial branches), stem, leaf, flower colors, stem hairiness, days to bract square formation and days to boll formation. Pollen grain viability test was carried out and the pollen size was measured using ocular graticle as described by Abubakar *et al.* (2015).

### Pollen Viability and Size

Mature flowers from each accession were harvested on the field and taken to the laboratory for microscopic analysis of the pollen grain. Pollen grains were picked from the anther of each flower using a pair of forceps and dusted on a glass slide. The pollen grains were then mixed with 1ml of water and two (2) drops of Lecto-phenol blue stain were added. The slide was then mounted on a microscope and viewed at X40 magnification. The deeply stained and rounded pollen were taken as viable and fairly, non-stained and distorted shape pollen as non-viable.

Diameters of 30 randomly selected pollens were measured for both intine and exine using the eye piece graticle and recorded in micrometre ( $\mu\text{m}$ ) (Abubakar *et al.*, 2015).

### Result and Discussion

The result from the analysis of variance showed significant difference at ( $P \leq 0.05$ ) among all the accessions for all the traits examined. The plant height for all the weeks of study showed great variations in the morphological appearance of the crop. SAMCOT 010 had the least plant height of 21.10cm at week 4, and SAMCOT 008 (49.75cm) had the highest. Though there was no significant differences at ( $P \geq 0.05$ ) in the plant height among SAMCOT 009 (31.00cm), SAMCOT 011 (31.25cm), and SAMCOT 013 (35.70cm). The morphological differences in plant height is in agreement with the findings of Mahmood *et al.* (2006) and maharam . (2014). They observed significant differences among cotton cultivars for plant height with most rapid growth occurring before flowering.

From the, result there is an indication that the height does not necessarily translate to many branches as recorded in SAMCOT 008 which has few number (2.00) of branches with the highest plant height of (49.75 ). Similar to



this result, Basbag and Gencer (2007) reported few number of branches in some varieties of cotton plant. This low values in number of branches per plant might be an advantageous trait for earliness. Baloch *et al.* (2014) reported that low mean value of branches is an indication of earliness in cotton.

Significant differences in number of branches were recorded among all the genotypes. SAMCOT 010 had the highest mean value (5.00) from week four (4) to week (6) and SAMCOT 013 recorded the highest value (6.00) at the end of week 7. This highest value ( $6.00 \pm 1.00$ ) was not significantly different from the values of SAMCOT 9, 10 and 12 (Table.1). The highest value recorded might be an indication of high yield since every branch carries a number of bolls

The formation of bract square is an indication that a cotton plant has attained a maturity age. In terms of earliness to maturity, SAMCOT 008 matured earliest, with 37 days and 55 days for bract square and boll formation, while SAMCOT 013 had the least maturity status of 56 days and 77 days for both bract square and boll formation from the date of crop emergence. This result is in agreement with the reports of Baloch *et al* (2014), Ahmad *et al.* (2012) and Jatoi *et al.* (2012).

The results of pollen viability and size are presented in Table 2. The results indicated that there was significant variation ( $p \geq 0.05$ ) in the pollen parameters of the cotton accessions. The highest percentage viability was recorded in SAMCOT 011 (93.48%) and the lowest was recorded in Samcot 008 with value of 77.98%. Significant variation in the pollen size (both intine and exine) among the accessions was obtained, with the lowest intine ( $20.50 \mu\text{m}$ ) and exine ( $27.00 \mu\text{m}$ ) in SAMCOT 011. The highest pollen intine was recorded in SAMCOT 012 ( $25.50 \mu\text{m}$ ) and exine ( $33.00$ ) in SAMCOT 008. However, there was no significant different in the values recorded in the intine for SAMCOT 009 (25.50), 10 (25.50) and 12 (25.50), and between the highest exine (33.00) in SAMCOT 008. The variation recorded in the intine and exine of the accessions could be due to aberration in microsporogenesis resulting from high degree of meiotic irregularity. This result is in agreement with the report of Abubakar *et al.* (2015) who recorded variation in the pollen size of *Celosia argentea*.

## CONCLUSION

The present study revealed that SAMCOT 008 with least values for bract square formation and days to boll formation posses the genes for earliness and SAMCOT 013 with high number of branches and with the least days of bract square formation might posses the genes for high yield in Minna which is a trait for selection in the breeding programs.

## REFERENCES

- Abubakar A., Falusi A. O., Daudu O. A. Y., Oluwajobi A. O., Dangana M. C., Abejide D.R. (2015). mutagenic effects of sodium azide and fast neutron irradiation on the cytological parameters of m2 lagos spinach (*celosia argenteavarcrystal*.) *World Journal of Agricultural Research*, (3),3, 107-112.
- Adeniji, O.B. (2011). Adoption of improved Technologies for Cotton Production in Katsina State, Nigeria. *Ph.D Thesis*, submitted to post graduate school. Ahmadu Bello University, Zaria.
- Ahmad, S., Ashraf, M., Khan, N., and Iqbal, (2012) Assessment of yield-related morphological measures for earliness in upland Cotton (*Gossypium hirsutum* L.) *Pak.J.Bot*, 40(3) 1201-1207.
- Akinyele, B. O., and Osekita, O. S. (2006). Correlation and path coefficient analyses of seed yield attributes in Okra (*Abelmoschus esculentus* (L) Moench). *African Journal of Biotechnology*, 5(14), 1330-1336.
- Asma, B. M. (2008). Determination of pollen viability, germination ratios and morphology of eight apricot genotypes. *African Journal of Biotechnology*, 7(23), 4269-4273.
- Baloch, M. J., Khan, N. U., Rajput, M. A., Jatoi, W. A., Gul, S., Rind, H., and veesar, N. F. (2014). Yield related morphological measures of short duration cotton gynotypes. *Journal of Animal and plant Sciences* 24,(4), 1198-1211.
- Basbag, S and Gencer, O. (2007). Investigation of some yield and fibre quality characteristics of interspecific hybrid (*Gossypium hirsutum* L. and *G. barbadense* L) cotton varieties *Hereditas* 144, 33-42.
- Cotton Australia. (2016). *Cotton Plant*. Retrieved: Aug. 12<sup>th</sup> 2016 from <http://cottonaustralia.com.au/cotton-library/fact-sheets/the-cotton-plant>
- Daudu, O.A.Y., Falusi, O.A., Dangana, M.C., Abubakar, A., Yahaya, S.A. and Abejide,



- D.R. (2015). Collection and evaluation of Roselle (*Hibiscus sabdariffa* L.) germplasm in Nigeria. *African Journal of Food Science*, 9(3), 92-96.
- Falusi, O.A., Dangana, M.C., Daudu, O.A.Y., Oluwajobi, A. O., Abejide, D.R., and Abubakar, A. (2014). Evaluation of some Roselle (*Hibiscus sabdariffa* L.) germplasm in Nigeria. *International Journal of Biotechnology and Food Science*, 2(1):117-121.
- Hornyak, G.L., Tibbals, H.F., Dutta, J. and Moore, J.J. (2008). *Introduction to nanoscience and nanotechnology*. CRC Press, First Edition, United States of America, pp. 378-387.
- Jatoi, W.A., Baloch, M.J., Panhwar, A., Veesar, N. F. and Panhwar, S.A. (2012) Characterisation and identification of early maturing up land cotton varieties. *Sarhad journal of Agriculture* 28,1,53-56.
- Mahmood, S., Mezhar Irfan, M., Raheel, F., and Hussain, A. (2006). Characterization of cotton (*Gossypium hirsutum* L.) varieties for growth and productivity traits under water deficit conditions. *International Journal of Agriculture & Biology* 08,(6) 796-800
- Maharam, F.M., Bronson, K.F. & Maas, S.J. (2014). Inter-relationships of cotton plant height, canopy width, ground cover and plant nitrogen status indicators. Publications from USDA ARS / UNL Faculty. Paper 147
4. <http://digitalcommons.unl.edu/usdaarsfacpub/1474>
- Ogunlela, V.B. (2004). Improved technology for increased production of cotton in Nigeria, paper presented at the trainers workshop on presidential initiative on vegetable oils, cotton and groundnut production lake Chad research institute 10-12 May, Maiduguri.
- Rahman, M., Shaheen, T., Tabbasam, N., Iqbal, M. A., Zafar, Y. & Paterson, A. H. (2012). Cotton genetic resources: A review. *Agronomical Sustainability Development*, 32, 419-432.
- Raji, A. A., Nor, H. Z., Nor, H. H. & Mohamad, O. (2012). Studies on Pollen Viability and Germinability in Accessions of *Stevia rebaudiana* Bertoni. *International Journal of Biology*, 4(3), 117-123.
- Wendel, J. F. & Cronn, R. C. (2003). Polyploidy and the evolutionary history of cotton. *Advances in Agronomy*, 78, 139-186.

**Table 1: Plant Height (cm) for the Cotton Accessions at 4<sup>th</sup>, 5<sup>th</sup>, 6<sup>th</sup> and 7<sup>th</sup> week after sowing.**

Treatments	accession	Week 4	Week 5	Week 6	Week 7
Plant Height (cm)	SAMCOT 008	21.60±8.40 <sup>b</sup>	25.25±9.25 <sup>b</sup>	36.35±6.35 <sup>bc</sup>	49.75±3.13 <sup>c</sup>
	SAMCOT 009	22.1±8.10 <sup>b</sup>	28.00±2.00 <sup>b</sup>	31.00±2.10 <sup>b</sup>	35.50±2.05 <sup>b</sup>
	SAMCOT 010	21.10±2.10 <sup>b</sup>	23.40±3.10 <sup>b</sup>	25.65±3.35 <sup>ab</sup>	26.75±1.26 <sup>ab</sup>
	SAMCOT 011	26.50±0.50 <sup>d</sup>	30.15±3.85 <sup>bc</sup>	31.25±4.25 <sup>b</sup>	32.10±1.08 <sup>b</sup>
	SAMCOT 012	24.25±1.25 <sup>c</sup>	42.00±1.50 <sup>c</sup>	42.75±1.25 <sup>c</sup>	43.65±6.20 <sup>c</sup>
	SAMCOT 013	24.90±0.10 <sup>c</sup>	33.00±3.00 <sup>bc</sup>	33.45±3.05 <sup>b</sup>	35.70±0.98 <sup>b</sup>
Number of Branches per plant	SAMCOT 008	2.00±0.01 <sup>b</sup>	2.00±1 <sup>ab</sup>	2.00±1 <sup>a</sup>	2.00±1 <sup>a</sup>
	SAMCOT 009	2.00±1.00 <sup>b</sup>	3.00±1 <sup>b</sup>	4.00±1 <sup>b</sup>	5.00±1 <sup>c</sup>
	SAMCOT 010	4.00±0 <sup>d</sup>	5.00±1	5.00±1 <sup>c</sup>	5.00±1 <sup>bc</sup>
	SAMCOT 011	4.00±0 <sup>d</sup>	2.00±1 <sup>ab</sup>	4.00±1 <sup>b</sup>	4.00±1 <sup>b</sup>
	SAMCOT 012	3.00±1 <sup>c</sup>	3.00±1 <sup>b</sup>	4.00±1 <sup>b</sup>	5.00±1 <sup>bc</sup>
	SAMCOT 013	3.00±0 <sup>c</sup>	4.00±1 <sup>c</sup>	4.00±1 <sup>b</sup>	6.00±1 <sup>c</sup>

Values are Mean ± Standard Error of mean. Values with the same superscript along the column are not significantly different from each other at P ≤ 0.05 according to DMRT.


**Table 2: Percentage of pollen viability and non-viability of some cotton cultivars**

Accessions	% Pollen viability	% Pollen non-viability	Intine( $\mu\text{m}$ )	Exine( $\mu\text{m}$ )
Sam cot 008	77.78 $\pm$ 0.75 <sup>a</sup>	22.22 $\pm$ 0.62 <sup>cd</sup>	24.25 $\pm$ 2.64 <sup>b</sup>	33.00 $\pm$ 1.43 <sup>c</sup>
Sam cot 009	94.25 $\pm$ 0.42 <sup>c</sup>	5.75 $\pm$ 0.64 <sup>a</sup>	25.25 $\pm$ 0.69 <sup>c</sup>	30.50 $\pm$ 0.62 <sup>ab</sup>
Sam cot 010	81.22 $\pm$ 0.45 <sup>b</sup>	18.78 $\pm$ 0.36 <sup>c</sup>	25.50 $\pm$ 0.73 <sup>c</sup>	31.50 $\pm$ 0.55 <sup>b</sup>
Sam cot 011	93.48 $\pm$ 0.96 <sup>c</sup>	6.52 $\pm$ 0.74 <sup>a</sup>	20.50 $\pm$ 1.17 <sup>a</sup>	27.00 $\pm$ 1.11 <sup>a</sup>
Sam cot 012	86.14 $\pm$ 0.34 <sup>b</sup>	13.86 $\pm$ 1.41 <sup>bc</sup>	25.50 $\pm$ 1.17 <sup>c</sup>	31.50 $\pm$ 1.19 <sup>b</sup>
Sam cot 013	90.16 $\pm$ 1.26 <sup>c</sup>	9.84 $\pm$ 0.43 <sup>b</sup>	24.50 $\pm$ 0.82 <sup>b</sup>	30.50 $\pm$ 0.73 <sup>ab</sup>

Values with the same superscript along the column are not significantly different from each other at  $P \leq 0.05$  according to DMRT.

**Table 3. Days to bracts square and boll formation**

Accessions	Days to bracts square formation	Days to boll formation
SAM COT 8	37	55
SAM COT9	40	59
SAMCOT10	45	64
SAMCOT11	48	67
SAMCOT12	53	71
SAMCOT13	56	77


**Figure 1: Phenotypic Features of Selected Cotton Accessions**

A: bract square formation  
C: Fertile pollen and

B: Ball formation  
D: Pollen size